



# **STEELHEAD (ONCORHYNCHUS MYKISS) POPULATION AND HABITAT MONITORING IN LOWER YAKIMA RIVER TRIBUTARIES**

2014

BPA Project # 1996-035-01



**Report covers work performed under BPA contract #(s) 56662 REL 50, 56662 REL 25**

**Report was completed under BPA contract #(s) 56662 REL 50, 56662 REL 64**

1/1/2013 - 12/31/2014

Tim Resseguie, Yakama Confederated Tribes, Toppenish, WA, 98948

Report Created 10-2014

This report was funded by the Bonneville Power Administration (BPA), U.S. Department of Energy, as part of BPA's program to protect, mitigate, and enhance fish and wildlife affected by the development and operation of hydroelectric facilities on the Columbia River and its tributaries. The views in this report are the author's and do not necessarily represent the views of BPA.

This report should be cited as follows: Tim Resseguie, STEELHEAD (ONCORHYNCHUS MYKISS) POPULATION AND HABITAT MONITORING IN LOWER YAKIMA RIVER TRIBUTARIES, 1/1/2013 - 12/31/2014 Annual Report, 1996-035-01

# Table of Contents

Table of Contents .....	2
1. Executive Summary .....	3
Fish Population Monitoring .....	3
Tributary Habitat RME .....	3
2. Introduction .....	3
3. Methods: Protocols, Study Designs, and Study Area .....	8
4. Results .....	13
a. Fish Population Status Monitoring (RM&E) .....	13
b. Tributary Habitat RM&E .....	18
5. Synthesis of Findings: Discussion/Conclusions .....	21
a. Fish Population Status Monitoring (RM&E) .....	21
a. Tibutary Habitat (RM&E) .....	22
References .....	23
Appendix A: Use of Data & Products .....	25
Appendix B: Detailed Results .....	26
Appendix C: List of Metrics and Indicators .....	32

## 1. Executive Summary

To monitor the trends and status of the steelhead populations in Toppenish, Satus and Ahtanum Creeks we conducted spawning ground surveys including a redd count to index adult spawning escapement. We operated rotary screw traps to monitor juvenile outmigration below the spawning and rearing habitat on these watersheds. We also monitored water temperature, an important limiting factor for steelhead production in the Yakima River Basin (Yakima Fish and Wildlife Recovery Board 2004, 2009)

### Fish Population Monitoring

The number of steelhead redds (n=267) increased in Satus Creek in 2014 compared with 223 redds observed the previous year but were lower than peak years of 2010 and 2011 (Figure 5). Redd numbers (n=123) increased in Toppenish Creek and were the highest since 2002 (Figure 7). Good conditions for the identification of steelhead redds were present in both creek and played a role in the increase in observed redd numbers. Spatial distribution of redds appeared to be typical in Satus (Figure 6) and Toppenish Creeks (Figure 8).

The estimated number (n=46,470 se 5008) of juvenile outmigrants from Toppenish Creek (Toppenish Creek screw trap) in 2014 was higher than the previous year and the highest since 2010 (Figure 11). In Satus Creek, 316 steelhead smolts were captured in the rotary screw trap and no estimate was calculated for this site due to low recapture numbers. The number of smolts captured (n=52) at the Ahtanum screw trap in 2014 continues to be too low to estimate juvenile outmigration from that watershed.

### Tributary Habitat RME

High summer water temperature (into the mid 20 degree celsius range) appears to be a limiting factor for steelhead early life history stages in the lower reaches of Satus, Toppenish, Simcoe and Ahtanum Creeks (Tables 9-15).

## 2. Introduction

The Yakama Reservation Watersheds Project (YRWP; 1996-035-01) originated as several separate BPA-funded projects, the earliest of which began in Satus Creek in 1996. Projects in Toppenish and Ahtanum Creeks were added soon afterward in an attempt to address the declining steelhead populations in three steelhead-producing tributaries of the lower Yakima River. All three of these watersheds are located on the Yakama Reservation. In March 1999, the Middle Columbia River Steelhead Distinct Population Segment was listed as threatened under the Endangered Species Act (ESA). Four populations (Satus, Toppenish, Naches, and Upper Yakima) are recognized by NOAA's National Marine Fisheries

Service for recovery purposes in the Yakima Basin and comprise the Yakima MPG (Major Population Group) (Yakima Fish and Wildlife Recovery Board 2009). Ahtanum Creek steelhead are grouped under the Naches population--although it is geographically removed may function as a separate population.

Monitoring the steelhead populations in Toppenish, Satus and Ahtanum creeks has proved to be difficult in many years. Adult steelhead are counted as they migrate up the Yakima River at Prosser Dam in the town of Prosser Washington. These counts using mostly video footage collected at the fish ladders (as well as a Denil fish ladder and trap for biological sampling) are believed to produce a fairly accurate adult migrant count for the Yakima MPG. Another counting facility (Roza) is located on the Yakima River farther upstream above the Naches River confluence and provides a direct count of adult steelhead migrating upstream of the boundary delineating the Upper Yakima population. No counting facilities are present to enable delineation of the Naches, Toppenish, or Satus Creek populations. Instead, redd count surveys provide an index of adult escapement on these Yakima River tributaries. Spawner surveys including a redd count and similar methods to those used at present have been performed in Satus Creek since 1988, in Toppenish Creek since 1997, and in Ahtanum Creek beginning in 2001. The accuracy of these counts varies from year to year and is dependent on snowpack and stream conditions during the spawning season. The poor accessibility and spring flooding of the upper section of the Toppenish Creek watershed and much of the Ahtanum Creek watershed has been the greatest obstacle to completing a redd survey that captures the entire spawning season in those tributaries.

## **Study Area**

### *Satus Creek*

Satus Creek is located in south-central Washington and drains the southeast portion of the Yakama Reservation with a watershed area of 710 mi<sup>2</sup> (Hubble 1992), more than 10 percent of the Yakima Basin area. It is the largest tributary of the Yakima River located on the Yakama Reservation. It is also the most downstream of the steelhead-producing tributaries in the Yakima basin and enters the Yakima River at river mile 69.6 (Columbia Basin Inter-Agency Committee 1964). Elevation ranges from 5800 feet near Potato Butte to 650 feet at the mouth. Logy Creek and Dry Creek are the largest tributaries of Satus Creek. Smaller tributaries include Mule-Dry Creek, Kusshi Creek, Shinando Creek, Bull Creek, and Wilson Charley Creek. A large section of Satus Creek located above Logy Creek becomes intermittent during the summer. Logy Creek itself drains a substantial area of the porous Simcoe Volcanic Field, and is perennial. Logy Creek provides summer stream flow for the lower portion of Satus Creek. Dry Creek, the largest tributary of Satus Creek, is intermittent in its lower 15 miles (below the "elbow") during the summer. However, the upper portion of this stream flows year round. Many of the smaller tributaries are also intermittent. Unlike the other lower tributaries of the Yakima River, irrigation withdrawals are limited to small pumps. The last irrigation dam (unused since 1981) was removed by YRWP in 2009. The watershed is also largely uninhabited with less than a dozen structures along its upper reaches, where grazing and logging are primary land uses. The lowest 10 miles of Satus Creek are within the Yakima River floodplain; here much of the land is farmed and irrigated with water withdrawn from the Yakima River near Parker Washington. Satus Creek has a particularly "flashy" hydrograph when compared to other streams in the Yakima basin due to its relatively low elevation and sparse vegetation. The highest recorded flows in Satus, Toppenish and Ahtanum creeks have been associated with midwinter rain-on-snow events.

### *Toppenish Creek*

Toppenish Creek is located in south central Washington and the entire watershed is situated within the boundary of the Yakama Reservation. Toppenish Creek is also a major tributary of the Yakima River with nearly as large a watershed area (622 mi<sup>2</sup>) as Satus Creek. The headwaters of Toppenish Creek are located on Lost Horse Plateau at a maximum elevation of 5200 feet. Simcoe Creek, the main tributary of Toppenish Creek, joins at river mile 32 (about the halfway point). The forks of Simcoe Creek and its main tributaries Agency and Wahtum Creeks also arise from Lost Horse Plateau, but at a slightly lower elevation. Toppenish Creek enters the Yakima River about 7 miles south east of the town of Toppenish, Washington at river mile 80 at an elevation of 650 feet. Along its approximately 70-mile length, stream morphology and watershed topography changes substantially. Through nearly the upper third of its length, Toppenish Creek flows through a remote forested canyon. Most of the upper Toppenish Creek watershed is tribal trust land managed for timber production and cultural resources. The middle third of Toppenish Creek is dominated by an alluvial fan. This area is managed for multiple uses including livestock grazing and some agriculture. Irrigation withdrawals begin in this region of the watershed, at the head of the alluvial fan. The lower portion of Toppenish Creek is heavily influenced by agriculture with a variety of crops grown (e.g. corn, wheat, hops, mint, orchard fruit and grapes). Flows and water quality are altered drastically by irrigation withdrawals, spills and return flows in the lower portion of Toppenish Creek. Much of the land adjacent to the lower third of the creek is devoted to waterfowl production and hunting. The USFWS Toppenish Creek Wildlife Refuge, Yakama Nation wildlife areas, and a number of private duck clubs provide a substantial amount of off-channel wetland habitat (managed to attract waterfowl) on the lower 30 miles of Toppenish Creek.

The hydrograph of Toppenish Creek is similar to other streams on the east slopes of the Cascade Range. Peak flows typically occur in early to mid-spring resulting from snowmelt, although rain-on-snow events during winter can cause substantial floods—sometimes the peak flow for the season. Flows decrease rapidly during late spring and early summer as the snowpack is depleted. Flows (and closely-related water temperature) are probably the limiting factor for steelhead production in portions of the Toppenish and Simcoe watersheds. Irrigation spills and return flows in the lower portions of Toppenish Creek are substantially greater than natural flow and are believed to be detrimental to salmonids because of their temperature, suspended solids, and the potential homing issues raised by flows originating from outside the Toppenish Creek watershed.

Marion Drain and its tributary Harrah Drain carry irrigation drainage from the agricultural area north of Toppenish Creek eastward to the Yakima River. During the irrigation season, most of their flow is mixed with Toppenish Creek flow and diverted south of the creek to the Satus Area of the Wapato Irrigation Project. Steelhead can access Marion and Harrah drains via the Yakima River outlet or Toppenish Creek. Redds are counted in both drains every year, but survival of steelhead eggs and parr may be low enough to consider Marion and Harrah drain a population sink.




Figure 1. Hydrograph on Toppenish Creek at river mile 40 for Water Year 2014.

### Ahtanum Creek

Ahtanum Creek is the next major Yakima River tributary located upstream from the Toppenish Creek watershed (Marion Drain). It forms the northern boundary of the Yakima Reservation and enters the Yakima River at river mile 106.9. Of the three steelhead-producing tributaries monitored by YRWP, Ahtanum Creek is the smallest with a watershed size of 173 mi<sup>2</sup>, but its base flow is comparable to that in perennial sections of Satus and Toppenish creeks because of its relatively higher watershed elevation. Elevations range from 6000 feet near Green Lake in the North Fork Ahtanum Creek watershed to 940 feet at the mouth. Flowing through the city of Union Gap and other small suburbs of the city of Yakima it is the most urban of the three watersheds and is where urbanization poses one of the greatest threats compared with other Yakima River tributaries. At one time irrigation withdrawals dewatered significant portions of the stream which is naturally perennial; however, regulated diversions now allow continuous flows. Irrigation withdrawals and returns still influence water quantity and quality.

### USGS 12502500 AHTANUM CREEK AT UNION GAP, WA




Figure 2. Hydrograph on Ahtanum Creek near the mouth for Water Year 2014.

### 3. Methods: Protocols, Study Designs, and Study Area

#### Adult steelhead abundance and distribution

**Protocol Title:** YRWP Adult steelhead abundance (1996-035-01) v1.0

**Protocol Link:** <http://www.monitoringmethods.org/Protocol/Details/110>

#### Protocol Summary:

Steelhead redd counts are used as an index of adult spawner abundance on the tributary spawning grounds in Lower Yakima River tributaries (Satus, Toppenish and Ahtanum Creeks).

In each lower Yakima River tributary (Satus, Toppenish, and Ahtanum), we attempt to perform a census survey on all recognized spawning habitat in each tributary. Yakama Nation Fisheries has attempted various survey methods (ground, raft, aerial) in some areas outside of these recognized spawning reaches but have not documented any redds.

The procedure for conducting steelhead redd counts has not changed significantly during past 30 years that the Yakama Nation has performed them. A three pass census count using the following technique is used. Two surveyors typically cover each 2 to 6 miles survey reach, walking in a downstream direction. In some smaller streams only one surveyor conducts the survey. Surveyors wear polarized glasses to aid in spotting redds. Each identified redd is marked with a GPS with an accuracy of +/- 30 feet. Redds are marked with fluorescent flagging to prevent counting redds identified on previous passes. Each redd is measured and its location in relation to the stream bank and thalweg are recorded. The presence or absence of direct cover is also noted on data sheets. It is unlikely that resident rainbow trout redds (or redds from other redd building species) are mistaken for anadromous steelhead redds because of the small size of all non-adult steelhead *O. mykiss* observed in these watersheds during population surveys (i.e. redd counts, snorkel surveys). The number of live steelhead adults and carcasses are also recorded. When possible, the sex of live steelhead and carcasses is noted. Surveyors will take care not to disturb spawning fish or possible staging pools when conducting spawner surveys. The survey begins after the sun breaks over the horizon to ensure that there is enough light to detect redds.

#### *Redd Count Expansion*

To translate our census count into an estimate of steelhead abundance we utilized the method outlined in Gallagher et al. (2007). The cumulative redd count was multiplied by the standard 2.5 fish per redd used for Washington streams for an estimate of spawning escapement. Other expansion coefficients were also considered.

#### *Ahtanum Bull Trout surveys*

The Yakama Nation has performed Bull trout surveys in the South Fork Ahtanum Creek annually since 2002. Each year they are conducted in the South Fork of Ahtanum Creek within an index reach from approximately RM 7.7 to RM 10.4. This section borders the Yakama Reservation and includes a small section of Tract C. upstream from Reservation Creek. Surveys are conducted as part of a program to track the status and trends of this species within the Yakima River watershed. Index reaches are situated in prime spawning reaches. In watersheds outside the South Fork Ahtanum, WDFW performs the surveys in cooperation with other agencies (e.g. USFWS, Joint Board, Yakama Nation, Yakima Basin Fish and Wildlife Recovery Board).



We perform these surveys using a protocol developed by WDFW (Eric Andersen; unpublished document-2011). In Summary, surveyor (s) walk upstream record, and flag redds during multiple passes. They are categorized as definite, probable, or possible. On the 3rd or last pass, GPS waypoints are collected. Live bull trout are documented during each pass as well. Only redds identified as probable or definite are included in the final count

### **Water temperature monitoring**

**Protocol Title:** YRWP continuous temperature monitoring (1996-035-01) v1.0

**Protocol Link:** <http://www.monitoringmethods.org/Protocol/Details/695>

### **Protocol Summary:**

High summer water temperature limits steelhead production in many reaches of the lower Yakima River tributaries (Toppenish, Satus, and Ahtanum)

We deployed data-loggers in the Ahtanum, Toppenish, and Satus watersheds to monitor water temperatures continuously during the warmer seasons when water temperatures can be a limiting factor for salmonid survival and growth. The Yakama Reservation Watersheds Project utilize this data to identify reaches where restoration projects would be most beneficial to salmonid populations and also to aid in management decisions that may affect water temperatures (i.e. management of irrigation diversions, riparian harvest, water withdrawals, etc.).

We deployed a total of forty nine devices in the three watersheds. Data-loggers (Onset Optic Stowaways and Onset Water Temp Pro v2) were launched in spring of 2014 and were programmed to collect water temperatures at 1 hour intervals. The units were encased in protective cages and secured to trees and roots using nylon coated aircraft cable. They were generally placed in pool tailouts that were less likely to dewater during the summer. Although some data-loggers were deployed beginning in April 2014, we only used data during the period between April 15<sup>th</sup> and October 15<sup>th</sup> to calculate descriptive statistics to evaluate in-stream conditions for salmonids. Several data-loggers were left in place year round to monitor water temperatures during the peak migration and spawning periods for steelhead (i.e. winter and spring).




Figure 3 Locations of temperature monitoring stations established between 1997 and 2014 in Satus, Toppenish and Ahtanum Creeks on the Yakama Reservation

### Juvenile steelhead out-migration

**Protocol Title:** YRWP Juvenile steelhead outmigration (1996-035-01) v1.0

**Protocol Link:** <http://www.monitoringmethods.org/Protocol/Details/111>

### Protocol Summary:

Rotary screw traps are used on Satus, Toppenish, and Ahtanum to estimate juvenile steelhead outmigrant abundance, timing, and downstream survival.




Figure 4. Location of rotary screw traps and PIT tag antenna arrays operated by YRWP in Satus, Toppenish, and Ahtanum Creeks on the lower Yakima River.

### Trapping and tagging Procedures

In many years we have not been able to obtain reliable estimates due to high flows that often occur in winter and spring. The heavy debris loads associated with these high-flow events have the potential to clog the screw trap cone and impinge and kill out-migrating juvenile fish. Our screw trapping protocol was similar at all four of the screw traps that we operate (upper Toppenish, lower Toppenish, Ahtanum, and Satus). Each trap was visited at least once a day usually between 6:30 AM and 11:00 AM. Fish were netted out, identified and target fish were held in 5-gallon buckets. Aeration using battery operated pumps was applied if needed. All juvenile steelhead were anesthetized in MS-222

before being handled. They were then enumerated, measured (mm), and weighed (g). Scales were collected on 100 individuals per season and location (up to 10 per day). We also collected fin clips from 100 individuals from Toppenish Creek. These samples were sent to CRITFC for DNA analysis to be used in several ongoing studies. On several occasions when large catches occurred ( $N > 300$ ) only a random sub-sample (first 100) were measured and weighed. We inserted PIT tags into a subsample (first 100) of captured steelhead smolts over 80 mm in length. PIT tagged fish were released several hundred meters upstream from the trap to estimate trap efficiency (i.e. mark-recapture). The upstream release site alternated between the right and left banks of the stream. Efficiency releases were made 4 times per week (Monday-Thursday) and release numbers and recaptures for the week were pooled. We set a target of 4000 steelhead juveniles to PIT tag and attempted to space our tagging effort throughout the season so the total out-migration was represented appropriately. Due to variable seasonal catches, there is no clear formula to achieve this and the target number tagged per day had to be adjusted several times during the season; however, we still ended the season short on PIT tagged fish.

Scales were collected from 100 individuals from each stream. We collected fin clips from 100 steelhead smolts for use in DNA studies as well. DNA samples are analyzed by CRITFC for their steelhead kelt reconditioning study. After handling we released steelhead juveniles, along with all recaptures and undersized fish 100 meters downstream. Physical data (water temperature, air temperature, and percent cloud cover) were recorded. The trap rotation rate (seconds per revolution) was recorded to evaluate operating efficiency.

#### *Juvenile out-migration estimate*

We utilized a Petersen's stratified capture-recapture model to estimate juvenile steelhead outmigrant abundance.

The assumptions of this model are:

- 1) The population is closed;
- 2) All fish have an equal probability of capture in the first period;
- 3) Marking does not affect catchability;
- 4) The fish do not lose their marks; and
- 5) All recovered marks are reported (Volkhardt et al. 2007)

Due to changes in factors shown to affect trap efficiency (i.e. stream flow, temperature, increasing smolt size) that occur as out-migration season progresses, we stratified our estimate temporally by week. Darr 2.02 software for R statistical software utilizing Darroch's (1961) maximum likelihood estimator for stratified data was used to obtain an estimate and its associated variance (Bjorkstedt 2005 and 2009). A one-trap study design was used. Since at least 5 recaptures are typically necessary to converge, Darr 2.02 incorporates an algorithm that automatically pools adjacent strata as needed. We enabled this algorithm because many weekly strata outside the peak migration period had fewer than five recaptures.

During periods of high flow when operation of the trap was not possible and trapping was suspended for several days, we interpolated daily catches by calculating the average of the three days before and the three days after the missed period.

## 4. Results

### a. Fish Population Status Monitoring (RM&E)

#### Adult abundance and distribution

4141 adult steelhead were enumerated at the Prosser Dam facility between July 1, 2013 and June 30, 2014. This is lower than the dam count from the previous 4 years which ranged from 4787 to 6793; however, still above the 10-year average (<http://ykfp.org/>).

#### *Satus Creek*

More redds were identified in 2014 (n=267) than in 2013 (n=223) or 2012 (n=152); but less than both 2011 (n=293) and 2010 (n=465). The largest number of redds were identified in the mainstem Satus (n=120) followed by Dry Creek (n=71). In the 2nd largest tributary, Logy Creek, 59 redds were identified in 2014. Also, redds were identified in some of the smaller tributaries of Satus Creek and numbers ranged from zero in Shinando Creek to 12 in Kusshi Creek. Fewer redds were identified in the upper part of Satus Creek where a fire burned in 2013 than in previous years when this area often had the highest density of redds.




Figure 5 Number of steelhead redds in the Satus Creek watershed 1997 to 2014




Figure 6 Locations of steelhead redds in the Satus Creek watershed 2014

*Toppenish Creek*

For the 2014 spawning season lasting from March 2014 through June 2014 a total of 134 steelhead redds were identified in the Toppenish Creek watershed during a season when conditions for counting redds was generally good. Only two passes of the upper 3 reaches of Toppenish Creek could be completed due to high runoff early in the season.




Figure 7. Number of steelhead redds per year in the Toppenish Creek watershed since 1997



Thirty-four percent of these redds were found in the Simcoe Creek watershed the largest tributary and the remaining 66 percent were found in the mainstem and smaller tributaries such as the North Fork Toppenish Creek and Willy Dick Creek.




Figure 8. The Location of steelhead redds in the Toppenish Creek watershed in 2014.

#### *Ahtanum Creek*

In 2014, no steelhead redd counts could be completed due to the unavailability of staff during a short period in early April when redd counts were possible. After this period, the flows were too high to conduct redd count surveys.

Bull trout were surveyed in late summer and autumn 2014 in the South Fork Ahtanum watershed which forms the northern boundary of the Yakama Reservation. During our 2014 bull trout surveys, we completed individual passes on Sept-4, Sept-17, and Oct-2. Two redds and 1 live adult bull trout were documented in 2014--an increase from 0 the previous year.

#### *Redd Count Expansion*

The cumulative redd count was multiplied by 2.5 fish per redd for an estimate of spawning escapement of 667 adult steelhead for the Satus watershed and 335 for the Toppenish watershed for the 2014 season. The 2.5 fish per redd expansion used for our estimate is similar to those obtained through studies of other Middle-Columbia steelhead populations in Oregon ranging from 2.1 to 2.6 fish per redd (Poxon et. al 2011). Better methods of enumerating adult steelhead utilizing PIT tagged adults and antenna arrays are currently under development.

## Juvenile Out-migration

### *Satus Creek*

The Satus Creek screw trap is located within 1 mile from the mouth of the stream and is located in a position to intercept fish from all the spawning and rearing habitat.

Although the trap was deployed on October 22, 2013 for the 2014 season, we did not catch the first steelhead juvenile until November 10. The peak catch occurred at the end of February when 54 steelhead juveniles were captured although another spike in outmigration occurred in April and May (Figure 10). A total of 316 steelhead juveniles were captured in the screw trap between November, 2013 and June 2014. Of these juveniles, we PIT tagged 283 and released most of these as part of a mark-recapture study to obtain an outmigrant abundance estimate. Six of these released fish were recaptured over the season producing an average (pooled) trap efficiency of 2.31%. The average length (114 mm) was smaller than the previous year but the frequency distribution was similar to other years.

In addition to steelhead juveniles, we captured 31 chinook juveniles and 3 lamprey in the Satus Creek trap.




Figure 9. Number of steelhead juveniles captured per day compared with steam stage at the Satus Creek screw trap in 2014.

### Upper Toppenish Creek Screw Trap

We operated a rotary screw trap on Toppenish creek from October 2013 through June 2014, a period which encompasses most of the outmigration period. We captured a total of 4509 steelhead parr/smolts and PIT tagged 2448. The daily catch of steelhead juveniles peaked in early December (Figure 9) as typically observed. Most of the PIT tagged steelhead juveniles were released several hundred meters above the screw trap as part of a mark-recapture study. An estimate of 46,470 SE 5008 juvenile steelhead outmigrants was obtained for the 2014 season using this method. The



mean length for all measured outmigrants was just over 126 mm (fork length) with a frequency distribution similar to previous years with the majority of captured steelhead between 100 and 200 mm. We also captured 4 chinook juveniles and 3 lamprey juveniles at this site during the 2013 and 2014 season.




Figure 10. Steelhead juvenile outmigration timing. Daily catch at the Toppenish Creek trap compared with stream stage for the 2014 season.




Figure 11. Estimated number of steelhead juvenile outmigrating from spawning and rearing habitat on Toppenish Creek. Error bars represent 95% CI

Lower Toppenish Creek Screw Trap

We operated a second rotary screw trap near the mouth of Toppenish Creek at a location that has been in use since 2011. This trap was deployed in mid November 2013 and retrieved in June 2014. We captured 11 steelhead juveniles, 172 chinook juveniles. Unlike the screw trap located upstream at river mile 30 on Toppenish Creek where outmigration typically peaks in late fall or early winter, most juvenile steelhead were captured at the lower screw trap during the spring. This pattern, similar to previous years, indicates that significant overwintering in the lower reaches of Toppenish Creek between the traps.

*Ahtanum Creek*

In 2014, we captured 52 juvenile *O.mykiss* in between the months of November and June and most of these (n=38 were tagged). Sample size is too low to obtain an estimate or draw many conclusions from these data. In addition to *O.mykiss*, we captured 18 chinook, 6 coho, and 7 lamprey juveniles in the Ahtanum Creek screw trap.

**b. Tributary Habitat RM&E**

Water Temperature Monitoring

*Satus Creek*

Mean daily averages in the mainstem Satus Creek ranged from 9.9°C downstream from the falls to 18.02 °C below Dry Creek (Table 1) in 2014. The greatest instantaneous maximum for the Satus Creek watershed was 27.1°C below the Dry Creek Confluence (RM 18.7) as well as the highest maximum 7-day maximum water temperature (26.3 °C). Water temperature was also the highest at this site in 2012, although the water temperatures were slightly lower (Table 1).

Table 1. Descriptive statistics for water temperature for 2014 for the period between April 15th and October 15th at 17 locations in the Satus Creek watershed. Maximum weekly average temperature are in bold text.

<b>Location</b> (river mile in parenthesis)	Instantaneous Maximum (°C)	Instantaneous Minimum (°C)	Mean Daily Maximum (°C)	Mean Daily Average (°C)	Mean Daily Minimum (°C)	Maximum Daily Average (°C)	<b>Maximum 7-Day Maximum</b> (°C)	Maximum 7-Day Average (°C)
Falls (44)	16.96	3.35	11.55	9.94	8.48	14.45	<b>15.89</b>	14.05
Satus at Wilson Charley (39.5)	22.78	3.67	15.24	12.36	10.21	18.69	<b>21.82</b>	18.27
4th Crossing (34.1)	23.95	3.96	17.02	14.68	12.57	21.33	<b>22.85</b>	20.54
High Bridge	25.48	4.22	17.62	15.02	12.85	22.08	<b>24.42</b>	21.31

(32.4)									
Above Logy Creek (23.6)	25.87	6.03	19.6	16.85	14.35	22.29	<b>25.04</b>	21.58	
Below Dry Creek (18.7)	28.2	6.94	20.01	17.16	14.65	24.01	<b>26.63</b>	22.96	
Plank Rd. (7.4)	27.16	8.74	19.67	18.02	16.6	24.53	<b>26.11</b>	23.83	
N. Satus Rd. (1.2)	24.82	9.36	18.57	17.3	15.96	22.89	<b>24.03</b>	22.08	
Logy at Falls(12.5)	lost								
Logy at Fourth Crossing (8.8)	20.51	4.01	14.01	12.48	10.98	18.44	<b>19.39</b>	17.56	
Logy Mouth (0.5)	24	5.54	16.49	14.85	13.07	21.77	<b>22.8</b>	20.83	
Dry Creek at Falls (25.7)	16.92	4.04	12.55	10.46	8.42	13.34	<b>15.68</b>	12.85	
Dry Creek at Elbow Crossing (18.5)	20.06	6.36	15.75	14.2	13.06	18.07	<b>19.7</b>	17.79	
Dry Creek at Mouth (1.2)	21.25	7.67	16.72	15.02	13.48	18.91	<b>20.82</b>	18.54	
Section corner source (4.6)	8.97	7.24	8.48	7.92	7.62	8.18	<b>8.84</b>	8.12	
Section corner mid crossing (2.7)	lost								
Section corner lower crossing (1.2)	14.58	4.9	11.77	9.74	7.94	11.81	<b>13.82</b>	11.35	

### *Toppenish Creek*

In 2014, mean daily average temperatures in the mainstem Toppenish Creek ranged from 8.2°C on Panther Creek in the head waters of Toppenish Creek (RM 65.7) to 17.52°C above the Unit 2 diversion (RM 26.5). The highest instantaneous maximum of 24.61°C occurred at the Cleparty diversion on the alluvial fan at river mile 36.1 as well as the highest MWMT (26.2°C). By most standards the summer water temperature in 2014 in the lower portion of Toppenish Creek (below the Unit 2 diversion) was higher than most salmonids can tolerate (Table 2). In Simcoe Creek, the highest instantaneous maximum water temperature occurred higher up in the watershed at Simcoe Creek road; however, this unit was placed in a shallow area outside of the main current of the stream which likely affected the measurements during the low flows of August. The lowest instantaneous maximum temperature (16.7°C) in the Simcoe Creek watershed was recorded in Agency Creek downstream from a large spring near the Jeldwin Mill (Table 2).

Table 2. Descriptive statistics for water temperatures at 13 locations in the Toppenish Creek watershed for 2014 during

the period from April 15th to October 15th. Maximum Weekly Maximum Temperature in bold text.

Location (river mile in parenthesis)	Instantaneous Maximum	Instantaneous Minimum	Mean Daily Maximum	Mean Daily Average	Mean Daily Minimum	Maximum Daily Average	Maximum 7-Day Maximum	Maximum 7-Day Average
Panther Creek	14.1	2.1	10.1	8.2	6.4	12.2	<b>13.5</b>	11.5
Topp. at N. Fork confluence (55.9)	19.79	3.7	13.9	11.79	10.17	16.49	<b>18.76</b>	15.99
Topp. at swim hole (47.2)	21.6	4.48	15.49	14.03	12.73	19.32	<b>20.71</b>	19.01
1 mile below swim hole (45.9)	22.82	6	16.8	14.96	13.42	20.05	<b>21.96</b>	19.63
1 mile above lateral (45.1)	22.35	5.1	16.58	14.99	13.63	20.14	<b>21.62</b>	19.79
Topp. above lateral (44.2)	22.5	4.9	16.2	14.6	13.3	20.2	<b>21.7</b>	19.8
At three way (43.1)	23.14	5.1	16.96	15.23	13.7	20.75	<b>22.32</b>	20.24
Topp. At Cleparty Diversion (36.1)	24.61	7.65	18.74	16.34	14.5	21.12	<b>23.76</b>	20.62
Topp. At Shaker Church Rd. (36.1)	22.68	6.56	16.51	15.41	14.4	20.4	<b>22.03</b>	19.83
Topp. at Unit 2 (26.5)	24.03	7.75	18.14	17.52	16.86	23.6	<b>23.22</b>	22.74
Simcoe below Forks (18.9)	21.01	4.04	14.51	13.18	11.84	18.81	<b>20.06</b>	18.39
Simcoe at Simcoe Cr. Rd. (15.3)	26.4	4.53	18.21	14.86	12.13	21.54	<b>25.48</b>	21.05
Simcoe below Stephensen Rd.. (8.1)	23.83	6.15	17.35	15.55	13.77	21.33	<b>22.13</b>	20.27
Barkes Rd (2.7)	24.65	6.99	18.25	16.8	15.29	22.83	<b>23.7</b>	21.78
Agency Creek at Wesley (0.5)	16.7	8.52	14.04	12.3	10.92	14.3	<b>16.18</b>	13.8

Mean daily averages ranged from 11.8 C in the South Fork of Ahtanum Creek one mile above the confluences to 17.3°C at the USGS gage several hundred meters upstream from the mouth in Union Gap (Table 3). The highest instantaneous maximum of 29.8°C was also recorded at this location.

Table 3. Descriptive statistics for water temperatures at 3 locations in the Ahtanum Creek watershed for 2014. Maximum Weekly Maximum Temperature in bold text.

Location (river mile in parenthesis)	Instantaneous Maximum	Instantaneous Minimum	Mean Daily Maximum	Mean Daily Average	Mean Daily Minimum	Maximum Daily Average	<b>Maximum 7-Day Maximum</b>	Maximum 7-Day Average
South Fork Ahtanum at Mouth(1.0)	21.77	2.98	14.73	11.82	9.42	17.64	<b>20.87</b>	17.14
AID Diversion (18.9)	25.14	3.49	16.67	14.02	11.62	22.86	<b>24.32</b>	20.32
At USGS Gauge (0.5)	29.82	6.28	19.51	17.34	15.3	26.23	<b>28.16</b>	25.29

## 5. Synthesis of Findings: Discussion/Conclusions

### a. Fish Population Status Monitoring (RM&E)

Information on Adult Steelhead Abundance and Juvenile out-migrant abundance is valuable in evaluating restoration efforts, managing natural resources (fish, water, etc.) and in determining if recovery goals for this species are being met.

In 2012 Redd count surveys were obstructed by substantial flooding, probably causing a substantial underestimation of spawning activity. Although conditions improved in 2013 and 2014, We have concluded that redd counts are not the best method to monitor adult steelhead spawner abundance. We do however intend to continue redd count surveys because they provide useful information on spawning distribution and they provide an opportunity for reconnaissance of critical steelhead spawning and rearing habitat. Spatial information (GPS coordinates) collected during redd count surveys using the same protocol over multiple years have been used to identify reaches where spawning habitat should be protected or restored. Alternative methods to using PIT tagged adult steelhead and instream antenna arrays to determine spawner abundance and fish per redd are under development with PIT tag antenna arrays that were installed in 2011 and 2014 in the Satus, Toppenish and Ahtanum watersheds.

Our screw trap in the upper Toppenish Creek and the method of operation that we have employed for the last four years appears to provide relatively precise estimates (CV coefficient of variation ranging from 6.78% to 16.71%) for the number of steelhead migrating from the spawning and rearing areas upstream. However, the other three screw traps do not produce adequate estimates of outmigrant abundance, although some useful information have been collected on

timing, growth, and survival of smolts captured and PIT tagged at these traps. One change proposed for the 2016 season is to discontinue trapping at the lower Toppenish site and move that trap to Marion Drain to determine if this constructed tributary to Toppenish Creek is producing steelhead juveniles. An absence of captured smolts, as we expect, would support our efforts to isolate this drain (where adult steelhead are observed spawning each year) from Toppenish Creek.

### **a. Tributary Habitat (RM&E)**

With Maximum 7 day average water temperatures consistently above 23 degrees C , water temperature appears to be a limiting in the lower portions of Toppenish, Simcoe, Ahtanum and Satus Creeks.

## References

- Bjorkstedt (2005 and 2009) Darrochs (1961) Petersen Estimator (Darr v 2.02 R code) available at <http://swfsc.noaa.gov/textblock.aspx?Division=FED&id=3346>.
- Carlson, S. R., L. G. Coggins, C.O. Swanton. 1998. A simple stratified design for smolt mark-recapture estimation for salmon smolt abundance. *Alaska Fishery Research Bulletin* 5(2): 88-102.
- Gallagher, S.P., P.K. Hahn, and D.H. Johnson. 2007. Redd Counts. Pages 197-231 in D.H. Johnson, B.M. Shrier, J. A. Knutzen, X Augerot, T.A. O'Neil, and T. N. Pearsons. 2007. *Salmonid field protocols handbook: techniques for assessing status and trends in salmon and trout populations*. American Fisheries Society, Bethesda Maryland.
- Hubble, J. D. 1992. A study of summer steelhead, *Oncorhynchus mykiss*, in several intermittent tributaries of the Satus Creek Basin. M.S. Thesis, Central Washington University, Ellensburg WA. 86 pp.
- Johnson D.H., B.M. Shrier, J.S. O'Neil, J.A. Knutzen, X Augerot, T.A. O'Neil, T.N. Pearsons. 2007. *Salmonid field protocols handbook: techniques for assessing status and trends in salmon and trout populations*. American Fisheries Society, Bethesda, Maryland.
- Junk, W.J., P.B. Bayley, and R.E. Sparks. 1989. The Flood Pulse Concept in River Floodplain Systems. p. 110-127. In D. P. Dodge [ed.] *Proceedings of the International Large River Symposium*. Can. Spec. Publ. Fish. Aquat. Sci. 106 pp.
- McClure, Michelle, Tom Cooney. 2005. Updated population delineation in the interior Columbia Basin, NOAA Fisheries Memorandum, [http://www.nwfsc.noaa.gov/trt/updated\\_population\\_delineation.pdf](http://www.nwfsc.noaa.gov/trt/updated_population_delineation.pdf). 16 pp.
- Northwest Power Planning Council. 2000. *Columbia River Basin Fish and Wildlife Program*. 56pp.
- Rawding, Dan and P. Cochran. 2007. *Wind River Winter and Summer Steelhead Adult and Smolt Population Estimates from Trapping Data, 2006*. Project No. 199801900 Contract No. 19617 For The Bonneville Power Administration Portland, Oregon
- Schuett-Hames, D, A. Pleus, L. Bullchild and S. Hall, eds, 1994. *Timber-Fish-Wildlife Ambient Monitoring Program Manual*. TFW-AM9-94-001. Northwest Indian Fisheries Commission, Olympia, Washington.
- Susac, G. L., S. E. Jacobs. 1999. Evaluation of spawning ground surveys for indexing the abundance of winter steelhead in Oregon coastal basins. Oregon Department of Fish and Wildlife Federal Aid in Sport Fish Restoration Program: F-145-R-08.
- Stanford, J. A. and J.V. Ward. 1992. Management of aquatic resources in large catchments: recognizing interactions between ecosystem connectivity and environmental disturbance, In. Naiman, R.J. (Ed.), *Watershed Management*. Springer-Verlag, New York. Pp 91-123.
- Terraqua, Inc. 2009. *A Field Manual of Scientific Protocols for Downstream Migrant Trapping within the Upper Columbia Monitoring Strategy*. BPA, ISEMP. (.pdf)
- Terraqua, Inc. 2009. *A Field Manual of Scientific Protocols for Steelhead Redd Surveys within the Upper Columbia Monitoring Strategy*. BPA, ISEMP. (.pdf)
- Thedinga, J.F. 1994. Determination of salmonid smolt yield with rotary screw traps in the Situk River, Alaska, to predict the effects of glacial flooding. *North American Journal of Fisheries Management* 14: 837-851.

*Volkhardt, G.C., S.L. Johnson, B.A. Miller, T.E. Nicholson, and D.E. Seiler. 2007. Rotary Screw Traps and Inclined Plane Traps. Pages 235-266 in D.H. Johnson, B.M. Shrier, J. A. Knutzen, X Augerot, T.A. O'Neil, and T. N. Pearsons. 2007. Salmonid field protocols handbook: techniques for assessing status and trends in salmon and trout populations. American Fisheries Society, Bethesda Maryland.*

*Yakima Subbasin Fish and Wildlife Planning Board. 2004. Yakima Subbasin Plan. 557pp. + appendices. <http://www.nwcouncil.org/fw/subbasinplanning/yakima/plan/>.*

*Yakima Subbasin Fish and Wildlife Planning Board. 2004. Yakima Subbasin Plan Supplement. 47pp. <http://www.nwcouncil.org/fw/subbasinplanning/yakima/plan/Supplement.pdf>.*

*Yakima Subbasin Fish and Wildlife Recovery Board. 2009. Draft 2009 Yakima Steelhead Recovery Plan. 288pp. <http://www.ybfwrp.org/RecoveryPlan/YakimaSteelheadPlan.pdf>*



## **Appendix A: Use of Data & Products**

Steelhead redd count, juvenile outmigrant abundance, and juvenile outmigrant survival data for Toppenish Creek are located at:

<http://ykfp.org/docsindex.htm>

## Appendix B: Detailed Results

Table B-1. Number of steelhead trout redds per reach in the Satus Creek watershed in 2014. River miles are in parentheses.

<b>Stream</b>	<b>Start location, RM</b>	<b>End location, RM</b>	<b>Distance (miles)</b>	<b># of Redds</b>
<b>SATUS</b>	Falls (44.1)	Wood Bridge (40.8)	4.2	<b>4</b>
(3 passes)	Wood Bridge (40.8)	County Line (36.4)	4.4	<b>6</b>
	County Line (36.4)	High Bridge (32.4)	4	<b>33</b>
	High Bridge (32.4)	Holwegner(28.4)	4.8	<b>40</b>
	Holwegner (28.4)	2nd X-ing (23.7)	3.9	<b>11</b>
	2nd X-ing (23.7)	1st Xing (20.2)	3.5	<b>1</b>
	1st X-ing (20.2)	Gage (17.4)	2.8	<b>0</b>
	Gage (17.4)	Rd 23 (13.1)	4.3	<b>0</b>
<b>Total</b>			<b>31.9</b>	<b>120</b>
<b>LOGY</b>	Falls (14)	Spring Cr (11)	3	<b>14</b>
(3 passes)	Spring Cr (11)	S. C. Ford (9.5)	1.5	<b>4</b>
	S. C. Ford (9.5)	3rd Xing (3.5)	6	<b>26</b>
	3 <sup>rd</sup> Xing (3.5)	Mouth (0.0)	3.5	<b>15</b>
<b>Total</b>			<b>14</b>	<b>59</b>
<b>DRY</b>	South Fk. (27.8)	Saddle ( 24)	3.6	<b>17</b>
(3 passes)	Saddle (24)	Elbow Xing (18.25)	5.75	<b>28</b>
	Elbow Xing (18.25)	Seattle Cr (14)	4.25	<b>13</b>
	Seattle Cr (14)	Rd 75 bend (8.75)	5.25	<b>6</b>
	Rd 75 bend (8.75)	Power Line Ford (2.5)	6.25	<b>5</b>
	Power Line Ford (2.5)	Mouth (0.0)	2.75	<b>2</b>
<b>Total</b>			<b>27.85</b>	<b>71</b>
<b>W. CHARLEY</b>	Forks (1.9)	Mouth (0.0)	<b>1.9</b>	<b>3</b>
<b>KUSSHI</b>	Top (11th) Xing (4.5)	Mouth (0.0)	<b>4.5</b>	<b>12</b>
<b>SHINANDO</b>	Ford (0.5)	Mouth (0.0)	<b>0.5</b>	<b>0</b>
<b>MULE DRY</b>	Yakima Chief Rd. (15.4)	Rd. 39 (4)	<b>11.4</b>	<b>2</b>

Table B2. Number of steelhead redds in the Toppenish Creek watershed in 2014.				
<b>Upper Toppenish Creek watershed</b>			<b>Distance</b>	<b>Number of</b>
(River Miles at Confluence in Parentheses)			<b>miles</b>	<b>Redds</b>
<b>Toppenish</b>	O Connor Cr (65.7)	"East Bank" (61.1)	4.6	<b>18</b>
	"East Bank" (61.1)	NF confluence (55.4)	5.7	<b>14</b>
	North Fork (55.4)	Washout (50.9)	4.5	<b>13</b>
	Washout (50.9)	Willy Dick Cr (48.5)	2.5	<b>6</b>
	Willy Dick Cr (48.5)	Olney Lateral (44.2)	4.3	<b>19</b>
	Olney Lateral (44.2)	Pom Pom Rd. (38.9)	5.3	<b>8</b>
	Pom Pom Rd. (38.9)	Shaker Church Rd. (35.9)	3	<b>8</b>
<b>Total</b>			<b>29.9</b>	<b>86</b>
<b>N. Fork Toppenish</b>	NF Falls (4)	NF confluence (0)	<b>4</b>	<b>3</b>
<b>Willy Dick</b>	old logging site (4)	Confluence (0)	<b>4</b>	<b>0</b>
<b>Simcoe Creek Watershed</b>				
<b>Simcoe</b>	NF at 2nd crossing (6.5)	Diamond Dick Cr (3.4)	3.1	<b>2</b>
	NF at Diamond Dick Cr (3.4)	NF/SF confluence (0)	3.4	<b>5</b>
	SF 6 mi above confluence (6.2)	SF 3 mi above confluence (3)	3.2	<b>9</b>
	SF 3 mi above confluence (3)	NF/SF confluence (0)	3	<b>0</b>
	NF/SF confluence (18.9)	Simcoe Creek Rd. (15.3)	3.6	<b>2</b>

	Simcoe Creek Rd. (15.3)	Wesley Rd. (10.1)	5.2	<b>0</b>
	Wesley Rd. (10.1)	N. White Swan Rd. (8.1)	2.0	<b>4</b>
	N. White Swan Rd. (8.1)	Stephenson Rd. (5.9)	2.2	<b>6</b>
<b>Total</b>			<b>25.7</b>	<b>32</b>
<b>Agency</b>	Falls (8.9)	Lateral Canal (4.4)	4.5	<b>1</b>
	Lateral Canal (4.4)	Confluence (0)	4.4	<b>2</b>
<b>Total</b>			<b>8.9</b>	<b>3</b>
<b>Wahtum</b>	Yesmowit Rd. (3.6)	Confluence (0)	<b>3.6</b>	<b>10</b>
<b>Total</b>			<b>76.1</b>	<b>134</b>

Table B3. Steelhead juvenile catch and mark-recapture data stratified weekly for the upper Toppenish creek screw trap for the MY 2014 season. Recaptures for each week are adjusted using PIT tag codes to exclude fish tagged in previous seasons and to assign fish to appropriate release group (e.g. fish in the trap recapture week  $i + 1$  column were recaptured the week following the week they were released above the trap)

Statistical Week ( $i$ )	Dates	Number Captured (week $i$ )	Number Released Upstream (week $i$ )	Number Recaptured (week $i$ )	Number Recaptured Week (week $i+1$ )+
Week 1	9/30-10/6	29	19		
Week 2	10/07-10/13	5	5		
Week 3	10/14-10/20	3	0		
Week 4	10/21-10/27	0	0		
Week 5	10/28-11/03	25	24	1	1
Week 6	11/04-11/10	45	34	5	
Week 7	11/11-11/17	25	16	4	
Week 8	11/18-11/24	0	0	0	
Week 9	11/25-12/01	24	40	0	

Week 10	12/02-12/08	2106	284	44	
Week 11*	12/09-12/15	84	24	1	
Week 12	12/16-12/22	51	20	5	
Week 13	12/23-12/29	27	27	1	
Week 14	12/30-01/05	1	0	0	
Week 15	01/06-01/12	20	14	1	
Week 16	01/13-01/19	15	14	1	1
Week 17	01/20-1/26	7	7	3	
Week 18	01/27-02/02	2	2	0	
Week 19	02/03-02/09	10	10	3	1
Week 20	02/10-02/16	129	127	5	10
Week 21	02/17-02/23	678	173	19	
Week 22	02/24-03/02	399	187	63	1
Week 23	03/03-03/09	66	29	2	
Week 24	03/10-03/16	52	34	0	
Week 25	03/17-03/23	87	66	0	
Week 26	03/24-03/30	128	113	9	
Week 27	03/31-04/06	30	18	1	
Week 28	04/07-04/13	30	15	0	
Week 29	04/14-04/20	22	18	1	
Week 30	04/21-04/27	18	18	0	0
Week 31	04/28-05/04	80	46	0	0
Week 32	05/05-05/11	72	50	1	0
Week 33	05/12-05/18	121	119	7	2
Week 34	05/19-05/25	133	79	7	0
Week 35	05/26-06/01	40	26	5	
Week 36	06/02-06/08	2	0		
<b>Total</b>	<b>09/30-06/03</b>	<b>4566</b>	<b>1655</b>	<b>189</b>	<b>16</b>

\*number captured during this week includes some dates when the trap was not operated and numbers were interpolated using

an average of the catch 3 days before and after the period of inoperability.

+two juvenile steelhead were recaptured more than 1 week after its release. one fish was tagged the previous year and was excluded.




Figure B1. Length frequency distribution of steelhead juveniles captured in the upper Toppenish rotary screw trap between October 1st 2013 and June 10th 2014.




Figure B2. Length frequency distribution of steelhead juveniles captured in the Satus Creek rotary screw trap between October 1st 2013 and June 10th 2014.




Figure B3. Length frequency distribution of steelhead juveniles captured in the Ahtanum Creek rotary screw trap between October 1st, 2013 and June 10th, 2014.

## Appendix C: List of Metrics and Indicators

Category	Subcategory	Subcategory Focus 1	Subcategory Focus 2	Specific Metric Title
Fish	Abundance of Fish	Fish Life Stage: Adult - Outmigrant	Fish Origin: Natural	Estimate of juvenile steelhead outmigrant abundance
Fish	Abundance of Fish	Fish Life Stage: Juvenile - Migrant	Fish Origin: Natural	Rotary screw trap juvenile steelhead daily catch
Fish	Abundance of Fish	Fish Life Stage: Adult - Spawner	Fish Origin: Natural	Steelhead redd count (an index of adult spawner abundance)
Fish	Abundance of Fish	Fish Life Stage: Juvenile - Fry/Parr	Fish Origin: Natural	Steelhead parr density: Steelhead parr / 100m <sup>2</sup>
Landscape Form & Geomorphology	Abundance of Habitat Types	Habitat Type: Channels		Stream habitat type composition
Landscape Form & Geomorphology	Abundance of Instream Wood Structures			Large Woody Debris
Landscape Form & Geomorphology	Cover	Habitat Type: Riparian Zone		Canopy Cover
Sediment/Substrate/Soils	Composition: Substrate/Soil-Dominant Size			Wolman Pebble Count
Water Quality	Water Temperature			Continuous water temperature
Water Quality	Water Temperature			Maximum Weekly Maximum Water Temperature
Hydrology/Water Quantity	Flow			Stream discharge
Hydrology/Water Quantity	Ground Water Level Change			Ground water lever