



YAKIMA/KLICKITAT FISHERIES PROJECT MONITORING AND EVALUATION Yakima Subbasin



PROJECT NUMBER 1995-063-25

Report covers work performed under BPA contract #00056662 REL 214

Report was completed under BPA contract #00056662 REL 223

FINAL REPORT For the Performance Period May 1, 2020 through April 30, 2021 September 8, 2021

Joe Blodgett, Project Coordinator
Mark Johnston, Research Scientist/Project Lead
Bill Bosch, Research and Data Coordinator/Editor
Yakima/Klickitat Fisheries Project

THE CONFEDERATED TRIBES AND BANDS OF
THE YAKAMA NATION
Toppenish, WA 98948

This report was funded by the Bonneville Power Administration (BPA), U.S. Department of Energy, as part of BPA's program to protect, mitigate, and enhance fish and wildlife affected by the development and operation of hydroelectric facilities on the Columbia River and its tributaries. The views in this report are the authors' and do not necessarily represent the views of BPA.

This report should be cited as follows:

Blodgett, J., M. Johnston, and B. Bosch (editors). 2021. Yakima-Klickitat Fisheries Project Monitoring and Evaluation – Yakima Subbasin, Final Report for the performance period May/2020-April/2021, Project number 1995-063-25, 388 electronic pages.

Table of Contents

List of Tables	i
List of Figures	iii
Acknowledgements	1
Executive Summary	3
Introduction	7
Fish Population Status Monitoring	12
Status and Trend of Adult Fish Populations (Abundance)	12
Status and Trend of Adult Productivity	17
Status and Trend of Juvenile Abundance	24
Status and Trend of Juvenile Migration Survival to McNary Dam	29
Status and Trend of Juvenile Productivity (smolt-to-adult returns)	32
Status and Trend of Spatial Distribution (Redd Counts)	39
Status and Trend of Diversity Metrics	46
Habitat Monitoring	51
Status and Trend of Fine Sediment	51
Harvest Monitoring	54
Marine and Mainstem Columbia Fisheries	54
Yakima Subbasin Fisheries	58
Hatchery Research	61
Effect of Artificial Production on the Viability of Natural Fish Populations	61
Effectiveness of Hatchery Reform	66
Predation Management and Predator Control	71
Avian Predation Index	71
Fish Predation Index and Predator Control	80
Adaptive Management and Lessons Learned	90
References and Project-related Publications	92
APPENDICES	103

List of Tables

Table 1. Adult-to-adult productivity indices for upper Yakima wild/natural spring Chinook	19
Table 2. Adult-to-adult productivity indices for Naches River Subbasin wild/natural spring Chinook	21
Table 3. Adult-to-adult productivity indices for Cle Elum SRF spring Chinook	22
Table 4. Estimates of adult-to-adult productivity indices for Yakima Basin natural-origin coho	23
Table 5. CESRF total releases of Spring Chinook by brood year, treatment, and acclimation site	26
Table 6. Total releases of Coho by brood year, life stage, and brood source	27
Table 7. Total releases of fall-run Chinook by release year and release site	28
Table 8. Total releases ¹ of summer-run Chinook by release year and release site	28
Table 9. Estimated smolt passage at Prosser Dam for Yakima Basin wild/natural and hatchery-origin spring Chinook and coho	29

Table 10. Estimated smolt passage at Chandler and smolt-to-adult return indices (Chandler smolt to Yakima R. mouth adult) for Yakima Basin wild/natural and CESRF-origin spring Chinook.....	34
Table 11. Average combined hatchery- and natural-origin smolt counts at Prosser for fish returning at age-3, -4, and -5, combined adult returns to Prosser Dam of all age classes, and estimated Prosser smolt-to-adult return indices for Yakima River fall-run Chinook for adult return years 1988-2019.	35
Table 12. Preliminary estimates of Prosser-to-Prosser smolt-to-adult survival (SAR) indices for adult returns from hatchery- and natural-origin coho for the Yakima reintroduction program, juvenile migration years 2000-2018.	36
Table 13. Preliminary McNary Dam smolt to Bonneville Dam adult SAR-indices for hatchery-origin PIT-tagged summer and fall-run chinook released in the Yakima subbasin by brood year and life stage at release, 2006-2015.....	37
Table 14. Preliminary McNary Dam smolt to Bonneville Dam age-3 adult return (SAR) indices for hatchery-origin PIT-tagged coho released as smolt (sm) or parr ^a in Lower Yakima (LY), Naches (Na), and Upper Yakima (UY) mainstem or tributary areas, brood years 2003-2014.	37
Table 15. Yakima Basin spring Chinook redd counts and distribution, 1981 – present..	40
Table 16. Yakima Basin coho redd counts and distribution, 1998 – present.....	43
Table 17. Results from Taneum Creek adult out-plant study.	45
Table 18. Sex ratio of upstream migrating fall Chinook sampled at the Prosser Dam right bank denil ladder and fish trap, 2001-present.	47
Table 19. Sample size (N), mean fork and mid-eye to hypural plate (MEH) lengths (cm), and weights (pounds) of upstream migrating fall Chinook sampled at the Prosser Dam right bank denil ladder and fish trap, 2001-present.....	47
Table 20. Sex ratio of upstream migrating coho sampled at the Prosser Dam right bank denil ladder and fish trap, 2001-present.....	48
Table 21. Sample size (N), mean fork and mid-eye to hypural plate (MEH) lengths (cm), and weights (pounds) of upstream migrating coho sampled at the Prosser Dam right bank denil ladder and fish trap, 2001-present.	48
Table 22. Age composition of returning hatchery-origin PIT-tagged summer and fall-run chinook released in the Yakima subbasin as subyearling or yearling fish.	49
Table 23. Marine and freshwater recoveries of CWTs from brood year 1997-2014 releases of spring Chinook from the CESRF as reported to the Regional Mark Information System (RMIS).	55
Table 24. Estimated run size, harvest, and harvest rates of Yakima Basin spring Chinook in Columbia River mainstem and terminal area fisheries, 1983-present.	56
Table 25. Spring Chinook harvest in the Yakima River Basin, 1983-present.	59
Table 26. Estimated fall Chinook return, escapement, and harvest in the Yakima River, 1998-2019..	60
Table 27. Estimated Coho return, escapement, and harvest in the Yakima River, 1999-2019.....	60
Table 28. Escapement (Roza Dam counts less brood-stock collection and harvest above Roza) of natural- (NoR) and hatchery-origin (HoR) spring Chinook to the upper Yakima subbasin, 1982 – present.	69

Table 29. Avian predation river reach survey start and end locations and total reach length.....	71
Table 30. Yakima River Avian Predators.....	72
Table 31. Estimated consumption in 2019 by Avian species at three spring Chinook Salmon acclimation sites.....	80
Table 32. Fish Predator Survey River Miles and Distances.....	82
Table 33. Yakima River Fish Species.....	83

List of Figures

Figure 1. Yakima River Basin and Yakama Nation/YKFP-related artificial production and monitoring facilities (map provided by Paul Huffman).....	11
Figure 2. Estimated counts of natural- and Cle Elum Supplementation and Research Facility (CESRF-) origin spring Chinook (adults and jacks) at Prosser Dam, 1982-present.....	13
Figure 3. Estimated returns of adult and jack summer- and fall-run Chinook to the Yakima River mouth, 1983-present.....	14
Figure 4. Estimated counts of marked (presumed hatchery-origin) and unmarked (presumed natural-origin) Coho (adults and jacks) at Prosser Dam 1986-present... ..	14
Figure 5. Estimated counts of natural- and Cle Elum Supplementation and Research Facility (CESRF-) origin spring Chinook (adults and jacks) at Roza Dam, 1982-present.....	15
Figure 6. Average daily passage of Chinook and Coho (adults and jacks) at Prosser Dam, 2011-2020.....	15
Figure 7. Passage timing of adult and jack Chinook at Prosser Dam in 2020 by run (see Methods).....	16
Figure 8. Upper Yakima wild/natural spring Chinook return rate per spawner, before (brood years 1984-2000) and after (brood years 2001-2015) commencement of supplementation.....	20
Figure 9. Naches subbasin spring Chinook return rate per spawner, before (brood years 1984-2000) and after (brood years 2001-2015) commencement of supplementation in the Upper Yakima River.....	20
Figure 10. Productivity indices for age-3 natural-origin coho, brood years 2001-2017... ..	23
Figure 11. Box plot showing the 5-year average survival probabilities of natural-origin (Natural) and hatchery-origin (Hatchery) spring Chinook Salmon smolts.	31
Figure 12. Average annual survival rate (release to McNary Dam) of juvenile Summer Chinook smolts migrating from 2010 through 2020.....	32
Figure 13. Redd Counts upstream of Prosser Dam in the Yakima River Basin by species, 1981-present.....	39
Figure 14. Teanaway River Spring Chinook redd counts, 1981-2020 (vertical lines denote pre- and post-supplementation periods) and the proportion of natural-origin (NO) carcasses observed in intensive spawning ground surveys, 2002-2010.....	41
Figure 15. Distribution of summer and fall run Chinook redds in the Yakima River Basin (above Prosser Dam) based on redd observations from 2014 to 2018.....	41
Figure 16. Fall Chinook redd counts above and below Prosser Dam, 1961-present, for years in which surveys were conducted and data are available.....	42
Figure 17. Distribution of coho redds in the Yakima River Basin.....	42

Figure 18. Adult return timing at Prosser Dam of PIT-tagged summer- and fall-run Chinook reared at the Marion Drain and Prosser Hatcheries and released as subyearlings, pooled for return years 2009-2018.	50
Figure 19. Overall Fine Sediment (<0.85mm) Trends with 95% confidence bounds in the Little Naches River Drainage, 1992-2020.	52
Figure 20. Fine Sediment Trends in the South Fork Tieton River, 1999-2015..	53
Figure 21. Overall average percent fine sediment (< 0.85 mm) in spawning gravels of the Upper Yakima River, 1997-2020.....	54
Figure 22. Distribution of coded-wire tag recoveries of Yakima Basin summer/fall run Chinook releases in marine, mainstem Columbia River, and Yakima Basin fisheries. Data retrieved from the regional mark information system (RMIS) for brood year 1997-2007 recoveries.....	57
Figure 23. Map of the Yakima River Basin, Cle Elum Supplementation and Research Facility (CESRF) locations, and timeline of the spring Chinook supplementation program.	63
Figure 24. Spring Chinook redd counts in the supplemented Upper Yakima (red bar) relative to the un-supplemented Naches (control; blue bar) for the pre- (1981-2000) and post-supplementation (2001-2020) periods.....	63
Figure 25. Natural-Origin returns of Spring Chinook in the supplemented Upper Yakima (red bar) relative to the un-supplemented Naches (control; blue bar) for the pre- (1982-2004) and post-supplementation (2005-2020) periods.....	64
Figure 26. Hypothetical outcomes of trait divergence (domestication effects) over time for a segregated (hatchery-control or HC) line of fish, compared to an integrated (supplementation or S) line of fish and a wild (wild-control or WC) line of fish	67
Figure 27. Estimated genetic divergence (variation) for integrated (INT blue), segregated (SEG red), and wild founder (black) spring Chinook in the CESRF program after 4 parental-generations of the hatchery program (P1=1998, F1=2002, F2=2006, F3=2010, F4=2014; updated from Figure 4 in Waters et al. 2015).	70
Figure 28. Avian Predator Survey Locations.....	73
Figure 29. Avian Predator Totals by Reach.....	75
Figure 30. Parker Reach Total Avian Predators by Week.....	75
Figure 31. Parker Reach Avian Predator Species Counts.....	76
Figure 32. Granger Reach Avian Predator Species Counts.....	76
Figure 33. Below Prosser Avian Predator Species Counts.....	77
Figure 34. Benton Reach Avian Predator Species Counts.....	77
Figure 35. Lower Yakima Reach Avian Predator Species Counts.....	78
Figure 36. Avian Predator Counts at Chandler “hotspot”.....	79
Figure 37. Avian Predator Counts at Wanawish Dam “hotspot”.....	79
Figure 38. Fish Predator Survey Locations.....	82
Figure 39. Fish Predator Counts by Reach and Species.....	84
Figure 40. Parker Reach Fish Counts by Species.....	85
Figure 41. Granger Reach Fish Counts by Species.....	85
Figure 42. Above Prosser Dam Fish Counts by Species.....	86
Figure 43. Below Prosser Dam Fish Counts by Species.....	86
Figure 44. Benton Reach Fish Counts by Species.....	87
Figure 45. Lower Yakima Reach Fish Counts by Species.....	87

Figure 46. Total Count of Fish Predators below Prosser Dam.	88
Figure 47. Adult Smallmouth Bass Totals by Reach.	89
Figure 48. Adult and Juvenile Smallmouth Bass Total below Prosser Dam.	90

Acknowledgements

Monitoring and evaluation efforts in the Yakima River Basin are the result of a cooperative effort by many individuals from a variety of agencies including the Yakama Nation Fisheries Program, the Washington Department of Fish and Wildlife, the United States Fish and Wildlife Service, the Columbia River Inter-Tribal Fish Commission, the National Oceanic and Atmospheric Administration Fisheries department as well as some consultants and contractors. We also wish to acknowledge and thank the Yakama Nation Tribal Council, U.S. Bureau of Reclamation, Yakima Subbasin Fish and Wildlife Recovery Board, Pacific States Marine Fisheries Commission, U.S. Forest Service Naches Ranger District, University of Idaho, Lars Mobrand and associates, University of Washington, and Central Washington University for their many contributions to this project including both recommendations and data services.

The core project team includes the following individuals: Joe Blodgett, Joe Hoptowitz, Mark Johnston, Bill Bosch, Todd Newsome, Melinda Davis, Michael Porter, Sara Sohappy, Chris Frederiksen, Shubha Pandit, Andrew Matala, Daylen Isaac, Jim Matthews, and a number of technicians from the YN; Charles Strom and a number of assistants from the Cle Elum Supplementation and Research Facility; Michael Fiander and the crew at the Prosser Hatchery; Anthony Fritts, Gabe Temple, Christopher Johnson, and a number of assistants from the WDFW; We also receive considerable support from: Sharon Lutz, Susan Gutenberger and assistants from the USFWS; Don Larsen, Andy Dittman, Charlie Waters and assistants from NOAA Fisheries; Zach Penney, Doug Hatch, and many others at the Columbia River Inter-Tribal Fish Commission; and Kerry Naish and others at the University of Washington. The following individuals are now retired but we acknowledge their many and invaluable contributions over the years: Dave Fast, David Lind, Bill Fiander, and Paul Huffman of the YN, Curt Knudsen from Oncorh Consulting and Doug Neeley from IntSTATS Consulting.

Special acknowledgement and recognition is owed to all of the dedicated Yakama Nation and YKFP personnel working on various tasks including Ray Decoteau, Joe Yallup, Joe Jay Pinkham III (retired), Leroy Senator (retired), Sy Billy, Wayne Smartlowit, Morales Ganuelas, Pharamond Johnson, Steve Salinas, Shiela Decoteau, Jimmy Joe Olney, Conan Northwind, Quincy Wallahee, Andrew Lewis, Nate Pinkham, Gene Sutterlick, Germaine Hart, Jamie Bill, Nate Pinkham, William Manuel, Terrance Compo, Levi Piel, Winna Switzler, Florence Wallahee, Ted Martin and Arnold Barney. The accomplishments and achievements documented here are a direct result of their dedication and desire to seek positive results for the betterment of the resource. Also, these achievements are attainable because of the efficient and

essential administrative support received from all of the office and administrative support personnel for the YKFP including: Rubi Rodriguez, Joey Estrada, Shirley Alvarado, and Adrienne Wilson.

We would especially like to thank former contributors to the Yakima/Klickitat Fisheries Project including Bruce Watson, Joel Hubble, Steve Schroder, Todd Pearsons, Craig Busack, Ray Brunson, Joy Evered, and Bill Hopley. These individuals put in countless hours of hard work during the planning, design, and implementation of this project. Their contributions helped to lay a solid foundation for this project and our monitoring and evaluation efforts. Dan Barrett (retired) served as the manager of the CESRF from 1997-2002. He helped to lay a solid foundation for the critical work done day in and day out at the Cle Elum facility.

This work is funded by the Bonneville Power Administration (BPA) through the Northwest Power and Conservation Council's (NPCC) Fish and Wildlife Program. Michelle O'Malley is BPA's contracting officer and technical representative (COTR) for this project. David Byrnes and Patricia Smith served as COTRs in the past and contributed substantially to the project over the years.

Executive Summary

The Yakima-Klickitat Fisheries Project (YKFP) is a joint project of the Yakama Nation (lead entity) and the Washington State Department of Fish and Wildlife (WDFW) and is sponsored in large part by the Bonneville Power Administration (BPA) with oversight and guidance from the Northwest Power and Conservation Council (NPCC). It is among the largest and most complex fisheries management projects in the Columbia Basin in terms of data collection and management, physical facilities, habitat enhancement and management, and experimental design and research on fisheries resources. The YKFP is attempting to evaluate all stocks historically present in the Yakima Subbasin and apply a combination of habitat restoration and hatchery supplementation or reintroduction, to restore the Yakima Subbasin ecosystem with sustainable and harvestable populations of salmon, steelhead and other at-risk species. This project and report address regional monitoring and evaluation strategies and sub-strategies as they apply to spring Chinook, summer/fall Chinook, and coho work in the Yakima Subbasin. This project (199506325) is related to numerous other projects in the Yakima Subbasin; additional information is available in the annual reports of these related projects.

The YKFP began a spring Chinook salmon hatchery program at the Cle Elum Supplementation and Research Facility (CESRF) near Cle Elum on the upper Yakima River in 1997. This program is a supplementation effort targeting the upper Yakima River population and is designed to test whether artificial propagation can be used to increase natural production and harvest opportunities while limiting ecological and genetic impacts. It is an integrated hatchery program because only natural-origin brood-stock is used and returning hatchery-origin adults are allowed to spawn in the wild. The program employs “best practice” hatchery management principles including reduced pond densities, strict disease management protocols, random brood-stock selection, and factorial mating to maximize effective population size. Fish are reared at the central facility, but released from three acclimation sites located near the central facility at: Easton approximately 25km upstream of the central facility, Clark Flat about 25km downstream of the central facility, and Jack Creek about 12km upstream from the Teanaway River’s confluence with the Yakima River. The CESRF collected its first spring Chinook brood-stock in 1997, released its first fish in 1999, and age-4 adults have been returning since 2001. The first generation of offspring of CESRF and wild fish spawning in the wild returned as adults in 2005. The program uses the adjacent, un-supplemented Naches River population as an environmental and wild control or reference system.

Adult returns of fall Chinook to the Yakima River Basin consist mostly of hatchery-origin fish returning from releases averaging 1.6 million Upriver Brights annually from

the Prosser Hatchery which have occurred since 1983. Summer-run Chinook were extirpated from the Yakima Basin by 1970. To increase the temporal and spatial distribution of summer/fall run Chinook in the Yakima River Subbasin, the program began releases of Wells Hatchery summer-run Chinook in the Yakima River Basin in 2009. Coho were extirpated from the Yakima Subbasin by the early 1980s. Pursuant to *U.S. v. Oregon* court-mandated agreements, substantial numbers (annual average > 700,000) of hatchery-reared coho salmon were released into the Yakima River since the mid-1980s. Prior to 1996 the primary purpose of releases was harvest augmentation and fish were released in sub-optimal spawning and rearing areas below Wapato Dam. With the inception of the YKFP in 1996, the objective of the coho program became “to determine the feasibility of reestablishing a naturally spawning coho population” and releases were moved upriver to more suitable habitats for natural coho.

Annual abundance of spring Chinook at Prosser Dam has increased from a 1982-2000 average of about 4,000 fish to a 2001-2020 average of about 9,500 fish. These increases can be attributed to returns from the Cle Elum supplementation program beginning in 2001, improved freshwater passage conditions, improved marine survival, and habitat restoration and enhancement work. Annual abundance of summer/fall Chinook at the Yakima River mouth has increased from a 1983-1999 average of about 1,200 fish to a 2000-2020 average of about 6,500 fish. While this increase coincides with improved ocean conditions, some of the increase may also be due to improved passage in the mainstem Columbia River, and improvements in spawning and rearing protocols. Approximately 250 summer-run Chinook were estimated to pass above Prosser Dam in 2020. The 2020/2021 adult passage over Prosser Dam was approximately 2,300 coho. An additional 970 adults returned directly into the Prosser Fish Hatchery. The hatchery is located approximately 1 mile below the dam and the returning adults are used for brood stock. Coho returns to Prosser averaged over 5,600 fish from 1998-2020 (an order of magnitude improvement from the average for years prior to the project) including estimated returns of wild/natural coho averaging over 800 fish annually since 2001.

Trends in adult productivity indices for Yakima Basin natural-origin spring Chinook appear to be very similar for both Upper Yakima and Naches populations. Trends in adult productivity indices for natural-origin coho are not as clear. Under present conditions, productivity for spring Chinook appears to peak at about 1,000 to 1,500 spawners and decline as spawner abundance approaches 2,000 fish or greater. These data indicate that density-dependent limiting factors depress natural productivity at fairly low population abundance in the Yakima River Basin. Until these factors are fully addressed, supplementation yields higher overall productivity rates and can be used to return adults to fisheries and to augment natural spawning populations.

For smolt migration years 2000 to present, annual abundance estimates of juvenile smolts migrating downstream at Prosser Dam averaged 209,900 wild/natural spring Chinook, 331,200 CESRF-origin spring Chinook, 44,000 wild/natural-origin coho, and 262,800 hatchery-origin coho. Preliminary smolt-to-adult survival indices averaged approximately 2.4% and 2.9% for natural-origin spring Chinook and coho, respectively. Because of many complexities associated with the production of smolt indices, these data are useful for analysis of trends but should not be used as direct citations of, or for comparisons of marked and unmarked, smolt-to-adult survival rates. Substantial juvenile mortality occurs as smolts migrate through the Yakima River system. Strategies have been proposed to address limiting factors and improve survival of emigrating Yakima Basin juveniles. As these strategies are implemented, we expect smolt and smolt-to-adult survival to improve.

Spatial distribution of spring Chinook spawners has increased as a result of acclimation site location, salmon homing fidelity and more fully seeding preferred spawning habitats. Spring Chinook redd counts in the Teanaway River increased from a pre-supplementation average of 3 redds per year to a post-supplementation average of 55 redds per year. Fall Chinook redd distribution in the Yakima River Basin appears to be experiencing a transition with an increasing proportion of redds observed above Prosser Dam in the most recent decade. This change is primarily attributed to substantial changes in lower Yakima River habitats in recent years. Redd counts and spatial distribution of coho have increased substantially in recent years, with over 200 redds enumerated annually on average in tributaries in the upper watersheds since 2004. In 2020/2021, 95 coho redds were observed in tributaries in the Naches and Upper Yakima Subbasins. Approximately 50 redds were found in the Naches River and over 70 were found in the mainstem Yakima River above Roza Dam.

Monitoring and evaluation of diversity metrics is primarily focused on the CESRF spring Chinook program in the Upper Yakima River. Generally, we have detected small, but significant differences between hatchery- and natural-origin fish in some juvenile and adult traits with many results already published in the peer-reviewed literature.

Overall average fine sediment levels in the Naches and Upper Yakima River subbasins over many years of sampling continue to trend downward.

We believe Yakima Basin spring Chinook contribute minimally to marine fisheries as their spatial and temporal ocean migration patterns do not appear to intersect with marine fisheries. However, Yakima Basin fall- and summer-run Chinook and coho do

contribute substantially to marine fisheries and to mainstem Columbia River fisheries from the mouth to the Hanford Reach area. Recreational spring Chinook fisheries have returned to the Yakima River Basin after a 40-year absence. This has contributed to improved relationships between all the Basin's stakeholders and increased opportunities for collaboration.

Supplementation has increased spring Chinook redd abundance in the Upper Yakima relative to the Naches control system. We observed an average increase in redd counts in the upper Yakima about 54% greater than that in the Naches system from the pre- to post-supplementation periods. Natural-origin returns of adult spring Chinook in the post-supplementation period (2005-2020) are trending downward relative to the pre-supplementation period (1982-2004) in both the Upper Yakima and Naches Rivers but the trend in the Naches control system is a steeper decline. After several generations of study, the results (many of which are published in the peer-reviewed literature) from the spring chinook supplementation program in the Upper Yakima River demonstrate that a well-designed and carefully managed integrated hatchery program using 100% natural-origin broodstock can produce fish for harvest and return fish to the natural spawning grounds with minimal negative impacts to the target ecosystem. Coho re-introduction research in the published literature suggests that hatchery-origin coho, with a legacy of as many as 10 to 30 generations of hatchery-influence, can reestablish a naturalized population after as few as 3 to 5 generations of outplanting in the wild.

YKFP efforts to monitor and evaluate hatchery reform focus on the CESRF spring Chinook program which was designed explicitly for this purpose from its inception. By designing the program to use only natural-origin fish for brood-stock, the program has demonstrated reduced genetic divergence for the integrated program compared to a traditional segregated hatchery program. The CESRF is also meeting or exceeding scientific recommendations for proportionate natural influence (PNI) on an annual basis with a 20-year mean annual PNI of 66%. The project is thus far meeting or exceeding most other established objectives related to hatchery reform.

Major piscivorous predators in the Yakima River Basin include: common mergansers, American white pelicans, double-crested cormorants, gulls, great blue herons, northern pike minnows, and smallmouth bass. The project has initiated efforts to control the pike minnow and smallmouth bass populations.

Project results are communicated broadly through the annual [science and management conference](#), technical reports and peer-reviewed journal publications (see references and project-related publications), and via several related web sites described in Appendix A.

Introduction

The Yakima-Klickitat Fisheries Project (YKFP) is a joint project of the Yakama Nation (lead entity) and the Washington State Department of Fish and Wildlife (WDFW) and is sponsored in large part by the Bonneville Power Administration (BPA) with oversight and guidance from the Northwest Power and Conservation Council (NPCC). It is among the largest and most complex fisheries management projects in the Columbia Basin in terms of experimental design and research on fisheries resources, physical facilities, habitat enhancement and restoration, and data collection and management. Consistent with [Wy-Kan-Ush-Mi Wah-Kish-Wit](#) (CRITFC 1995) and using principles of adaptive management (BPA 1996; Salafsky et al. 2001), the YKFP is attempting to evaluate all stocks historically present in the Yakima Subbasin and apply a combination of habitat restoration and hatchery supplementation or reintroduction, to restore the Yakima Subbasin ecosystem with sustainable and harvestable populations of salmon, steelhead and other at-risk species.

The original impetus for the YKFP resulted from the landmark fishing disputes of the 1970s, the ensuing legal decisions in *United States versus Washington* and *United States versus Oregon*, and the region's realization that lost natural production needed to be mitigated in upriver areas where these losses primarily occurred. The YKFP was first identified in the NPCC's 1982 Fish and Wildlife Program (FWP) and supported in the *U.S. v Oregon* 1988 Columbia River Fish Management Plan (CRFMP). A draft Master Plan was presented to the NPCC in 1987 and the Preliminary Design Report was presented in 1990. In both circumstances, the NPCC instructed the Yakama Nation, WDFW and BPA to carry out planning functions that addressed uncertainties in regard to the adequacy of hatchery supplementation for meeting production objectives and limiting adverse ecological and genetic impacts. At the same time, the NPCC underscored the importance of using adaptive management principles to manage the direction of the Project. The 1994 FWP reiterated the importance of proceeding with the YKFP because of the added production and learning potential the project would provide. The YKFP is unique in having been designed to rigorously test the efficacy of hatchery supplementation. Given the current depressed status of many salmon and steelhead stocks, and the heavy reliance on artificial propagation as a recovery tool, YKFP monitoring results have great region-wide significance.

Supplementation is envisioned as a means to enhance and sustain the abundance of wild and naturally-spawning populations at levels exceeding the cumulative mortality burden imposed on those populations by habitat degradation and by natural cycles in environmental conditions. A supplementation hatchery is properly operated as an

adjunct to the natural production system in a watershed. By fully integrating the hatchery with a naturally-producing population, high survival rates for the component of the population in the hatchery can raise the average abundance of the total population (hatchery component plus naturally-producing component) to a level that compensates for the high mortalities imposed by human development activities and fully seeds the natural environment. However, it is important to recognize that “rebuilding natural populations will ultimately depend on improving habitat quality and quantity” ([ISRP 2011](#), Venditti et al. 2017) of which habitat connectivity is an essential component (CRITFC 1995, Milbrink et al. 2011). Hatchery programs, even “state of the art” integrated supplementation programs designed to follow all of the best management practice recommendations (Cuenco et al. 1993, Mobernd et al. 2005), do not directly affect any of these habitat parameters which are vital to improving natural productivity. Therefore, the YKFP is working with partners in multiple forums to implement habitat restoration and water resource management projects designed to address factors limiting productivity (see Yakima [Subbasin Recovery](#), and [Integrated](#) plans).

The objectives of the YKFP are to: enhance existing stocks; re-introduce extirpated stocks; protect and restore habitat in the Yakima Subbasin; operate using a scientifically rigorous process that will foster application of the knowledge gained about hatchery supplementation and habitat restoration throughout the Columbia River Basin; and use Ecosystem Diagnosis and Treatment (EDT) and other modeling tools to facilitate planning for project activities. In strictly scientific terms the stated purpose of the project is, “to test the assumption that new artificial production can be used to increase harvest and natural production while maintaining the long-term genetic fitness of the fish population being supplemented and keeping adverse genetic and ecological interactions with non-target species or stocks within acceptable limits” (RASP 1992, BPA 1996). WDFW is addressing some critical uncertainties (see [Columbia River Basin Research Plan](#) and [Critical Uncertainties for the Columbia River Basin Fish and Wildlife Program](#)) related to genetic and ecological interactions under project [1995-064-25](#). We are working jointly with WDFW and CRITFC ([2009-009-00](#)) to address fish propagation, predation, harvest, and monitoring and evaluation methodology uncertainties including:

Fish Propagation Question 1. Are current propagation efforts successfully meeting harvest and conservation objectives while managing risks to natural populations?

1.2. Can hatchery production programs meet adult production and harvest goals (integrated and segregated) while protecting naturally spawning populations?

1.4. What is the magnitude of any demographic benefit or detriment to the production of natural-origin juveniles and adults from natural spawning of hatchery-origin supplementation adults?

1.5. What are the range, magnitude and rates of change of natural spawning fitness of integrated (supplemented) populations, and how are these related to management rules including the proportion of hatchery fish permitted on the spawning grounds, and the proportion of natural origin adults in the hatchery broodstock?

Predation Question 1. Are the current efforts to address predation and reduce numbers of predators effective?

Predation Question 2. Are there actions other than removing predators that could reduce predation on listed species?

Harvest Question 1. Do current harvest and escapement strategies provide the expected results in supporting recovery efforts and providing harvest opportunities?

Monitoring and evaluation methods Question 1. Are current methods to ... count fish and to measure productivity adequate to cost effectively inform decisions?

Monitoring and evaluation methods Question 2. Are there innovative methods for counting fish and measuring their productivity that would better inform decisions?

YKFP-related project research in the Yakima River Basin has resulted in the publication of over 60 manuscripts in the peer-reviewed literature (see References and Project-Related Publications). The status of ongoing research relative to the above uncertainties is presented as part of this report.

This report includes sections on the following regional research, monitoring, and evaluation (RME) strategies: fish population status, harvest, hatchery, and predation. Each section addresses all relevant sub-strategies that apply to this project. The report addresses these strategies and sub-strategies as they apply to spring Chinook (*Oncorhynchus tshawytscha*), summer/fall Chinook (*O. tshawytscha*), and coho (*O. kisutch*) RM&E work in the Yakima subbasin. Steelhead (*O. mykiss*) RME work is addressed in related VSP ([2010-030-00](#)), on-reservation watersheds ([1996-035-01](#)), and Kelt Reconditioning (CRITFC [2008-458-00](#) and [2007-401-00](#)) projects. WDFW is addressing hatchery uncertainties related to genetic and ecological interactions under project [1995-064-25](#). YKFP-related habitat activities for the Yakima Subbasin are addressed under projects [1997-051-00](#) and [1996-035-01](#) (except for sediment sampling

which is addressed here). Hatchery Production Implementation (O&M) is addressed under project [1997-013-25](#). **Data and findings presented in this report should be considered preliminary until results are published in the peer-reviewed literature.**

Study Area

The project study area is the Yakima River Basin [WRIA 37/38/39](#) (Figure 1).




Figure 1. Yakima River Basin and Yakama Nation/YKFP-related artificial production and monitoring facilities (map provided by Paul Huffman).

Fish Population Status Monitoring

Status and Trend of Adult Fish Populations (Abundance)

Methods: Adult salmon populations in the Yakima River Basin are enumerated at Prosser Dam using video equipment installed in all three adult fish ladders (monitoringresources.org methods 143, 144, 307, 418, 515). At both Prosser and Roza Dams, adult fish traps are also used on a seasonal basis for biological sampling and enumeration (monitoringresources.org methods 135). When the Roza adult trap is not in operation, video equipment is also employed at the adult fish ladders there. However, camera placement and actual viewing area are limited; these combined with water clarity issues during certain river conditions all affect video enumeration at Roza Dam. Automatic Passive Integrated Transponder (PIT) tag detectors are also employed at all fish ladders at both dams (see sites RZF and PRO in ptagis.org). For the safety and protection of personnel and equipment, video and PIT-detection equipment are removed during periods of high river flow. In these instances, biologists attempt to extrapolate fish counts using data from before and after the high flow event. Although adult passage over spillways is believed to occur when flows are favorable, Prosser Dam counts are generally considered by Yakama Nation biologists to be within +/- 5% of actual fish passage. Roza Dam counts during trap operation (generally the entire spring Chinook counting period, March-September) are considered virtually 100% accurate; however, during the late fall and winter counting period when video equipment is used at least part of the time, accuracy may fall to only 50-75% of actual fish passage based on preliminary evaluation of PIT tag detection data. Fish are denoted as hatchery- or natural-origin based on presence or absence respectively, of observed external or internal marks or tags (monitoringresources.org method 342). Chinook are denoted as spring-, summer-, or fall-run based on review of PIT-detection data and visual observations of coloration and body morphometry.

At Prosser Dam, time-lapse video recorders (VHS) and a video camera were used at viewing windows at each of the three fishways. Digital video recorders (DVR) and progressive scan cameras (to replace the VHS systems) were tested at each of the three Prosser fishways in 2007 and became fully operational in February of 2008. The new system functions very similarly to the VHS system but provides digital video data readily downloadable to the viewing stations in Toppenish. This new system also allows technicians in Toppenish to scan rapidly to images of fish giving more timely and accurate fish counts. The technicians review the images and record various types of data for each fish that migrates upstream via the ladders. The data are entered into a Microsoft Access database. Similarly, adult trap sample data for operations at both

Prosser and Roza Dams are entered into Microsoft Access databases. Post-season, counts are reviewed and adjusted for data gaps and knowledge about adult and jack lengths from sampling activities with corrections made to our master data sets. Daily dam count (including trap and video counts) reports and Yakima Basin adult trap sampling (login required) data for the Prosser and Roza data sets are available at: <https://yakamafish-nsn.gov/fish-data>.

Spring Chinook began returning from the Cle Elum Supplementation and Research Facility (CESRF) in 2000 (jacks) and 2001 (adults). All CESRF-origin spring Chinook are marked. Due to physical and logistical constraints at the Prosser Hatchery it is not possible to mark all hatchery releases of summer/fall run Chinook without jeopardizing fish health and survival but these issues are being addressed through the Master Planning process (Yakama Nation 2019). Thus, enumeration of hatchery- and natural-origin summer/fall run Chinook adult returns is not presently available but will be available in the future. New marking protocols made it possible to distinguish hatchery- and natural-origin coho beginning with return year 2001.

Results:




Figure 2. Estimated counts of natural- and Cle Elum Supplementation and Research Facility (CESRF-) origin spring Chinook (adults and jacks) at Prosser Dam, 1982-present.




Figure 3. Estimated returns of adult and jack summer- and fall-run Chinook to the Yakima River mouth, 1983-present.




Figure 4. Estimated counts of marked (presumed hatchery-origin) and unmarked (presumed natural-origin) Coho (adults and jacks) at Prosser Dam 1986-present.




Figure 5. Estimated counts of natural- and Cle Elum Supplementation and Research Facility (CESRF)-origin spring Chinook (adults and jacks) at Roza Dam, 1982-present.




Figure 6. Average daily passage of Chinook and Coho (adults and jacks) at Prosser Dam, 2011-2020.




Figure 7. Passage timing of adult and jack Chinook at Prosser Dam in 2020 by run (see Methods).

Discussion:

Annual abundance of spring Chinook at Prosser Dam has increased from a 1982-2000 average of about 4,000 fish to a 2001-2020 average of about 9,500 fish (Figure 2). Annual abundance of spring Chinook at Roza Dam has increased from a 1982-2000 average of about 2,300 fish to a 2001-2020 average of approximately 6,400 fish (Figure 5). These increases beginning in 2001 coincide with the first adult returns from the Cle Elum supplementation program. However, freshwater passage conditions, marine survival, and habitat restoration and enhancement work also affect survival and return rates. The lower adult returns observed in 2003 and 2007 coincide with notable droughts during the corresponding smolt outmigration years of 2001 and 2005. Returns in several recent years (beginning in 2015) were affected by thermal barriers in the lower Yakima River during the adult migration timeframe. Discussion of uncertainties relating to the Cle Elum spring Chinook supplementation program is included under Hatchery Monitoring later in this report. Additional data and detail on the Cle Elum spring Chinook supplementation program and the status of natural- and CESRF-origin spring Chinook in the Yakima River Basin are provided in Appendix B.

Although some natural production is occurring, adult returns of fall Chinook to the Yakima River Basin consist mostly of hatchery-origin fish returning from annual releases of Upriver Brights from the Prosser Hatchery which have occurred since 1983 and averaged about 1.9 million since 1999 (Yakama Nation 2019). In addition, the Yakama Nation has a goal of re-establishing Summer-run Chinook which were

extirpated from the Yakima Basin by 1970. Pursuant to this goal we began releases of Wells Hatchery summer-run Chinook in the Yakima River Basin in 2009. Annual abundance of summer/fall Chinook at the Yakima River mouth has increased from a 1983-1999 average of about 1,200 fish to a 2000-2020 average of about 6,500 fish (Figure 3). While this increase coincides with improved ocean conditions, some of the increase may also be due to improved passage in the mainstem Columbia River, and improvements in spawning and rearing protocols. By re-establishing the summer-run component we seek to increase the temporal (Figures 6 and 7) and spatial distribution of summer/fall run Chinook in the Yakima River Subbasin (Yakama Nation 2019). Approximately 250 summer-run Chinook were estimated to pass above Prosser Dam in 2020 (Figure 7).

Coho were extirpated from the Yakima Subbasin by the early 1980s. Pursuant to *U.S. v. Oregon* court-mandated agreements, substantial numbers (annual average > 700,000) of hatchery-reared coho salmon were released into the Yakima River since the mid-1980s. Prior to 1996 the primary purpose of releases was harvest augmentation and fish were released in sub-optimal spawning and rearing areas below Wapato Dam. With the inception of the YKFP in 1996, the objective of the coho program became “to determine the feasibility of reestablishing a naturally spawning coho population” and releases were moved upriver to more suitable habitats for natural coho. Monitoring of these efforts to re-introduce a sustainable, naturally spawning coho population in the Yakima Basin have indicated that coho returns averaged 5,600 fish from 1998-2020 (an order of magnitude improvement from the average for years prior to the project) including estimated returns of wild/natural coho averaging over 800 fish annually since 2001 (Figure 4).

Status and Trend of Adult Productivity

Methods:

We used recruit-per-spawner relationships (Ricker 1975) to describe adult-to-adult productivity indices. Species-specific methods were as follows.

Spring Chinook

Estimated natural-origin spawners for the Upper Yakima River were calculated as the estimated escapement above Roza Dam plus the estimated number of spawners between the confluence with the Naches River and Roza Dam. Total natural-origin returns to the Upper Yakima River were developed using run reconstruction techniques (Appendix B). Age composition for Upper Yakima returns was estimated from spawning ground carcass scale samples (monitoring resources.org method [112](#)) for the years 1982-1996 and from Roza Dam brood-stock collection samples (Knudsen et al. 2006; Appendix B) for the years 1997 to present. Since age-3 fish

(jacks) are not collected for brood-stock in proportion to the jack run size, the proportion of age-3 fish in the upper Yakima for 1997 to present was estimated using the proportion of jacks (based on visual observation) counted at Roza Dam relative to the total run size.

Estimated spawners and total returns for Naches River Subbasin natural-origin spring Chinook were calculated using run reconstruction techniques (Appendix B). Age composition for Naches Basin age-4 and age-5 returns were estimated from spawning ground carcass scale samples (monitoring resources.org method [112](#)). The proportion of age-3 fish was estimated after reviewing jack count (based on visual observations) data at Prosser and Roza dams.

Estimated spawners at the CESRF were the total number of wild/natural fish collected at Roza Dam and taken to the CESRF for production brood-stock (Knudsen et al. 2006; Appendix B). Total returns of CESRF-origin fish were based on run reconstruction and Roza dam sampling operations. Age composition for CESRF fish was estimated using scales and PIT tag detections from CESRF fish sampled passing upstream through the Roza Dam adult monitoring facility (Knudsen et al. 2006; Appendix B).

Coho

From central British Columbia south, the vast majority of coho salmon adults are 3-year-olds, having spent approximately 18 months in fresh water and 18 months in salt water (Loeffel and Wendler 1968, Wright 1970). Therefore, we estimated a natural-origin productivity (recruits per spawner) index by dividing natural-origin returns to Prosser Dam by the estimated returns to Prosser Dam three years prior. We computed this index for both adult and combined adult and jack returns per adult and combined adult and jack spawner. Note that this method will bias productivity estimates high, as it assumes no natural production from hatchery-origin spawners.

Summer/Fall Run Chinook

Adult fall Chinook returning to the Yakima Basin consist of hatchery-origin returns from releases at and above Prosser Dam and natural-origin returns from fish spawning naturally in the Yakima River. Due to fiscal, physical, logistical, and policy considerations, only a small proportion of hatchery-origin releases have been externally marked. Therefore, it is impossible at present to know the origin of unmarked adult fall Chinook counted at Prosser. Additional marking is proposed for hatchery-origin releases as part of the Master Plan (Yakama Nation 2019), which will allow development of a comprehensive brood/cohort age at return table for natural- and hatchery-origin returns. Methods and results for evaluating adult productivity of summer/fall run Chinook will be included in future reports and publications as the data become available.

Results:

Table 1. Adult-to-adult productivity indices for upper Yakima wild/natural spring Chinook.

Brood Year	Estimated Spawners	Estimated Yakima R. Mouth Returns			Total	Returns/ Spawner
		Age-3	Age-4	Age-5		
1984	1,715	92	1,348	139	1,578	0.92
1985	2,578	114	2,746	105	2,965	1.15
1986	3,960	171	2,574	149	2,893	0.73
1987	2,003	53	1,571	109	1,733	0.87
1988	1,400	53	3,138	132	3,323	2.37
1989	2,466	68	1,779	9	1,856	0.75
1990	2,298	79	566	0	645	0.28
1991	1,713	9	326	22	358	0.21
1992	3,048	87	1,861	95	2,043	0.67
1993	1,925	66	1,606	57	1,729	0.90
1994	573	60	737	92	890	1.55
1995	364	59	1,036	129	1,224	3.36
1996	1,657	1,059	12,882	630	14,571	8.79
1997	1,204	621	5,837	155	6,613	5.49
1998	390	434	2,803	145	3,381	8.68
1999	1,021 ¹	164	722	45	930	0.91
2000	11,864	856	7,689	127	8,672	0.73
2001	12,087	775	5,074	222	6,071	0.50
2002	8,073	224	1,875	148	2,247	0.28
2003	3,341	158	1,036	63	1,257	0.38
2004	10,377	207	1,547	75	1,828	0.18
2005	5,713	293	2,630	14	2,936	0.51
2006	3,378	868	2,887	133	3,888	1.15
2007	2,322	456	3,976	65	4,498	1.94
2008	4,343	1,135	3,410	123	4,668	1.07
2009	7,056	283	2,572	109	2,964	0.42
2010	8,383	923	3,854	59	4,836	0.58
2011	8,584	832	3,908	144	4,883	0.57
2012	5,483	197	2,445	20	2,662	0.49
2013	4,984	299	1,622	36	1,957	0.39
2014	6,751	241	814	12	1,067	0.16
2015	5,466	66	620	14 ²	701 ²	0.13 ²
2016	4,281	99	905 ²			
2017	3,342	75 ²				
2018	1,817					
2019	1,508					
2020	1,664 ²					
Mean	4,031	329	2,679	106	3,117	1.43

1. The geometric mean jack (age-3) proportion of spawning escapement from 1999-2020 was mean 0.17.

2. Preliminary.




Figure 8. Upper Yakima wild/natural spring Chinook return rate per spawner, before (brood years 1984-2000) and after (brood years 2001-2015) commencement of supplementation.




Figure 9. Naches subbasin spring Chinook return rate per spawner, before (brood years 1984-2000) and after (brood years 2001-2015) commencement of supplementation in the Upper Yakima River.

Table 2. Adult-to-adult productivity indices for Naches River Subbasin wild/natural spring Chinook.

Brood Year	Estimated Spawners	Estimated Yakima R. Mouth Returns					Returns/ Spawner
		Age-3	Age-4	Age-5	Age-6	Total	
1984	383	110	706	564	0	1,381	3.60
1985	683	132	574	396	0	1,102	1.61
1986	2,666	68	712	499	15	1,294	0.49
1987	1,162	27	183	197	0	407	0.35
1988	1,340	32	682	828	0	1,542	1.15
1989	992	28	331	306	0	665	0.67
1990	954	24	170	74	0	269	0.28
1991	706	7	37	121	57	222	0.31
1992	852	29	877	285	0	1,191	1.40
1993	1,145	45	593	372	0	1,010	0.88
1994	474	14	164	164	0	343	0.72
1995	124	40	164	251	0	455	3.66
1996	887	179	3,983	1,620	0	5,782	6.52
1997	762	207	3,081	708	0	3,996	5.24
1998	503	245	1,460	1,128	0	2,833	5.63
1999	358 ¹	113	322	190	0	626	1.75
2000	3,862	71	2,060	215	0	2,346	0.61
2001	3,912	126	1,254	471	0	1,850	0.47
2002	1,861	59	753	153	0	965	0.52
2003	1,400	52	237	175	0	464	0.33
2004	2,197	107	875	218	0	1,199	0.55
2005	1,439	167	653	116	0	936	0.65
2006	1,163	192	838	254	0	1,283	1.10
2007	463	125	1,649	514	0	2,288	4.94
2008	1,074	414	827	290	0	1,531	1.42
2009	903	84	448	65	0	597	0.66
2010	1,024	209	653	198	0	1,059	1.03
2011	1,942	137	1,088	305	0	1,530	0.79
2012	1,110	64	419	260	0	743	0.67
2013	750	110	660	148	0	919	1.23
2014	746	142	376	13	0	532	0.71
2015	1,285	26	34	206 ²		266 ²	0.21 ²
2016	790	6	523 ²				
2017	971	32 ²					
2018	500						
2019	51						
2020	740 ²						
Mean	1,140	101	830	353	3	1,301	1.57

1. The geometric mean jack (age-3) proportion of spawning escapement from 1999-2020 was 0.09.

2. Preliminary.

Table 3. Adult-to-adult productivity indices for Cle Elum SRF spring Chinook.

Brood Year	Estimated Spawners	Estimated Yakima R. Mouth Returns			Total	Returns/ Spawner
		Age-3	Age-4	Age-5		
1997	261	741	7,753	176	8,670	33.22
1998	408	1,242	7,939	602	9,782	23.98
1999	738 ¹	134	714	16	864	1.17
2000	567	1,103	3,647	70	4,819	8.50
2001	595	396	845	9	1,251	2.10
2002	629	345	1,886	69	2,300	3.66
2003	441	121	800	12	932	2.11
2004	597	805	3,101	116	4,022	6.74
2005	510	1,305	3,052	21	4,378	8.58
2006	419	3,038	5,812	264	9,114	21.75
2007	449	1,277	5,174	108	6,558	14.61
2008	457	2,344	4,567	65	6,976	15.27
2009	486	461	2,663	58	3,181	6.55
2010	336	1,495	3,183	30	4,707	14.01
2011	377	1,233	2,340	34	3,607	9.57
2012	374	221	1,492	10	1,723	4.61
2013	398	802	1,993	0	2,795	7.02
2014	384	1,008	1,447	7	2,463	6.41
2015	442	314	877	0 ²	1,191 ²	2.70 ²
2016	376	287	771 ²		1,058 ²	2.81 ²
2017	382	349 ²				
2018	294					
2019	306					
2020	405 ²					
Mean	443	906	3,003	88	4,020	6.84 ³

1. 357 or 48% of these fish were jacks.
2. Preliminary.
3. Geometric mean.

Table 4. Estimates of adult-to-adult productivity indices for Yakima Basin natural-origin coho.

Return Year	Prosser Dam Counts		Return per Spawner Indices	
	Adults	Jacks	With Jacks	Without Jacks
2001	1,432	21		
2002	309	245		
2003	1,523	135		
2004	1,820	25	1.27	1.27
2005	472	120	1.07	1.53
2006	1,562	114	1.01	1.03
2007	1,049	32	0.59	0.58
2008	459	587	1.77	0.97
2009	982	173	0.69	0.63
2010	573	37	0.56	0.55
2011	802	24	0.79	1.75
2012	550	33	0.50	0.56
2013	424	79	0.83	0.74
2014	1,082	18	1.33	1.35
2015	362	9	0.64	0.66
2016	103	45	0.29	0.24
2017	1,162	15	1.07	1.07
2018	125	32	0.42	0.35
2019	301	8	2.09	2.92
2020	744	107	0.72	0.64
Mean	792	93	0.92	0.99




Figure 10. Productivity indices for age-3 natural-origin coho, brood years 2001-2017.

Discussion:

Recruit per spawner data for the Upper Yakima and Naches spring Chinook populations are highly correlated (Tables 1 and 2; Pearson’s correlation

coefficient=0.87) and analysis of variance indicates the means (\pm one standard error) in the 32-year data set are not different (Upper Yakima= 1.47 ± 0.39 ; Naches= 1.57 ± 0.31 ; $P=0.85$). Trends in adult productivity indices for Yakima Basin natural-origin spring Chinook are also very similar for both Upper Yakima (Figure 8) and Naches (Figure 9) populations. Under present conditions, productivity for spring Chinook appears to peak at about 1,000 to 1,500 spawners and declines as spawner abundance approaches 2,000 fish or greater (Figures 8-9). The trend in adult productivity indices for natural-origin coho (Figure 10) is not as obvious, and 2014 marked the first year that we observed high coho spawner escapements (when hatchery-origin spawning escapement is included) similar to those we have observed with spring Chinook in some recent years. These data indicate that density-dependent limiting factors (see YSFWPB 2004) depress natural productivity at fairly low population abundance in the Yakima River Basin, as is the case for most salmon populations throughout the Columbia River Basin (ISAB 2015). Until these factors are fully addressed, supplementation yields higher overall productivity rates and can be used to return adults to fisheries and to augment natural spawning populations (Table 3). While higher spawner abundances under present conditions do not yield increased adult production, these fish still contribute to more fully seeding available habitats, increased spatial and temporal diversity, and nutrient enhancement that should eventually lead to increased natural food supply and higher productivity in the future (NRC 1996, see especially pp. 368-369; Kiffney et al. 2014).

Status and Trend of Juvenile Abundance

Methods: The Yakama Nation releases a number of hatchery-origin smolts annually pursuant to *U.S. v Oregon* Management Agreements. Adult returns from these releases serve to mitigate for lost harvest opportunity (due to alteration of the Columbia River ecosystem and associated losses in natural production and productivity), to augment the number of fish spawning naturally (supplementation), or a combination of the two. Juveniles are released from many locations as yearlings or subyearlings depending on the goals of the specific programs. As these juveniles migrate downstream, they are mixed with naturally produced juveniles.

Above Prosser Dam, a portion of the river flow is diverted into the Chandler canal to generate electrical power and serve irrigation districts downstream. Juvenile fish are diverted into the Canal (and subsequently the Chandler juvenile monitoring facility-CJMF, Figure 1) at different rates depending on river and canal flow. Smolt sampling efforts at the CJMF near Prosser Dam were conducted annually from early winter through early summer corresponding with salmon smolt out-migrations. A portion of entrained salmon outmigrants (regulated by a timed gate) was manually counted and sampled for biological data on a daily basis and all PIT tagged fish were interrogated.

Sampling methods were described in Busack et al. (1997) and were consistent with monitoringresources.org methods 1562, 1563, 1595, and 1614.

Paired releases of PIT-tagged smolts were made in order to estimate the fish entrainment and canal survival rates in relation to river conditions and canal operations. For outmigration years 1999 through 2014, these data were used to generate a multi-variate river flow/canal entrainment relationship (D. Neeley 2010 and 2012a). Over a range of flow diversion rates, juvenile fish entrainment rates generally fit a logistic curve: at low diversion rates, the entrainment rate is lower than the diversion rate, and at high diversion rates the entrainment rate is higher than the diversion rate. In recent years it became difficult to adapt the model to higher winter and spring flows and to river channel changes, partly because at low diversion rates it was difficult to capture enough fish to get many point estimates of entrainment rate. The releases that were made, however, still tended to support a low entrainment rate relative to diversion rate at high river flows. For some years, Prosser smolt passage estimates produced by this model were outside of what were considered reasonable bounds (e.g., entrainment-based Prosser passage estimates approached or even exceeded known releases for hatchery-origin spring Chinook far upstream). This required us to reevaluate and change our methodology. The proportions of all PIT-tagged smolts released above Prosser and detected at mid-Columbia dams that were previously detected in the Chandler Canal bypass now serve as estimates of bypass-detection efficiency. Expanded Prosser passage estimates were then derived using the juvenile sample counts and detection efficiencies as described in Appendix C. These methods were generally consistent with monitoringresources.org methods 435, 623 and 1743.

Results and Discussion:

At the CESRF, the number of release groups and total number of spring Chinook released diverged from the facility goal of 810,000 smolts in some years. In brood year 1997, the Jack Creek acclimation facility was not yet complete and project policy and technical teams purposely decided to under-collect brood stock to allow a methodical testing of the new facility's operations with less risk to live fish, which resulted in the stocking of only 10 of the 18 raceways. In brood year 1998, the project did not meet facility release goals due to a biological specification that no more than 50% of returning wild fish be taken for brood stock. As a result, only 16 raceways were stocked with progeny of the 1998 brood. In the same year, raceway 4 at the Jack Creek acclimation site suffered mechanical failures causing loss of flow and reduced oxygen levels and resulted in the loss of approximately one-half the fish in this raceway prior to release. In the drought year of 2001, a large number of returning adults presented with high enzyme-linked immunosorbent assay (ELISA) levels of *Renibacterium salmoninarum*, the causative agent of bacterial kidney disease (BKD). The

progeny of these females were purposely destroyed. As a result, only nine raceways were stocked with fish. The project decided to use the fish from an odd raceway for a predator avoidance training sub-experiment (these fish were subsequently acclimated and released from the Easton acclimation site).

Table 5. CESRF total releases of Spring Chinook by brood year, treatment, and acclimation site.

Brood Year	Control ¹	Treatment ²	Acclimation Site ³			Total
			CFJ	ESJ	JCJ	
1997	207,437	178,611	229,290	156,758		386,048
1998 ⁴	284,673	305,010	221,460	230,860	137,363	589,683
1999	384,563	374,226	232,563	269,502	256,724	758,789
2000	424,554	409,731	285,954	263,061	285,270	834,285
2001 ⁵	183,963	186,273	80,782	39,106	250,348	370,236
2002	420,764	416,140	266,563	290,552	279,789	836,904
2003	414,175	410,517	273,377	267,711	283,604	824,692
2004 ⁶	378,740	406,708	280,598	273,440	231,410	785,448
2005	431,536	428,466	287,127	281,150	291,725	860,002
2006	351,063	291,732	209,575	217,932	215,288	642,795
2007	387,055	384,210	265,907	254,540	250,818	771,265
2008	421,290	428,015	280,253	287,857	281,195	849,305
2009	418,314	414,627	279,123	281,395	272,423	832,941
2010	395,455	399,326	264,420	264,362	265,999	794,781
2011	382,195	386,987	255,290	248,454	265,438	769,182
2012	401,059	401,657	256,732	276,210	269,774	802,716
2013	No Experiment		215,933	214,745	216,077	646,755
2014	337,548	347,682	232,440	226,257	226,533	685,230
2015	331,316	323,631	208,239	218,225	228,483	654,947
2016	339,816	329,392	230,490	218,676	220,042	669,208
2017	351,656	359,013	244,236	233,449	232,984	710,669
2018	322,219	320,201	213,833	206,619	221,968	642,420
2019	270,242	280,156	153,575	193,042	203,781	550,398
Mean	356,347	353,741	237,729	235,387	244,865	707,335

1. Brood years 1997-2001: Optimum Conventional Treatment (OCT). Brood Years 2002-2004: Normal (High) growth. Brood Years 2005-2012: Normal feed at Cle Elum or accl. sites.
2. Brood years 1997-2001: Semi-natural Treatment (SNT). Brood Years 2002-2004: Slowed (Low) growth. Brood Year 2005, 2007-2012: saltwater transition feed at accl. Sites; 2014: BioPro vs BioVIT. Brood Year 2006: EWS diet at CESRF through May 3, 2007.
3. CFJ=Clark Flat; ESJ=Easton; JCJ=Jack Creek.
4. At the Jack Creek acclimation site only 4 of 6 raceways were stocked, and raceway 4 suffered mechanical failures resulting in the loss of about 20,000 OCT (control) fish.
5. High BKD incidence in adult broodstock reduced production to just 9 ponds (Clark Flat 1-2, Jack Creek, and Easton). Easton ponds were used for predator avoidance trained (PAT) fish and a single Cle Elum pond was spread between 6 ponds at Easton with crowders used to simulate pond densities for fish at other acclimation sites. These releases were excluded from mean pond density calculations by treatment.
6. At the Jack Creek acclimation site raceway 3 suffered mechanical failures resulting in the loss of about 45,000 high-growth (control) fish.

Table 6. Total releases of Coho by brood year, life stage, and brood source.

Brood Year	Smolts			Parr		Local Brood		Total Smolts	
	UppYak	Naches	Prosser	UppYak	Naches	Smolts	Parr	Non-Local	Local
1997	436,000	1,257,000							1,693,000
1998	502,155	502,239							1,004,394
1999	498,872	429,318							928,190
2000	187,659	379,904							567,563
2001	263,288	357,530							620,818
2002	403,000	407,002							810,002
2003	313,207	291,494							604,701
2004	322,417	332,455							654,872
2005	338,127	554,784	50,000						942,911
2006	426,632	516,753	81,114						1,024,499
2007	358,412	440,783	219,098						1,018,293
2008	304,638	269,936	182,719	12,000	25,000	324,598	37,000	432,695	757,293
2009	407,184	341,414	245,455	13,000	12,000	610,423	25,000	383,630	994,053
2010	443,030	131,972	190,836	15,000	15,000	522,027	30,000	243,811	765,838
2011	311,102	359,067	322,100	365,035	73,572	992,269	438,607		992,269
2012	339,034	305,197	221,567	10,555	29,565	446,295	40,120	419,503	865,798
2013	353,139	373,072	367,382	9,000	18,232	524,967	27,232	568,626	1,093,593
2014	408,112	298,619	267,830	93,525	92,023	974,561	185,548		974,561
2015	141,000	141,000	204,358			204,358		282,000	486,358
2016	407,196	369,521	205,967			205,967		776,717	982,684
2017	438,331	267,211	470,000	114,141	138,624	641,589	252,765	533,953	1,175,542
2018			929,388	139,925	114,735	400,000	254,660	528,388	929,388
2019			897,233	3,000	3,000	354,000	6,000	543,243	
Mean ¹	355,277	285,701	375,404	77,518	52,175	516,755	129,693	471,257	1,426,306

¹ 2008-2019 average.

Table 7. Total releases of fall-run Chinook by release year and release site.

Release Year	Prosser On-Station Release				Billy's Pond ²	Stiles Pond ²	Marion Drain	Total Release
	LWH ¹	PRH ¹	Subyrl ²	Yrlng ²				
1997	1,694,861							1,694,861
1998	1,695,399							1,695,399
1999	1,690,000		192,000					1,882,000
2000	1,695,037		306,000			16,000		2,017,037
2001	1,699,136		427,753			12,000		2,138,889
2002	1,704,348		286,158			4,000		1,994,506
2003	1,771,129		365,409			18,000		2,154,538
2004	1,748,200		561,385			52,223		2,361,808
2005	1,700,000		466,000		75,000 ³	38,890	41,000	2,320,890
2006	1,683,664		130,002			118,835	2,000	1,934,501
2007	1,700,000 ⁴		50,000		5,000	75,000	15,731	1,845,731
2008	789,993		519,486 ⁵	1,833	11,308	72,296	5,253	1,400,169
2009	1,647,275		299,574	7,516			24,245	1,978,610
2010	1,680,045		290,282	12,167			22,945	2,005,439
2011	1,699,944	503,772	620,952	22,857				2,847,525
2012	1,200,000	405,000	269,633	19,432			72,258	1,966,323
2013	1,506,725		184,949	22,735				1,714,409
2014	1,542,702	379,970	445,347					2,368,019
2015	1,653,495	479,078	584,397					2,716,970
2016	1,593,090		562,472					2,155,562
2017	1,789,400		423,920	159,470				2,213,320
2018	1,638,300		328,620	208,660				1,966,920
2019			457,691	224,961				682,652
2020	1,701,369		696,937					2,398,306

1. Transfers from LWH=Little White Salmon NFH; PRH=Priest Rapids Hatchery.
2. Releases from local brood source adults collected at Prosser Dam or Hatchery.
3. Released from Edler Pond (approximately 2 miles downstream from Billy's Pond).
4. Of which approximately 500,000 were reared on-station at Prosser under accelerated growth conditions.
5. Of which approximately 5,400 were released from SKOV pond.

Table 8. Total releases¹ of summer-run Chinook by release year and release site.

Release Year	Prosser	Stiles Pond		Nelson		Roza	Total Release
		Subyrl	Yrlng	Springs	Wapatox		
2009		180,911					180,911
2010		200,747					200,747
2011			176,364	39,406			215,770
2012	98,300			98,803			197,103
2013				88,208		48,355	136,563
2014				179,901		74,980	254,881
2015	55,000			99,600		122,848	277,448
2016						37,000	37,000
2017	169,499					75,000	244,499
2018				44,000		30,000	74,000
2019	581,000			50,000	100,000	75,000	806,000
2020	932,843 ²			100,000	100,000	175,000	1,307,843

1. All fish released as subyearlings unless otherwise noted.
2. Includes Marion Drain facility acclimation

For smolt migration years 2000 to present, annual abundance estimates of juvenile smolts migrating downstream at Prosser Dam averaged 209,900 wild/natural spring Chinook, 331,200 CESRF-origin spring Chinook, 42,200 wild/natural-origin coho,

and 271,300 hatchery-origin coho (Table 9). These are the years for which our data and methods are considered most reliable. Juvenile passage estimates for earlier years are provided below under “Status and Trend of Juvenile Productivity”; however, the reader should be aware that we have less confidence in these data because we have refined data collection protocols and passage estimation methods over time. As the majority of fall Chinook smolt migrants are unmarked hatchery-origin fish, we provide only the gross abundance indices below under “Status and Trend of Juvenile Productivity”. The reader is cautioned to pay particular attention to the factors complicating estimates of juvenile abundance and productivity described under “Status and Trend of Juvenile Productivity”.

Table 9. Estimated smolt passage at Prosser Dam for Yakima Basin wild/natural and hatchery-origin spring Chinook and coho.

Brood Year	Smolt Migr. Year	Spring Chinook		Coho	
		Wild/Natural	Hatchery (CESRF)	Wild/Natural	Hatchery
1998	2000	199,416	303,688	37,359	331,503
1999	2001	148,460	281,256	40,605	134,574
2000	2002	467,359	366,950	19,859	155,814
2001	2003	308,959	154,329	9,092	139,135
2002	2004	169,397	290,950	18,787	148,810
2003	2005	134,859	236,443	31,631	204,728
2004	2006	133,238	300,508	8,298	204,602
2005	2007	99,341	351,359	18,772	260,455
2006	2008	120,013	265,485	40,170	416,708
2007	2009	237,228	415,923	23,858	496,594
2008	2010	220,950	382,878	33,408	341,145
2009	2011	304,322	442,564	22,908	333,891
2010	2012	258,106	391,446	17,667	244,503
2011	2013	365,486	372,079	56,947	483,122
2012	2014	263,266	408,222	159,642	337,988
2013	2015	125,150	332,715	20,757	134,084
2014	2016	185,442	403,938	227,163	233,374
2015	2017	208,929	273,248	12,031	108,570
2016	2018	131,489	290,644	38,451	299,535
2017	2019	175,427	319,579	41,696	246,178
2018	2020	151,265	371,069	8,057	442,881
	Mean	209,910	331,203	42,246	271,343

Status and Trend of Juvenile Migration Survival to McNary Dam

Methods: For all species, releases of PIT tagged smolts provided a means to estimate smolt survival to McNary Dam. For most releases, PIT-tag detectors were located in or near the exit(s) from the release sites (monitoringresources.org 1558) and allowed estimation of the number of PIT-tagged fish leaving the release sites. To estimate the survival of smolts detected leaving the release sites that eventually pass McNary Dam,

the proportion of PIT-tagged smolts detected leaving the release sites that were later detected at McNary Dam was divided by McNary Dam's detection efficiency. The estimated detection efficiency was the number of smolts detected passing dams downstream of McNary that were previously detected passing McNary divided by the total number of smolts passing the downstream dams, whether or not the smolts were previously detected at McNary. Our methods are described in detail in Appendix C and are generally consistent with Sandford and Smith (2002) and with monitoringresources.org methods 623 and 1536. We used weighted logistic or weighted least squares analysis of variance to analyze differences in survival metrics and indices between various release sites, years and treatments. Additional detail, results and discussion are provided in Appendices D (spring Chinook), E (coho), and F (summer-run Chinook). There were no PIT-tagged releases of fall-run Chinook in 2020; the latest results for this species were presented in Appendix G of Fiander et al. (2019).

Results and Discussion:

For spring Chinook, we compared survivals to McNary Dam of CESRF hatchery-and natural-origin PIT-tagged smolts released into the Roza Dam bypass and migrating downstream of Roza Dam contemporaneously on or after March 16. This date was selected because CESRF fish were not allowed to begin volitional emigration from the acclimation sites until March 15. Approximately 81% of natural-origin spring Chinook smolts PIT-tagged and released at Roza since 1999 migrated downstream of Roza Dam prior to March 16 (derived using queries of PTAGIS database 7/12/2013). Natural and hatchery-origin smolts contemporaneously migrating past Roza from March 16 on are referred to as "late" migrants. Survival from Roza Dam to McNary Dam was better for late-migrating natural-origin relative to hatchery-origin spring Chinook smolts and for late-migrating relative to early-migrating natural-origin smolts (Figure 11; Appendix D).

For coho, we estimated survival from acclimation site release to McNary Dam based on life stage, brood source, location, and timing of the releases (Appendix E). The average survival probability of Coho Salmon smolts from the release sites to McNary Dam in 2020 was 47.14 ± 5.78 %, which was higher than estimates for the past three years. The higher survival rate for 2020 migration is at least partly due to release location because most of the tagged smolts in 2020 were released at Prosser Dam in the lower Yakima River. In other years, smolts were also released from several locations upstream from Prosser Dam, with correspondingly lower survival rates. The survival rate of smolts to McNary Dam was higher for the Eagle Creek-stock releases ($52.6 \pm 7.4\%$) than for Yakima-origin release ($41.1 \pm 10.1\%$). In 2019 and 2020, there was no release of the Washougal stock. For the last 6 migration years (2015-2020), Coho parr were released at different locations from May to October. Release site-to-

McNary survival of the parr releases was higher for the population released in August ($14\% \pm 0.20$) and followed by the group of July releases ($3.1\% \pm 0.40$) and then June releases ($1\% \pm 0.4$).




Figure 11. Box plot showing the 5-year average survival probabilities of natural-origin (Natural) and hatchery-origin (Hatchery) spring Chinook Salmon smolts (S. Pandit, Appendix D). A. is the comparison of Late hatchery- and natural-origin smolts; and B. is the comparison between Early- and Late-migrating natural-origin smolts.

Juvenile survival rates to McNary Dam for summer-run Chinook varied by year over migration years from 2010 through 2020 (Figure 12). The highest average annual survival rate was in 2011 ($40.15\% \pm 1.94\%$) and the lowest was in 2015 ($0.73\% \pm 0.47\%$). For 2020, the average survival rate from the combined release locations to McNary Dam was $14.7\% \pm 2.5\%$, which was a substantial improvement over 2018-2019. The relationship between the average of May and June river flow measured below Prosser Dam and the annual survival rate (release location to McNary Dam from 2009 through 2020) was strong and statistically significant ($r^2=0.54$, $p=0.01$) indicating that survival rate was a function of river flow in May and June. Higher flow in these months results in higher survival of juvenile Summer Chinook outmigrants. We also found that the relationship of size to survival rate from Prosser to McNary dams was similar for April and May releases, but that releases in June depressed the Prosser-to-McNary survival rate over the entire range of fish sizes. A complete report of our study of juvenile outmigration survival of

Yakima Basin Summer Chinook to Prosser and McNary dams is provided in Appendix F.




Figure 12. Average annual survival rate (release to McNary Dam) of juvenile Summer Chinook smolts migrating from 2010 through 2020 (S. Pandit, Appendix F).

The data indicate that there are substantial sources of juvenile mortality limiting survival of smolts migrating from release sites in the Yakima River basin. The YKFP is working with partners in multiple forums to implement habitat restoration and water resource management projects that address factors limiting survival and productivity (see Yakima [Subbasin](#), [Recovery](#), and [Integrated](#) plans).

Status and Trend of Juvenile Productivity (smolt-to-adult returns)

Methods:

Smolt abundance passage estimates at Prosser and the methods used to derive them were described above. For spring Chinook, adult return estimates to the Yakima River mouth were derived using Prosser and Roza adult abundance and harvest data (described in other sections of this report and in Appendix B) and run reconstruction techniques (Appendix B). For coho, we used Prosser adult abundance.

Adult fall Chinook returning to the Yakima Basin consist of hatchery-origin returns from releases at and above Prosser Dam and natural-origin returns from fish spawning naturally in the Yakima River. Due to fiscal, physical, logistical, and policy considerations, only a small proportion of hatchery-origin releases have been externally marked. Therefore, it is impossible at present to know the origin of unmarked adult fall Chinook counted at Prosser. Additional marking is proposed for hatchery-origin releases as part of the Master Plan (Yakama Nation 2019). To derive rough smolt-to-adult return indices for fall Chinook, aggregate (marked and unmarked combined) smolt passage estimates for the age-3, -4, and -5 components for a given return year were averaged and the aggregate adult passage estimate for that return year was divided by this average smolt passage estimate. For example, the “Prosser Average Smolts” for adult return year 1988 is the average of marked and unmarked Prosser smolt estimates for juvenile migration years 1983-1985.

We also queried the PTAGIS database for PIT-tagged summer- and fall-run Chinook and Coho that were released in the Yakima Subbasin in recent years and produced McNary Dam juvenile (smolt) to Bonneville Dam adult SAR indices using juvenile detections at or downstream of McNary and adult detections at or upstream of Bonneville Dams.

Results:

Table 10. Estimated smolt passage at Chandler and smolt-to-adult return indices (Chandler smolt to Yakima R. mouth adult) for Yakima Basin wild/natural and CESRF-origin spring Chinook.

Brood Year	Smolt Migr. Year	Mean Flow ¹ at Prosser Dam	Estimated Smolt Passage at Chandler		CESRF smolt-to-smolt survival ³	Yakima R. Mouth Adult Returns ⁴		Smolt-to-Adult Return Index ⁴	
			Wild/Natural ²	CESRF Total		Wild/Natural ²	CESRF Total	Wild/Natural ²	CESRF Total
1983	1985	3421	146,952			5,198		3.5%	
1984	1986	3887	227,932			3,932		1.7%	
1985	1987	3050	261,819			4,776		1.8%	
1986	1988	2454	271,316			4,518		1.7%	
1987	1989	4265	76,362			2,402		3.1%	
1988	1990	4141	140,218			5,746		4.1%	
1989	1991	n/a	109,002			2,597		2.4%	
1990	1992	1960	128,457			1,178		0.9%	
1991	1993	3397	92,912			544		0.6%	
1992	1994	1926	167,477			3,790		2.3%	
1993	1995	4882	172,375			3,202		1.9%	
1994	1996	6231	218,578			1,238		0.6%	
1995	1997	12608	52,028			1,995		3.8%	
1996	1998	5466	491,584			21,151		4.3%	
1997	1999	5925	584,016	187,669	48.6%	12,855	8,670	2.2%	4.6%
1998	2000 ⁵	4946	199,416	303,688	51.5%	8,240	9,782	4.1%	3.2%
1999	2001	1321	148,460	281,256	37.1%	1,764	864	1.2%	0.3%
2000	2002	5015	467,359	366,950	44.0%	11,434	4,819	2.4%	1.3%
2001	2003	3504	308,959	154,329	41.7%	8,597	1,251	2.8%	0.8%
2002	2004	2439	169,397	290,950	34.8%	3,743	2,557	2.2%	0.9%
2003	2005	1285	134,859	236,443	28.7%	2,746	1,020	2.0%	0.4%
2004	2006	5652	133,238	300,508	38.3%	2,802	4,482	2.1%	1.5%
2005	2007	4551	99,341	351,359	40.9%	4,295	5,004	4.3%	1.4%
2006	2008	4298	120,013	265,485	41.3%	6,004	10,577	5.0%	4.0%
2007	2009	5784	237,228	415,923	53.9%	7,952	7,604	3.4%	1.8%
2008	2010	3592	220,950	382,878	45.1%	7,385	8,036	3.3%	2.1%
2009	2011	9414	304,322	442,564	53.1%	3,766	3,606	1.2%	0.8%
2010	2012	8556	258,106	391,446	49.3%	6,602	5,592	2.6%	1.4%
2011	2013	4875	365,486	372,079	48.4%	7,343	4,160	2.0%	1.1%
2012	2014	4923	263,266	408,222	50.9%	3,969	1,932	1.5%	0.5%
2013	2015	1555	125,150	332,715	51.4%	3,415	3,139	2.7%	0.9%
2014	2016	5765	185,442	403,938	58.9%	1,800	2,865	1.0%	0.7%
2015	2017	7804	208,929	273,248	41.7%	1,171	1,319	0.6%	0.5%
2016	2018 ⁶	5652	131,489	290,644	43.4%	1,724 ⁶	1,220 ⁶	1.3% ⁶	0.4% ⁶
2017	2019 ⁶	3595	175,427	319,579	45.0%				
2018	2020 ⁶	2850	151,265	371,069	57.8%				

1. Mean flow (cfs) approaching Prosser Dam March 29-July 4 of juvenile migration year. In high flow years (flows at or > 5000 cfs) operation of the Chandler smolt sampling facility may be precluded during portions of the outmigration. Data courtesy of [U.S. BOR hydromet](#).
2. Aggregate of Upper Yakima, Naches, and American wild/natural populations.
3. Estimated smolt-to-smolt (release from upper Yakima River acclimation sites to Chandler) survival for CESRF juveniles.
4. Includes combined age-3 through age-5 returns. CESRF adult returns and smolt-to-adult survival values are understated relative to wild/natural values since these figures are not adjusted for differential harvest rates in mark selective fisheries in marine and lower Columbia River fisheries.
5. Available data were not sufficient to estimate juvenile flow-entrainment and passage of wild/natural fish.
6. Data for most recent year are preliminary; return data do not include age-5 adult fish.

Table 11. Average combined hatchery- and natural-origin smolt counts at Prosser for fish returning at age-3, -4, and -5, combined adult returns to Prosser Dam of all age classes, and estimated Prosser smolt-to-adult return indices for Yakima River fall-run Chinook for adult return years 1988-2020.

Adult Return Year	Prosser Average Smolts ¹	Prosser Total Adults	Prosser Smolt-to-Adult Return Index (SAR)
1988	1,029,429	224	0.02%
1989	1,469,019	670	0.05%
1990	1,664,378	1,504	0.09%
1991	1,579,989	971	0.06%
1992	1,811,088	1,612	0.09%
1993	2,034,865	1,065	0.05%
1994	1,976,301	1,520	0.08%
1995	1,329,664	1,322	0.10%
1996	1,023,053	1,392	0.14%
1997	1,097,032	1,120	0.10%
1998	1,533,093	1,148	0.07%
1999	1,786,511	1,896	0.11%
2000	1,716,156	2,293	0.13%
2001	1,867,966	4,311	0.23%
2002	1,946,676	6,241	0.32%
2003	2,108,238	4,875	0.23%
2004	2,653,056	2,947	0.11%
2005	2,707,132	1,942	0.07%
2006	2,724,824	1,528	0.06%
2007	2,312,562	1,132	0.05%
2008	2,450,308	2,863	0.12%
2009	2,353,675	2,972	0.13%
2010	2,118,702	2,888	0.14%
2011	1,780,670	2,718	0.15%
2012	1,806,572	4,477	0.25%
2013	1,939,754	7,706	0.40%
2014	2,411,076	7,792	0.32%
2015	2,476,483	7,380	0.30%
2016	2,436,111	5,355	0.22%
2017	2,348,973	1,613	0.07%
2018	2,527,520	763	0.03%
2019	2,544,821	691	0.03%
2020	2,422,840	1,724	0.07%
Mean	1,999,653	2,687	0.13%

¹ Average combined hatchery- and natural-origin smolt counts for the years which would comprise the age-3, -4, and -5 adult return components for each adult return year. For example, the “Prosser Average Smolts” for adult return year 1988 is the average of hatchery- and natural-origin Prosser smolt estimates for juvenile migration years 1983-1985.

Table 12. Preliminary estimates of Prosser-to-Prosser smolt-to-adult survival (SAR) indices for adult returns from hatchery- and natural-origin coho for the Yakima reintroduction program, juvenile migration years 2000-2019.

Juvenile Migration Year	Hatchery-origin			Natural-origin		
	Chandler Smolts ^a	Prosser Adults ^b	SAR Index	Chandler Smolts ^a	Prosser Adults ^b	SAR Index
2000	331,503	3,546	1.1%	37,359	1,432	3.8%
2001	134,574	166	0.1%	40,605	309	0.8%
2002	155,814	669	0.4%	19,859	1,523	7.7%
2003	139,135	505	0.4%	9,092	1,820	20.0%
2004	148,810	2,418	1.6%	18,787	472	2.5%
2005	204,728	2,898	1.4%	31,631	1,562	4.9%
2006	204,602	2,404	1.2%	8,298	1,049	12.6%
2007	260,455	4,131	1.6%	18,772	459	2.4% ^c
2008	416,708	8,835	2.1%	40,170	982	2.4% ^c
2009	496,594	5,153	1.0%	23,858	573	2.4% ^c
2010	341,145	7,216	2.1%	33,408	802	2.4% ^c
2011	333,891	4,948	1.5%	22,908	550	2.4% ^c
2012	244,503	2,703	1.1%	17,667	424	2.4%
2013	483,122	24,178	5.0%	56,947	1,082	1.9%
2014	337,988	2,943	0.9%	159,642	362	0.2%
2015	134,084	3,280	2.4%	20,757	103	0.5%
2016	233,374	2,693	1.2%	227,163	1,162	0.5%
2017	108,570	2,083	1.9%	12,031	125	1.0%
2018	299,535	3,566	1.2%	38,451	301	0.8%
2019	246,178	2,530	1.0%	41,696	744	1.8%
Mean	262,766	4,439	1.5%	43,955	792	2.9% ^d

^a Yakama Nation estimates of coho smolt passage at Chandler.

^b Yakama Nation estimates of age-3 coho returns to Prosser Dam for this juvenile migration cohort.

^c Average estimate derived from PIT-tag detections of Taneum Creek natural coho for juvenile migration years 2009-2011.

^d Excludes migration year 2003.

Table 13. Preliminary McNary Dam smolt to Bonneville Dam adult SAR-indices for hatchery-origin PIT-tagged summer and fall-run chinook released in the Yakima subbasin by brood year and life stage at release, 2006-2015 (PTAGIS query run May 6, 2019).

Brood Year	Subyearlings		Yearlings	
	Summer	Fall	Summer	Fall
2006		0.0%		8.5%
2007		2.3%		1.2%
2008	2.1%	0.5%		3.0%
2009	2.0%	1.1%		0.7%
2010	3.8%	0.0%	1.9%	1.6%
2011	1.7%	1.2%		1.6%
2012	1.3%	0.9%		
2013	1.1%	0.4%		
2014	0.0%	0.0%		
2015	0.2%	0.4%		
Pooled Mean	1.8%	1.1%	1.9%	1.7%

Table 14. Preliminary McNary Dam smolt to Bonneville Dam age-3 adult return (SAR) indices for hatchery-origin PIT-tagged coho released as smolt (sm) or parr^a in Lower Yakima (LY), Naches (Na), and Upper Yakima (UY) mainstem or tributary areas, brood years 2003-2014 (PTAGIS queries run April 16, 2019).

	LY_sm	Na_sm	UY_sm	Na_parr	UY_parr
2003	3.78%	6.14%	2.92%		
2004	2.28%	3.16%	3.67%	1.09%	
2005	3.11%	3.31%	2.36%	1.41%	1.96%
2006	9.76%	6.81%	4.17%	5.52%	7.84%
2007	8.16%	2.84%	4.35%	0.52%	3.16%
2008	4.10%	7.59%	8.80%	5.84%	8.30%
2009	0.20%	1.89%	3.37%	1.99%	3.20%
2010	1.67%	1.80%	1.76%	0.98%	3.23%
2011	6.57%	7.15%	11.64%	6.11%	10.49%
2012	1.15%	1.48%	2.58%	1.01%	2.59%
2013	3.35%	2.33%	4.91%		3.03%
2014	0.66%	3.01%	3.05%	3.73%	6.74%
Average	3.73%	3.96%	4.46%	2.82%	5.05%
Geomean	2.46%	3.40%	3.85%	2.03%	4.33%

^a PIT-tagged fish released as parr in brood year 2003, 2004 (Upp. Yak.), and 2013 (Naches) experienced very poor (<1%) survival to McNary Dam as juvenile smolts and were omitted from this analysis.

Discussion:

Calculation of smolt-to-adult survival rate indices for Yakima Basin anadromous salmonids are complicated by the following factors:

- 1) Smolt accounting at Prosser is based on statistical expansion of Chandler smolt trap sampling data using available PIT-detection and flow data and estimated Chandler entrainment rates. Chandler smolt passage estimates are prepared primarily for the purpose of comparing relative marked versus unmarked passage estimates and not for making survival comparisons. While these Chandler smolt passage estimates represent the best available data, there may be a high degree of error associated with

these estimates due to inherent complexities, assumptions, and uncertainties in the statistical expansion process. Therefore, these estimates are subject to revision.

2) Large numbers of Yakima Basin salmonid releases (all CESRF spring Chinook) are adipose-fin clipped and subjected to higher harvest rates than unmarked wild/natural fish in marine and Columbia River mark-selective fisheries. No adjustments have yet been made in the above SAR estimates to account for differential harvest rates in these mark-selective fisheries.

3) Due to issues such as water diversion permitting, size required for tagging, and allowing sufficient time for acclimation, release time for many hatchery-origin juveniles (including all CESRF spring Chinook) may be delayed relative to their wild counterparts. For example, spring Chinook from the CESRF are not allowed to voluntarily migrate until at least March 15 of their smolt outmigration year; however, juvenile sampling observations at Roza Dam indicate that a substantial number of wild/natural juveniles migrate downstream during the summer, fall, and winter months prior to their smolt outmigration year. Analysis of juvenile migrant PIT detections at Roza Dam (PTAGIS queries run 7/12/2013) indicated that approximately 81% of natural-origin spring Chinook migrated downstream of Roza in the fall or winter as juveniles (before CESRF fish would have the opportunity). Comparison of SAR data for non-contemporaneously migrating juveniles may be invalid.

Given these complicating factors, Tables 10-14 present available smolt-to-adult survival indices for Yakima River spring and summer/fall Chinook and coho. Because of the complexities noted above, these data are useful for analysis of trends but should not be used as direct citations of, or for comparisons of marked and unmarked, smolt-to-adult survival rates. The reader is encouraged to contact Yakama Nation technical staff to discuss these and other issues prior to any use of these data or any other estimation of Yakima Basin SARs that may be available through data obtained from public web sites such as RMPC, PTAGIS, DART, FPC or others.

Substantial juvenile mortality of subyearling releases of summer- and fall-run Chinook occurs in the Yakima River between their release sites and McNary Dam (Neeley 2012b). Strategies have been proposed to address limiting factors (YSFWPB 2004) and improve survival of these releases (Yakama Nation 2019). As these strategies are implemented, we expect SARs for summer- and fall-run Chinook to improve substantially from the estimates provided in Table 11 (Yakama Nation 2019). Additional discussion and results for Yakima Basin spring Chinook SARs are presented in Appendix B.

Status and Trend of Spatial Distribution (Redd Counts)

Methods: Regular foot and/or boat surveys (monitoringresources.org methods 30, 131, 285, 1508) were conducted within the established geographic range for each species (this is increasing for coho as acclimation sites are located upriver and as the run increases in size). Redds were individually marked during each survey and carcasses were sampled to collect egg retention, scale sample, sex, and body length information and to check for possible experimental marks. River conditions vary from year to year and preclude complete accounting, especially for fall Chinook and Coho. Other agencies (WDFW, Pacific Northwest National Laboratory, and private contractors) have also conducted foot, boat, or aerial surveys for fall Chinook redds in the Yakima River Basin and we have attempted to incorporate available information from those surveys here.

Results:




Figure 13. Redd Counts upstream of Prosser Dam in the Yakima River Basin by species, 1981-present.

Table 15. Yakima Basin spring Chinook redd counts and distribution, 1981 – present.

Year	Upper Yakima River System				Naches River System				
	Mainstem ¹	Elum	Teaway	Total	American	Naches ¹	Bumping	Little Naches	Total
1981	237	57	0	294	72	64	20	16	172
1982	610	30	0	640	11	25	6	12	54
1983	387	15	0	402	36	27	11	9	83
1984	677	31	0	708	72	81	26	41	220
1985	795	153	3	951	141	168	74	44	427
1986	1,716	77	0	1,793	464	543	196	110	1,313
1987	968	75	0	1,043	222	281	133	41	677
1988	369	74	0	443	187	145	111	47	490
1989	770	192	6	968	187	200	101	53	541
1990	727	46	0	773	143	159	111	51	464
1991	568	62	0	630	170	161	84	45	460
1992	1,082	164	0	1,246	120	155	99	51	425
1993	550	105	1	656	214	189	88	63	554
1994	226	64	0	290	89	93	70	20	272
1995	105	12	0	117	46	25	27	6	104
1996	711	100	3	814	28	102	29	25	184
1997	364	56	0	420	111	108	72	48	339
1998	123	24	1	148	149	104	54	23	330
1999	199	24	1	224	27	95	39	25	186
2000	3,349	466	21	3,836	54	483	278	73	888
2001	2,910	374	21	3,305	392	436	257	107	1,192
2002	2,441	275	110	2,826	366	226	262	89	943
2003	772	87	31	890	430	228	216	61	935
2004	2,985	330	129	3,444	91	348	205	75	719
2005	1,717	287	15	2,019	140	203	163	68	574
2006	1,092	100	58	1,250	136	163	115	33	447
2007	665	51	10	726	166	60	60	27	313
2008	1,191	137	47	1,375	158	165	102	70	495
2009	1,349	197	33	1,579	92	159	163	68	482
2010	2,199	219	253	2,671	173	171	168	40	552
2011	1,663	171	64	1,898	212	145	175	48	580
2012	1,276	125	69	1,470	337	196	189	89	811
2013	552	85	34	671	170	66	85	55	376
2014	962	138	53	1,153	129	65	158	27	379
2015	1,258	39	24	1,321	239	177	152	46	614
2016	512	83	22	617	149	106	74	37	366
2017	402	118	23	543	123	84	56	30	293
2018	339	13	0	352	27	56	44	1	128
2019	184	44	9	237	21	1	2	7	31
2020	189	44	8	241	44	25	71	6	146
Mean	980	119	26	1,125	153	157	109	45	464

¹ Including minor tributaries.




Figure 14. Teanaway River Spring Chinook redd counts, 1981-2020 (vertical lines denote pre- and post-supplementation periods) and the proportion of natural-origin (NO) carcasses observed in intensive spawning ground surveys, 2002-2010.




Figure 15. Distribution of summer and fall run Chinook redds in the Yakima River Basin (above Prosser Dam) based on redd observations from 2014 to 2018.




Figure 16. Fall Chinook redd counts above and below Prosser Dam, 1961-present, for years in which surveys were conducted and data are available. Data from YN, WDFW, and Pacific Northwest National Laboratory files. Note that survey completeness is highly variable due to annual flow and turbidity conditions; survey data are partial or incomplete for most years prior to 2000.




Figure 17. Distribution of coho redds in the Yakima River Basin.

Table 16. Yakima Basin coho redd counts and distribution, 1998 – present.

	Yakima River	Naches River	Tributaries	Total
1998	53	6	193	252
1999	104		62	166
2000	142	137	67	346
2001	27	95	25	147
2002	4	23	16	43
2003	32	56	55	143
2004	33	87	150	270
2005	57	72	153	282
2006	44	76	187	307
2007	63	87	195	345
2008	49	60	242	351
2009	229	281	485	995
2010	75	276	327	678
2011	82	243	196	521
2012	148	228	172	548
2013	45	69	67	181
2014	320	86	751	1157
2015	16	0	47	63
2016	27	37	54	118
2017	92	36	177	305
2018	46	103	100	249
2019	62	80	112	254
2020	71	50	95	216

Discussion:

Spatial distribution of spring Chinook spawners has increased as a result of acclimation site location, salmon homing fidelity and more fully seeding preferred spawning habitats (Dittman et al. 2010). Redd surveys in the Teanaway River conducted annually by Yakama Nation staff since 1981 demonstrate the benefits of reintroducing salmonids into underutilized habitat (Figure 14). The Jack Creek acclimation site began releasing CESRF spring chinook in 2000, with the first age-4 females returning from these releases in 2002. Redd counts in this tributary have increased from a pre-supplementation average of 3 redds per year to a post supplementation average of 55 redds per year. The proportion of natural-origin carcasses increased from less than one percent in 2002 (when CESRF fish first returned to the natural spawning grounds) to 42% in 2006 when the progeny of the 110 redds produced in 2002 (virtually 100% of which were produced by CESRF-origin fish) returned. These data clearly indicate that naturally-spawning CESRF spring Chinook were successful in returning natural-origin adults back to the Teanaway River. However, redd counts in the Teanaway River remain at or below pre-supplementation levels in some years, including 2018, indicating that habitat factors (primarily low late-summer and fall season flows) continue to deter returning fish and these fish are likely spawning in nearby mainstem and tributary reaches more conducive to survival of progeny (Fast et al. 2015).

Fall Chinook redd distribution in the Yakima River Basin appears to be experiencing a major transition in recent years. Historical redd survey data indicates that a substantial number of fall Chinook spawned below Prosser Dam in the lower Yakima River. However, from 2003-present, an average of approximately 80 percent (range 62 to 90 percent) of surveyed fall Chinook redds have been located above Prosser Dam (Figure 16). Biologists and habitat experts in the subbasin at least partially attribute this change in spawning distribution to the invasion of water stargrass (see Wise et al. 2009) in the lower 43 miles of the Yakima River. With the reintroduction of summer run Chinook, the Yakama Nation is expanding the distribution of summer/fall run Chinook spawners and redds into the middle reaches of the Yakima Basin between the town of Wapato upstream to the confluence with the Tieton River in the Naches subbasin and to Roza Dam in the Upper Yakima subbasin (Figures 1 and 15; Yakama Nation 2012). Summer-run Chinook have now spawned naturally in these habitats since 2013 after an absence of over 40 years.

Coho redd counts and spawner distribution have increased substantially since reintroduction efforts began (Table 16 and Figure 17). Many redds in the mainstem were located intermixed with fall chinook redds, tucked under cut banks or were found in side channels. Tributary redd enumeration and identification continues to be accurate due to the fall low water levels, improving interagency cooperation, and relatively good weather. One of the overall goals during the present implementation phase (Phase II) of the coho program is to evaluate the transition of redds from the mainstem river into historic tributaries. With the beginning of Phase II of the Coho Program we observed large increases in tributary spawning, with an annual average exceeding 200 redds counted in tributaries since 2004 (Table 16). Although, there were large numbers of potential spawners in 2014 (~9,000 females), river conditions were very unfavorable for finding redds. Winter anchor ice in early December kept surveys to a minimum. This was followed by winter freshets that reduced visibility in the Naches River to the point where visibility was near zero. However, the stability of low water conditions in 2015 might have contributed to good survival of coho eggs from the 2014-2015 spawning season. The 2020 redd count was difficult to perform because of the ongoing pandemic, but we did observe an increase in the overall redd counts for coho (Table 16). Coho continue to volunteer into many tributaries, and the fidelity of adults from summer parr plants has shown good results.

Adult Coho plants have also been used to evaluate the feasibility of increasing fish abundance in several tributaries. To determine the spawning success and effects on resident trout of these adult outplants, an intensive monitoring program was conducted in Taneum Creek for brood/spawn years 2007-2014. The results of this evaluation indicate that Coho spawned successfully and have the potential to produce large numbers of returning adult offspring per smolt that survive to McNary Dam as

juveniles (Table 17). The total biomass of all salmonids in the stream increased and there were no discernable impacts to resident trout (Temple et al. 2012, 2017). Adult coho were not planted in 2019 and 2020. However, there are plans for adult out plants in 2021.

Table 17. Results from Taneum Creek adult out-plant study.

Year	Number of Adult Females Outplanted	Redds	Number of Juvenile coho PIT Tagged	McNary Juvenile PIT Detections	McNary Juvenile & Adult PIT Detections	McNary Juvenile-Adult SAR
2007	150	75	1,299	94		
2008	150	50	1,868	82	7	8.5%
2009	150	130	4,515	177	4	2.3%
2010	150	134	1,054	73	3	4.1%
2011	150	100	743	30	4	13.3%
2012	60	54	1,941	70		
2013	9	5	231	0		
2014	360	200	752	12		
Pooled			12,403	538	18	3.3%

Status and Trend of Diversity Metrics

Methods:

Diversity metrics collected for the Cle Elum Supplementation and Research Facility spring Chinook program in the Upper Yakima River include parameters relating to: eggs (e.g., egg size, KD at emergence, emergence timing, etc.), juveniles (growth and survival, migration timing, fish health, etc.), and adults (size at age, sex composition, migration timing, etc.). Methods for monitoring the spring Chinook program were documented in: the YKFP Monitoring Plan ([Busack et al. 1997](#)), the project's "[Supplementation Monitoring Plan](#)" (Chapter 7 in 2005 annual report on project genetic studies), and numerous manuscripts in the published literature (see Results and References).

Diversity metrics for returning adult summer/fall Chinook and coho collected at the Prosser Dam denil fish trap include sex ratios, lengths, and weights ([monitoringresources.org](#) methods 454, 1454, 1548, 1549, 1551, 4008, 4041). We also queried the PTAGIS database for PIT-tagged summer- and fall-run Chinook that were released in the Yakima Subbasin in recent years and used PIT-detection data at Bonneville Dam for upstream migrants to estimate age composition and run timing of returning fish.

Results and Discussion:

A detailed presentation of current results for the spring Chinook monitoring program (YN-collected data) are included in Appendix B of this report and are discussed in greater detail in the annual report(s) for WDFW-companion project [1995-064-25](#). Generally, we have detected small, but significant differences between hatchery- and natural-origin fish in some juvenile and adult traits. Results in the published literature include: Busack et al. (2007), Knudsen et al. (2006, 2008), Larsen et al. (2004, 2006, 2010, 2013), and Pearsons et al. (2009).

Sex ratios, lengths, and weight data for fall Chinook and coho salmon sampled at the Prosser denil adult sampling facility from 2001-present are presented in Tables 18-21. Age composition of summer- and fall-run Chinook are presented in Table 22 and run timing in Figure 18. In addition, preliminary results of some diversity metrics relating to the effort to reestablish a natural spawning coho population in the Yakima Basin were published in Bosch et al. (2007). That study observed divergence in some diversity traits between hatchery- and natural-origin fish suggesting that some re-naturalization can be detected in just a few generations after outplanting of hatchery-origin fish in the wild.

Table 18. Sex ratio of upstream migrating fall Chinook sampled at the Prosser Dam right bank denil ladder and fish trap, 2001-present.

Return Year	Sample Size			Female Adult %	Female Total %	Sample Date Range	
F	J	M			First	Last	
2001	186	80	213	46.6%	38.8%	09/10/01	11/19/01
2002	389	61	512	43.2%	40.4%	09/09/02	11/25/02
2003	396	24	224	63.9%	61.5%	09/07/03	11/17/03
2004	185	40	201	47.9%	43.4%	09/06/04	11/23/04
2005	201	8	233	46.3%	45.5%	09/06/05	11/14/05
2006	107	11	84	56.0%	53.0%	09/13/06	11/06/06
2007	42	44	39	51.9%	33.6%	09/10/07	11/06/07
2008	81	23	101	44.5%	39.5%	09/08/08	11/13/08
2009	110	132	95	53.7%	32.6%	09/08/09	11/07/09
2010	239	4	162	59.6%	59.0%	09/08/10	11/03/10
2011	67	10	34	66.3%	60.4%	09/07/11	11/09/11
2012	249	109	264	48.5%	40.0%	09/04/12	11/06/12
2013	272	86	460	37.2%	33.3%	09/16/13	11/22/13
2014	681	78	725	48.4%	45.9%	09/04/14	12/10/14
2015	1047	69	1374	43.2%	42.0%	09/09/15	11/16/15
2016	158	22	128	55.2%	51.3%	09/09/16	11/12/16
2017	122	67	66	64.9%	47.8%	09/13/17	12/05/17
2018	78	23	114	40.6%	36.3%	09/12/18	11/05/18
2019	36	7	22	62.1%	55.4%	09/22/19	11/15/19
2020	20		25	44.4%	44.4%	09/23/20	11/20/20
			Mean	51.2%	45.2%		

Table 19. Sample size (N), mean fork and mid-eye to hypural plate (MEH) lengths (cm), and weights (pounds) of upstream migrating fall Chinook sampled at the Prosser Dam right bank denil ladder and fish trap, 2001-present.

Run Year	Females				Males (excluding Jacks)			
	N	Fork	POH	Weight	N	Fork	POH	Weight
2001	186	72.7	60.1	11.0	213	71.5	57.8	9.3
2002	389	78.4	63.9	13.5	512	76.1	60.2	12.1
2003	396	83.4	68.5	15.6	224	83.7	67.0	16.3
2004	185	82.3	67.8	15.1	201	73.9	60.0	11.2
2005	201	80.5	66.3	14.2	233	75.1	60.6	11.5
2006	107	81.5	66.3	15.6	84	81.3	64.6	15.3
2007	42	79.9	64.4	14.8	39	72.8	56.8	11.7
2008	81	70.1	56.5	9.8	101	67.8	54.0	8.9
2009	110	74.1	57.8	11.2	95	69.4	52.5	9.6
2010	239	73.3	57.8	11.3	162	70.9	54.7	9.7
2011	67	76.5	60.4	12.4	34	74.2	57.7	11.3
2012	249	70.1	53.3	9.5	264	66.4	49.6	7.9
2013	272	72.5	56.1	10.1	460	69.8	52.9	8.7
2014	681	76.1	60.8	11.9	725	69.0	53.2	8.6
2015	1047	76.2	59.5	11.4	1374	71.4	54.8	9.2
2016	158	75.3	59.5	9.7	128	71.6	55.3	8.1
2017	122	74.6	58.8	10.8	66	73.9	57.1	10.4
2018	78	72.3	54.4	9.6	114	67.2	48.9	7.5
2019	36	70.2	55.3	8.7	22	68.4	54.2	7.9
2020	20	71.9	51.7	9.1	25	71.4	51.9	8.5
Mean		75.6	60.0	11.8		72.3	56.2	10.2

Table 20. Sex ratio of upstream migrating coho sampled at the Prosser Dam right bank denil ladder and fish trap, 2001-present.

Return Year	Sample Size			Female	Female	Sample Date Range	
	F	J	M	Adult %	Total %	First	Last
2001	1147	44	1024	52.8%	51.8%	09/11/01	11/22/01
2002	72	201	71	50.3%	20.9%	09/11/02	11/25/02
2003	473	89	452	51.1%	46.6%	09/11/03	11/21/03
2004	586	49	509	53.5%	51.2%	09/07/04	11/16/04
2005	531	146	405	56.7%	49.1%	09/13/05	11/15/05
2006	826	97	586	58.5%	54.7%	09/17/06	11/19/06
2007	676	34	538	55.7%	54.2%	09/11/07	11/20/07
2008	666	930	514	56.4%	31.6%	09/08/08	12/04/08
2009	1644	76	1576	51.1%	49.9%	09/09/09	11/20/09
2010	999	35	673	59.7%	58.5%	09/08/10	11/19/10
2011	907	12	776	53.9%	53.5%	09/16/11	11/17/11
2012	1156	108	961	54.6%	52.0%	09/08/12	11/17/12
2013	523	146	528	49.8%	43.7%	09/20/13	11/22/13
2014	4302	135	3668	54.0%	53.1%	09/03/14	12/23/14
2015	656	67	683	49.0%	46.7%	09/13/15	12/09/15
2016	310	101	249	55.5%	47.0%	09/13/16	11/16/16
2017	694	132	752	48.0%	44.0%	09/13/17	12/19/17
2018	343	318	308	52.7%	35.4%	09/06/18	11/05/18
2019	758	28	692	52.3%	51.3%	09/04/19	12/31/19
2020	357	115	180	66.5%	54.8%	09/22/20	11/25/20
			Mean	54.1%	47.5%		

Table 21. Sample size (N), mean fork and mid-eye to hypural plate (MEH) lengths (cm), and weights (pounds) of upstream migrating coho sampled at the Prosser Dam right bank denil ladder and fish trap, 2001-present.

Run Year	N	Females			Males (excluding Jacks)			
		Fork	POH	Weight	N	Fork	POH	Weight
2001	1147	65.4	53.7	6.7	1024	65.6	52.4	6.5
2002	72	68.1	54.9	8.5	71	69.4	54.0	8.1
2003	473	65.3	52.9	7.0	452	65.7	51.4	6.8
2004	586	68.8	56.4	8.0	509	67.8	53.9	7.4
2005	531	67.5	54.9	8.0	405	67.6	53.5	7.8
2006	826	71.6	58.2	10.0	586	71.3	55.8	9.4
2007	676	66.3	52.1	7.0	538	65.5	49.9	6.6
2008	666	69.9	56.7	9.6	516	69.8	54.6	9.0
2009	1644	68.1	52.4	7.9	1576	67.2	49.7	7.2
2010	999	69.7	54.2	8.7	673	68.5	51.5	7.8
2011	907	68.6	53.7	8.2	776	68.5	51.7	7.7
2012	1156	64.3	49.5	6.8	961	62.6	46.4	6.0
2013	523	66.2	51.9	6.9	528	64.0	48.4	5.9
2014	4302	65.6	52.6	7.0	3668	63.5	49.8	6.1
2015	656	63.5	50.1	6.0	683	61.9	47.5	5.2
2016	310	66.9	52.7	6.9	249	67.4	51.6	6.4
2017	694	64.5	49.6	6.4	752	63.6	47.8	5.9
2018	343	66.6	51.0	6.8	308	66.0	49.2	6.4
2019	758	64.8	49.7	5.7	692	63.7	47.7	5.2
2020	357	67.4	49.8	7.9	180	66.4	47.9	7.0
Mean		67.0	52.9	7.5		66.3	50.7	6.9

Table 22. Age composition of returning hatchery-origin PIT-tagged summer and fall-run chinook released in the Yakima subbasin as subyearling or yearling fish (data from PTAGIS query run May 1, 2019).

Brood Year	Age at Return				
	2	3	4	5	6
Summer Chinook Subyearlings					
2008	12.5%	12.5%	50.0%	25.0%	0.0%
2009	5.4%	16.3%	63.6%	14.7%	0.0%
2010	0.2%	27.5%	61.4%	10.6%	0.2%
2011	0.0%	12.1%	67.5%	20.4%	0.0%
2012	1.0%	50.0%	40.8%	8.2%	0.0%
2013	5.6%	11.1%	77.8%	5.6%	0.0%
Mean	4.1%	21.6%	60.2%	14.1%	0.0%
Fall Chinook Subyearlings					
2007	9.7%	47.9%	35.8%	6.6%	
2008	13.3%	53.3%	33.3%	0.0%	
2009	18.9%	40.5%	32.4%	8.1%	
2010	0.0%	66.7%	16.7%	16.7%	
2011	11.6%	34.9%	50.0%	3.5%	
2012	9.7%	61.1%	26.4%	2.8%	
Mean	10.6%	50.7%	32.4%	6.3%	
Summer Chinook Yearlings					
2010 ¹	13.6%	31.2%	44.2%	3.9%	0.6%
Fall Chinook Yearlings					
2006	96.4%	0.0%	3.6%	0.0%	0.0%
2007	63.2%	16.2%	8.8%	11.8%	0.0%
2008	30.9%	36.2%	27.1%	5.8%	0.0%
2009	20.4%	19.4%	40.8%	19.4%	0.0%
2010	39.4%	26.8%	27.8%	6.1%	0.0%
2011	6.4%	16.7%	57.1%	14.7%	5.1%
Mean	42.8%	19.2%	27.5%	9.6%	0.9%

¹ 10 of 154 (6.5%) of detections occurred about 90 days post-release in adult ladders at Bonneville Dam and were assumed to be age-1 returns. However, only 2 of these 10 were confirmed as upstream detections based on later detections at dams upstream of Bonneville. The other 8 detections at Bonneville could have been late-migrating juveniles.




Figure 18. Adult return timing at Prosser Dam of PIT-tagged summer- and fall-run Chinook reared at the Marion Drain and Prosser Hatcheries and released as subyearlings, pooled for return years 2009-2018.

Habitat Monitoring

While the majority of YKFP habitat activities in the Yakima Basin are addressed in a separate project ([1997-051-00](#)), we are monitoring stream sediment loads associated with the operation of dams and other anthropogenic factors (e.g. logging, agriculture and road building) under this contract as sediment loads can affect survival of salmonids (see description and references [here](#) and [here](#)).

Status and Trend of Fine Sediment

Methods: Representative gravel samples (McNiel core samples, monitoring resources [1504](#)) were collected from various reaches in the Little Naches and Upper Yakima Rivers in the fall of 2020. Each sample was analyzed to estimate the percentage of fine or small particles present (<0.85 mm). The Washington State Timber, Fish, and Wildlife program established guidelines that specify the impacts that estimated sedimentation levels can have on salmonid egg-to-smolt survival. These impact guidelines will inform future analyses of “extrinsic” factors on natural production in the Yakima Basin.

Results and Discussion:

Little Naches

A total of 106 McNiel core samples were collected and processed from 9 spawning reaches in the Little Naches drainage this past year. Pyramid Creek has not been sampled since 2009 when the main road going into this reach was decommissioned. Other means to access this sampling site is needed. With this year’s monitoring work, the data set for the Little Naches drainage now covers a time period of 36 years for the two historical reaches, and 29 years for the expanded sampling area that includes several tributary streams.

The average percent fine sediment less than 0.85mm for the entire Little Naches drainage in 2020 was 7.3% which is the lowest watershed average observed on record (Figure 19). The overall trend remains downward and similar trends can be seen when looking at individual reach conditions over the longer term monitoring period since 1992.

The overall average fine sediment found in spawning substrate remains relatively low and should lessen mortality on incubating eggs and alevins. The reduced rate of fine sediment found can be partially attributed to less anthropogenic disturbance occurring in the watershed in recent years, other than recreational activity. Timber harvest activity and road building has been minimal for several years. Landowners have also improved roads and trails to reduce sediment delivery.

Further, enhanced stream protection measures have been instituted through the Northwest Forest Plan and the Central Cascades Habitat Conservation Plan for over 20 years. These factors have likely helped reduce fine sediment inputs to the stream system. However recreational activity, such as dispersed camping sites and off-road vehicle use near streams, continues to be a concern. Sediment delivery, bank erosion, and loss of riparian vegetation from recreational use have been observed in some localized areas.




Figure 19. Overall Fine Sediment (<0.85mm) Trends with 95% confidence bounds in the Little Naches River Drainage, 1992-2020.

South Fork Tieton

One reach on the South Fork Tieton River (in the vicinity of Minnie Meadows) has been sampled in the past by the U.S. Forest Service. To the best of our knowledge this reach has not been sampled since 2015. This stream reach typically receives significant bull trout spawning activity and the monitoring efforts provide valuable information on their spawning conditions. Average fine sediment in this reach was 8.9% in 2015, matching the previous low observed in 1999, and is well below the mean for sediment levels for the 17 years that were sampled (Figure 20).

Upper Yakima

A total of 60 samples were collected and processed from the Upper Yakima River drainage this past year (5 reaches, 12 samples from each reach). The same reaches (Stampede Pass, Easton, Camelot to Ensign Ranch, Elk Meadows, and Cle Elum) have been sampled annually for the past 24 years. The 24-year trend in average percent fine sediment less than 0.85mm for the combined Upper Yakima drainage remains downward, with 2020 being among the four lowest years on record (Figure 21).




Figure 20. Fine Sediment Trends in the South Fork Tieton River, 1999-2015. Note: Data for 2007 were collected from only 1 Riffle. Data courtesy of U.S. Forest Service.




Figure 21. Overall average percent fine sediment (< 0.85 mm) in spawning gravels of the Upper Yakima River, 1997-2020.

Summary

We continue to observe a general decreasing trend in average fine sediment levels in the Little Naches and Upper Yakima drainages. Increases observed in 2016-2019 in both drainages may have been due to effects from the large fires the region has experienced in recent years. Overall, the generally low rates of fine sediment should be conducive for egg and alevin survival and should favor salmonid spawning success.

The results of the USFS sampling in the South Fork Tieton River were also low over a 17-year sampling period. These conditions should be favorable for early life history survival of bull trout.

Detailed field data including additional tables and graphs for samples collected in the upper Yakima and Naches basins can be obtained from Jim Matthews, fisheries biologist for the Yakama Nation (matj@yakamafish-nsn.gov).

Harvest Monitoring

Marine and Mainstem Columbia Fisheries

Methods: We evaluated recoveries of coded-wire tags (CWTs) and PIT tags in out-of-basin fisheries using queries of regional mark information system ([RMIS](#)) and PIT Tag Information System ([PTAGIS](#)) databases. We coordinated with agencies responsible for harvest management (WDFW, ODFW, USFWS, CRITFC, etc.) to estimate the harvest of target stocks. We reviewed reports produced annually by the [Pacific Fisheries Management Council](#) (marine) and the *U.S. v Oregon* [Technical](#)

[Advisory Committee](#) (mainstem Columbia) to evaluate estimated harvest or exploitation rates on comparable stocks in these fisheries.

For spring Chinook, additional information was employed that is not readily available for fall Chinook and coho. Standard run reconstruction techniques (Appendix B) were employed to derive estimates of harvest from the Columbia River mouth to the Yakima River mouth for spring Chinook. Data from databases maintained by the *U.S. v Oregon* [Technical Advisory Committee](#) were used to obtain harvest rate estimates downstream of the Yakima River for the aggregate Yakima River spring Chinook population and to estimate passage losses from Bonneville through McNary reservoirs. These data, combined with the Prosser Dam counts and estimated harvest below Prosser, were used to derive a Columbia River mouth run size estimate and Columbia River mainstem harvest estimate for Yakima spring Chinook.

Results:

Table 23. Marine and freshwater recoveries of CWTs from brood year 1997-2015 releases of spring Chinook from the CESRF as reported to the Regional Mark Information System (RMIS) Jan. 5, 2021.

Brood Year	Observed CWT Recoveries			Expanded CWT Recoveries		
	Marine	Fresh	Marine %	Marine	Fresh	Marine %
1997	5	56	8.2%	8	321	2.4%
1998	2	53	3.6%	2	228	0.9%
1999		2	0.0%		9	0.0%
2000		14	0.0%		34	0.0%
2001		1	0.0%		1	0.0%
2002		7	0.0%		36	0.0%
2003		4	0.0%		10	0.0%
2004	2	154	1.3%	15	526	2.8%
2005	2	96	2.0%	2	304	0.7%
2006	14	328	4.1%	16	1160	1.4%
2007	8	145	5.2%	13	1139	1.1%
2008	5	245	2.0%	7	1634	0.4%
2009	4	91	4.2%	7	588	1.2%
2010	4	164	2.4%	9	948	0.9%
2011	5	186	2.6%	5	1030	0.5%
2012	4	73	5.2%	2	273	0.7%
2013	9	65	12.2%	20	534	3.6%
2014	4	71	5.3%	8	533	1.5%
2015 ¹	2	23	8.0%	2	49	3.9%

1. Reporting of CWT recoveries to the RMIS database typically lags actual fisheries by one to two years. Therefore, CWT recovery data for brood year 2015 are considered preliminary or incomplete.

Table 24. Estimated run size, harvest, and harvest rates of Yakima Basin spring Chinook in Columbia River mainstem and terminal area fisheries, 1983-present.

Year	Columbia R. Mouth Run Size	Col. R.	BON to McNary Harvest	Yakima R. Mouth Run Size	Yakima River Harvest	Columbia Basin Harvest Summary			Col. Basin Harvest Rate	
		Mouth to BON Harvest				Total	Wild	CESRF	Total	Wild
1983	2,460	118	113	1,441	84	316	316	0	12.8%	12.8%
1984	3,911	135	290	2,658	289	714	714	0	18.3%	18.3%
1985	5,276	192	197	4,560	865	1,254	1,254	0	23.8%	23.8%
1986	13,567	280	802	9,439	1,340	2,423	2,423	0	17.9%	17.9%
1987	6,160	96	378	4,443	517	991	991	0	16.1%	16.1%
1988	5,674	363	401	4,246	444	1,208	1,208	0	21.3%	21.3%
1989	8,919	213	683	4,914	747	1,642	1,642	0	18.4%	18.4%
1990	6,954	352	480	4,372	663	1,495	1,495	0	21.5%	21.5%
1991	4,650	184	291	2,906	32	507	507	0	10.9%	10.9%
1992	6,207	103	380	4,599	345	827	827	0	13.3%	13.3%
1993	5,132	44	315	3,919	129	488	488	0	9.5%	9.5%
1994	2,251	87	113	1,302	25	225	225	0	10.0%	10.0%
1995	1,394	1	69	666	79	149	149	0	10.7%	10.7%
1996	5,898	6	309	3,179	475	790	790	0	13.4%	13.4%
1997	5,192	3	348	3,173	575	926	926	0	17.8%	17.8%
1998	2,868	3	144	1,903	188	335	335	0	11.7%	11.7%
1999	4,154	4	192	2,781	604	800	800	0	19.3%	19.3%
2000	28,753	58	1,752	19,101	2,458	4,267	4,144	123	14.8%	14.8%
2001	32,307	971	4,281	24,149	4,630	9,882	5,685	4,197	30.6%	28.7%
2002	25,256	1,275	2,877	15,814	3,108	7,259	2,736	4,524	28.7%	23.9%
2003	10,278	286	903	7,227	440	1,628	987	641	15.8%	14.7%
2004	24,212	1,023	2,330	16,820	1,679	5,031	2,877	2,154	20.8%	16.2%
2005	13,317	354	906	9,589	474	1,735	1,375	360	13.0%	12.1%
2006	12,197	311	944	6,594	600	1,855	1,068	787	15.2%	13.5%
2007	5,223	174	457	4,457	279	910	449	461	17.4%	15.1%
2008	12,554	1,204	1,870	9,273	1,532	4,607	1,360	3,247	36.7%	25.2%
2009	13,693	1,210	1,089	11,395	2,353	4,651	1,318	3,333	34.0%	23.9%
2010	18,565	1,631	2,778	13,745	1,741	6,149	1,516	4,633	33.1%	21.8%
2011	23,316	1,098	1,794	18,520	4,380	7,272	2,590	4,682	31.2%	22.4%
2012	17,315	856	1,633	12,616	3,320	5,809	2,370	3,438	33.5%	26.6%
2013	14,933	880	974	10,602	2,653	4,507	1,817	2,690	30.2%	23.3%
2014	17,303	716	2,222	11,868	2,171	5,110	2,097	3,012	29.5%	22.5%
2015	11,991	476	1,440	9,848	815	2,731	1,457	1,274	22.8%	17.8%
2016	10,107	454	996	7,281	444	1,894	971	923	18.7%	15.5%
2017	12,196	493	920	7,544	1,272	2,685	853	1,831	22.0%	13.5%
2018	6,237	248	636	3,737	548	1,433	459	975	23.0%	16.4%
2019	3,784	68	260	2,251	40	368	131	238	9.7%	8.6%
2020 ¹	5,764	62	347	3,413	68	476	276	200	8.3%	7.7%
Mean	11,381	445	1,038	7,934	1,176	2,659	1,410	1,249	20.0%	17.0%

1. Preliminary.




Figure 22. Distribution of coded-wire tag recoveries of Yakima Basin summer/fall run Chinook releases in marine, mainstem Columbia River, and Yakima Basin fisheries. Data retrieved from the regional mark information system (RMIS) for brood year 1997-2007 recoveries.

Recovery data for Yakima River-origin coho are presently limited because few fish have been coded wire-tagged until recent years. We will continue to collect and analyze CWT-recovery data from regional databases and will report this information in the future. ‘All H Analyzer’ (AHA) modeling for Master Planning purposes assumed that natural- and hatchery-origin Yakima River coho have an exploitation rate of approximately 40 and 60 percent, respectively (Yakama Nation 2019). These estimates include coho caught in marine, Columbia River and Yakima River fisheries.

Discussion:

Based on available CWT information, harvest managers have long assumed that Columbia River spring Chinook are not harvested in any abundance in marine fisheries as their ocean migration does not generally overlap either spatially or temporally with the occurrence of marine fisheries (TAC 1997). Harvest recoveries of CESRF spring Chinook as reported to RMIS to date appear to confirm this, as marine harvest apparently accounts for only about 0-3% of the total harvest of Yakima Basin spring Chinook (Table 23). Adult returns of spring Chinook from the CESRF appear to be making substantial contributions to Columbia Basin fisheries (Table 24).

Yakima Basin summer/fall Chinook are harvested in marine fisheries from Alaska to southern Oregon, and in Columbia River fisheries from the mouth to the Hanford Reach (Figure 22). Approximately 71% of harvest recoveries from Yakima Basin fall Chinook releases for brood years 1997-2007 occurred in marine (44%) and mainstem Columbia (27%) fisheries. Out-of-basin harvest rates have not been estimated

specifically for Yakima Basin summer/fall run Chinook, but the 1982-89 brood year average ocean fisheries exploitation rate for mid-Columbia River summer/fall Chinook was 39%, with a total exploitation rate of 68% estimated for the same years (PSC 1994). Chapman et al. (1994) estimated that the 1975-87 brood year mean exploitation rate for fall Chinook released from Priest Rapids Hatchery was 64%. Harvest rates of these stocks in U.S. fisheries since the mid-1990s have been reduced due to Endangered Species Act (ESA) management concerns as these stocks are intermixed with ESA-listed Snake River fall Chinook populations (NMFS 1999a-d and 2000a-c). It is assumed that Yakima River summer/fall run Chinook are harvested at the same rate in these fisheries as other mid-Columbia River summer/fall Chinook stocks.

Yakima Subbasin Fisheries

Methods: The two co-managers, Yakama Nation and WDFW, are responsible for monitoring their respective fisheries in the Yakima River. Each agency employs fish monitors dedicated to creel surveys and/or fisher interviews at the most utilized fishing locations and/or boat ramps. From these surveys, standard techniques are employed to expand fishery sample data for total effort and open areas and times to derive total harvest estimates. Fish are interrogated for various marks. Methods are consistent with monitoringresources.org methods 404 and 960.

Results:

Table 25. Spring Chinook harvest in the Yakima River Basin, 1983-present.

Year	Tribal		Non-Tribal		River Totals		Total	Harvest Rate ¹
	CESRF	Natural	CESRF	Natural	CESRF	Natural		
1983		84		0		84	84	5.8%
1984		289		0		289	289	10.9%
1985		865		0		865	865	19.0%
1986		1,340		0		1,340	1,340	14.2%
1987		517		0		517	517	11.6%
1988		444		0		444	444	10.5%
1989		747		0		747	747	15.2%
1990		663		0		663	663	15.2%
1991		32		0		32	32	1.1%
1992		345		0		345	345	7.5%
1993		129		0		129	129	3.3%
1994		25		0		25	25	1.9%
1995		79		0		79	79	11.9%
1996		475		0		475	475	14.9%
1997		575		0		575	575	18.1%
1998		188		0		188	188	9.9%
1999		604		0		604	604	21.7%
2000	53	2,305		100	53	2,405	2,458	12.9%
2001	572	2,034	1,252	772	1,825	2,806	4,630	19.9%
2002	1,373	1,207	492	36 ²	1,865	1,243	3,108	20.6%
2003	134	306	0	0	134	306	440	6.3%
2004	289	712	569	109 ²	858	820	1,679	11.0%
2005	46	428	0	0	46	428	474	5.4%
2006	246	354	0	0	246	354	600	9.5%
2007	123	156	0	0	123	156	279	6.5%
2008	521	414	586	11 ²	1,107	426	1,532	17.8%
2009	1,089	715	541	8 ²	1,630	722	2,353	19.4%
2010	345	194	1,154	48 ²	1,499	241	1,741	13.2%
2011	1,361	1,261	1,579	179 ²	2,940	1,440	4,380	24.4%
2012	1,220	1,302	735	63 ²	1,955	1,364	3,320	27.5%
2013	846	975	786	46 ²	1,632	1,021	2,653	25.9%
2014	576	715	826	54 ²	1,402	769	2,171	19.2%
2015	121	271	385	38 ²	506	309	815	8.7%
2016	103	185	132	24 ²	235	209	444	6.4%
2017	217	201	750	104 ²	967	305	1,272	17.8%
2018	154	115	259	20 ²	413	136	548	15.2%
2019	24	16	0	0	24	16	40	1.8%
2020	26	42	0	0	26	42	68	2.0%
Mean	469	580	502	76	972	599	1,098	13.0%

1. Harvest rate is the total Yakima Basin harvest as a percentage of the Yakima River mouth run size.
2. Includes estimate of post-release mortality of unmarked fish.

Table 26. Estimated fall Chinook return, escapement, and harvest in the Yakima River, 1998-2020. Data from WDFW and YN databases.

Year	Total Return		Escapement				WA Recreational Harvest		
	Adult	Jack	Above Prosser Adult	Jack	Below Prosser Adult	Jack	Adult	Jack	Rate
1998	1,743	106	1,064	84	645	22	34	0	1.8%
1999	4,056	43	1,876	20	2,046	23	134	0	3.3%
2000	4,557	1,138	1,371	922	2,931	194	255	22	4.9%
2001	5,886	869	3,651	660	1,293	151	942	58	14.8%
2002	13,369	211	6,146	95	4,923	116	2,300	0	16.9%
2003	10,092	193	4,796	79	3,874	73	1,422	41	14.2%
2004	5,825	271	2,862	85	2,231	140	732	46	12.8%
2005	3,121	45	1,920	22	491	7	710	16	22.9%
2006	2,299	67	1,499	29	363	10	437	28	19.7%
2007	1,318	460	892	240	194	26	232	194	24.0%
2008	3,403	208	2,739	124	137	17	527	67	16.4%
2009	3,315	772	2,381	591	424	106	510	75	14.3%
2010	3,474	176	2,763	125	270	12	441	39	13.2%
2011	3,325	705	2,318	400	470	81	537	224	18.9%
2012	5,553	1,468	3,751	963	1098	211	704	294	14.2%
2013	13,005	1,541	8,537	995	1936	194	2,532	352	19.8%
2014	12,839	1,371	8,302	1,003	2,969	302	1,568	66	11.5%
2015	15,533	769	8,644	559	5,224	156	1,665	54	10.5%
2016	7,982	735	5,688	585	1,372	119	922	31	10.9%
2017	3,116	399	1,927	278	719	105	470	16	13.8%
2018	1,739	147	1,137	76	397	46	205	25	12.2%
2019	1,420	161	869	78	406	21	145	62	13.1%
2020	2,734	201	1,873	105	631	40	230	56	9.7%

Table 27. Estimated Coho return, escapement, and harvest in the Yakima River, 1999-2020. Data from WDFW and YN databases.

Year	Total Return		Escapement				WA Recreational Harvest		
	Adult	Jack	Prosser Dam Adult	Jack	Hatchery Denil Adult	Jack	Adult	Jack	Rate
1999	3,906	91	3,852	91			54	0	1.4%
2000	4,444	1,841	4,390	1,826			54	15	1.1%
2001	5,032	68	4,978	68			54	0	1.1%
2002	515	343	475	343			40	0	4.7%
2003	2,192	162	2,192	162			0	0	0.0%
2004	2,367	74	2,325	64			42	10	2.1%
2005	2,897	225	2,890	225			7	0	0.2%
2006	4,478	175	4,335	175	125	0	18	0	0.4%
2007	3,461	64	3,153	60	300	4	8	0	0.2%
2008	4,636	1,917	3,890	1,809	700	58	46	50	1.5%
2009	9,843	873	8,517	573	1300	300	26	0	0.2%
2010	5,776	567	4,811	183	915	384	50	0	0.8%
2011	8,073	171	6,424	121	1594	50	55	0	0.7%
2012	5,511	264	4,298	164	1200	100	13	0	0.2%
2013	3,173	848	2,290	395	837	412	46	41	2.2%
2014	25,368	584	20,997	427	4263	157	108	0	0.4%
2015	3,314	300	2,210	105	1095	195	9	0	0.2%
2016	3,383	374	1,693	188	1690	186	0	0	0.0%
2017	3,920	274	3,051	222	804	34	65	18	2.0%
2018	2,236	835	1,690	440	518	365	28	30	1.9%
2019	3,921	105	2,506	52	1361	46	54	7	1.5%
2020	3,274	3,228	2,303	524	971	2704	0	0	0.0%

Discussion:

Adult returns of spring Chinook from the CESRF have substantially increased fishing opportunity for all fishers in the Yakima Basin (Table 25) and returned recreational fisheries to the Basin after a 40-year absence. This has contributed to improved relationships between all the Basin's stakeholders and increased opportunities for collaboration.

Recreational fishers enjoy a successful annual fall Chinook fishery situated primarily near the mouth of the Yakima River (Table 26). Tribal fishers harvest a substantial, but unquantified number of Yakima Basin-destined fall Chinook (Figure 22) and coho in commercial gillnet fisheries in the Zone 6 fishing area. Because of the quantity and relatively higher quality of fall Chinook and coho available to tribal fishers in Zone 6 Columbia and Klickitat River fisheries, Yakima River tribal harvest is typically at or near zero even though regulations allowing fall season fisheries in the Yakima River are propagated annually by the Yakama Nation.

Hatchery Research

Effect of Artificial Production on the Viability of Natural Fish Populations

WDFW is addressing some critical uncertainties (see [Columbia River Basin Research Plan](#) and [Critical Uncertainties for the Columbia River Basin Fish and Wildlife Program](#)) related to genetic and ecological interactions under project [1995-064-25](#). We are working jointly with WDFW to address the following additional fish propagation uncertainties:

- 1.2. Can hatchery production programs meet adult production and harvest goals (integrated and segregated) while protecting naturally spawning populations?
- 1.4. What is the magnitude of any demographic benefit or detriment to the production of natural-origin juveniles and adults from natural spawning of hatchery-origin supplementation adults?
- 1.5. What are the range, magnitude and rates of change of natural spawning fitness of integrated (supplemented) populations, and how are these related to management rules including the proportion of hatchery fish permitted on the spawning grounds, and the proportion of natural origin adults in the hatchery broodstock?

Methods:

The YKFP began a spring Chinook salmon hatchery program at the CESRF near Cle Elum on the upper Yakima River (river kilometer 297, measuring from the confluence with the Columbia River; Figures 1 and 23) in 1997. This program is a supplementation effort targeting the upper Yakima River population and is designed to test whether artificial propagation can be used to increase natural production and harvest opportunities while limiting ecological and genetic impacts (RASP 1992). It is an integrated hatchery program (Mobrand et al. 2005) because only natural-origin brood-stock are used and returning hatchery-origin adults are allowed to spawn in the wild. The program employs “best practice” hatchery management principles (see Cuenco et al. 1993, Mobrand et al. 2005) including reduced pond densities, strict disease management protocols, random brood-stock selection, and factorial mating (Busack and Knudsen 2007) to maximize effective population size. Fish are reared at the central facility, but released from three acclimation sites located near the central facility at: Easton approximately 25km upstream of the central facility, Clark Flat about 25km downstream of the central facility, and Jack Creek about 12km upstream from the Teanaway River’s confluence with the Yakima River (Figure 23). The CESRF collected its first spring Chinook brood-stock in 1997, released its first fish in 1999, and age-4 adults have been returning since 2001. The first generation of offspring of CESRF and wild fish spawning in the wild returned as adults in 2005. The program uses the adjacent, un-supplemented Naches River population as an environmental and wild control system.

To evaluate demographic benefits for spring Chinook, we compared redd count and natural-origin adult return data for the supplemented Upper Yakima and un-supplemented (control) Naches populations using a Before/After Control/Impact (BACI) analysis (Stewart-Oaten et al. 1986; Smith et al. 1993). For redd counts, the before period was defined as 1981 to 2000 and the after period as 2001 to present (hatchery-origin age-4 adults first returned to integrate with natural-origin fish on the natural spawning grounds in 2001). The first natural-origin returns of age-4 fish from these integrated population redds did not occur until 2005, so the pre- and post-supplementation (before/after) periods for natural-origin return evaluation were defined as 1982 to 2004 and 2005 to present, respectively. The spring Chinook findings described below were published in Fast et al. (2015). We are working with WDFW to incorporate additional out-of-basin control populations in this evaluation and these results will be considered for publication at a later date.

To evaluate fitness parameters for an integrated spring Chinook population, we used methods described in Knudsen et al. (2008), Schroder et al. (2008, 2010, and 2012) and Waters et al. (2015; discussed further below under Hatchery Reform). For coho,

we conducted preliminary evaluation of both demographic benefits and some fitness parameters using methods described in Bosch et al. (2007).




Figure 23. Map of the Yakima River Basin, Cle Elum Supplementation and Research Facility (CESRF) locations, and timeline of the spring Chinook supplementation program.

Results:




Figure 24. Spring Chinook redd counts in the supplemented Upper Yakima (red bar) relative to the un-supplemented Naches (control; blue bar) for the pre- (1981-2000) and post-supplementation (2001-2020) periods.




Figure 25. Natural-Origin returns of Spring Chinook in the supplemented Upper Yakima (red bar) relative to the un-supplemented Naches (control; blue bar) for the pre- (1982-2004) and post-supplementation (2005-2020) periods.

Discussion:

Supplementation has increased spring Chinook redd abundance in the Upper Yakima relative to the Naches control system (Figure 24). Redd counts in the post-supplementation period (2001-2020) increased in the supplemented Upper Yakima (+74%; P=0.041) but the change observed in the un-supplemented Naches control system relative to the pre-supplementation period (1981-2000) was not significant (+20%; P=0.398). As noted above, spatial distribution of spring Chinook has also increased as a result of supplementation with dramatic increases in redd abundance observed in the Teanaway River (Figure 14) in some years.

Changes in mean natural-origin return abundance in the post-supplementation period (2005-2020) relative to the pre-supplementation period (1982-2004) were not significant in either the supplemented upper Yakima River (-5.5%; P=0.83; Figure 25) or the un-supplemented Naches River system (-24.6%; P=0.32; Figure 25). We have already noted that limiting factors appear to be inhibiting natural productivity (see status and trend of adult productivity) throughout the Yakima Basin.

With respect to spring Chinook fitness parameters we found the following. The relationships between reproductive traits and body length were not significantly altered by a single generation of hatchery exposure. However, because hatchery females had smaller body sizes, the distributions of linked traits, such as total gamete mass and fecundity, differed by as much as 0.6 SD, probably resulting in some fitness loss. Our data support the idea that a single generation of state-of-the-art conservation hatchery propagation can produce fish with reproductive traits similar to those of wild fish, given comparable body size (Knudsen et al. 2008). No differences were detected in the egg deposition rates of wild and hatchery origin females, but pedigree assignments based on microsatellite DNA showed that the eggs deposited by wild females survived to the fry stage at a 5.6% higher rate than those spawned by hatchery-origin females (Schroder et al. 2008). Behavior and breeding success of wild and hatchery-origin males were found to be comparable (Schroder et al. 2010). Large anadromous males produced 89%, jacks 3%, yearling precocious 7%, and sub-yearling precocious 1% of the fry in our tests suggesting that large anadromous males generate most of the fry in natural settings when half or more of the males present on a spawning ground use this life history strategy (Schroder et al 2012). For additional detail on Spring Chinook findings, see Fast et al. (2015). Finally, in addition to the relative reproductive success (RRS) results reported by Schroder et al. (2008 and 2010) for artificial spawning channel studies, we are also working with our project collaborators at WDFW and CRITFC to evaluate RRS for all integrated hatchery- and natural-origin spawners above Roza Dam for brood years 2007-2011 (see <https://www.cbfish.org/Document.mvc/Viewer/P159280> for the latest progress report on this project). Genotyping for this work has been completed and we are working to publish findings in 2021.

The YKFP is presently studying the release of over 1.0 million coho smolts annually from acclimation sites in the Naches and Upper Yakima subbasins. These fish are a combination of in-basin production from brood-stock collected in the vicinity of Prosser Dam plus out-of-basin stock generally reared at Willard or Eagle Creek National Fish Hatcheries and moved to the Yakima Subbasin for final rearing and release. Monitoring of these efforts to re-introduce a sustainable, naturally spawning coho population in the Yakima Basin have indicated that coho returns averaged over 5,600 fish from 1997-2020 (an order of magnitude improvement from the average for years prior to the project) including estimated returns of wild/natural coho averaging over 800 fish annually since 2001 (Figure 4). Coho re-introduction research has demonstrated that hatchery-origin coho, with a legacy of as many as 10 to 30 generations of hatchery-influence, can reestablish a naturalized population after as few as 3 to 5 generations of outplanting in the wild (Bosch et al. 2007). The project is working to further develop a locally adapted brood-stock and to establish specific release sites and strategies that optimize natural reproduction and survival.

Effectiveness of Hatchery Reform

Hatcheries have long been a part of the fisheries landscape in the Pacific Northwest with programs originally designed to provide abundant returns for harvest in river ecosystems that were becoming increasingly exploited to serve human needs (Lichatowich 1999). Historically, hatchery programs were designed to release a specified number of juveniles from a central facility, and adult survivors, after providing many fish for harvest during their marine and freshwater migrations, would return to swim-in ladders and adult holding ponds at that same facility to spawn successive generations. Over the past two decades or more, such programs have been the subject of much scientific study regarding risks, such as domestication, they pose to natural populations if these fish spawn in the wild.

The concepts of supplementation and hatchery reform, where hatchery programs could be (re)designed to serve conservation as well as harvest purposes, first began to appear in regional discussions and the literature in the late 1980s and early 1990s (e.g, RASP 1992; Cuenco et al. 1993). In Mobrand et al. (2005) and Paquet et al. (2011), the Hatchery Scientific Review Group (HSRG) described in more scientific detail several principles that should guide integrated (conservation-oriented) hatchery programs which purposefully allow fish to spawn in the wild (note that virtually all of the HSRG recommendations were designed into the integrated CESRF program described above). The HSRG reports also recommended that traditional, harvest-oriented hatchery programs should be segregated as much as possible from natural populations to minimize risks by limiting the number of returning fish that escape to natural spawning grounds.

YKFP efforts to monitor and evaluate hatchery reform focus on the CESRF spring Chinook program which was designed explicitly for this purpose from its inception (BPA 1996). To the extent that is practical, we will evaluate similar metrics for the summer/fall run Chinook and coho programs and publish those results in future reports as the Master Plan (Yakama Nation 2019) is implemented and the programs mature over time.

In addition to the integrated (supplementation-S) hatchery program described above for the CESRF, this facility also introduced a segregated “hatchery control” (HC) program in 2002 as recommended by independent scientific review. To protect the integrity of the integrated program evaluation described above, returning HC line fish were either harvested or trapped and removed at the Roza Adult Monitoring Facility (RAMF); no HC line fish were allowed to escape to the spawning grounds (determination of fish origin was based on a differential marking strategy for S and HC fish; unmarked fish were presumed wild). CESRF-project scientists hypothesized

that HC-line fish, which use only returning hatchery-origin fish as brood source, would increasingly diverge in phenotypic and genetic characteristics from wild (WC or wild control) fish with increasing generations of hatchery influence, whereas S-line fish, which use only wild or natural-origin fish for brood source, would remain relatively close in characteristics to wild fish (Figure 26). These hypothetical outcomes were based on hatchery reform theory which suggests that, by using only wild or natural-origin parents to spawn successive generations of fish in the hatchery environment, mean fitness of an integrated population in the natural environment can be maintained relatively close to that of a wild population (Mobernd et al. 2005).




Figure 26. Hypothetical outcomes of trait divergence (domestication effects) over time for a segregated (hatchery-control or HC) line of fish, compared to an integrated (supplementation or S) line of fish and a wild (wild-control or WC) line of fish (D. Fast, Yakama Nation).

This section reports on our efforts to evaluate the effectiveness of hatchery reform measures implemented in the CESRF program.

Methods:

Methods for enumerating natural- and CESRF-origin fish at Roza Dam were described above (Status and Trend of adult abundance) and in Knudsen et al. (2006). Methods for evaluating genetic differentiation between the wild founding, integrated, and segregated populations at the CESRF were described in Waters et al. (2015).

A recently developed parameter to monitor the mean fitness of an integrated population in the natural environment is called Proportionate Natural Influence (PNI). PNI is an approximation of the rate of gene flow between the natural environment and the hatchery environment (Busack et al. 2008). The equation describing PNI is

$$PNI = \frac{pNOB}{pNOB + pHOS}$$

where pNOB is the proportion of natural-origin brood-stock and pHOS is the proportion of hatchery-origin spawners. We evaluated PNI for the CESRF program using a pNOB value of 1.0 as only natural-origin fish were used for the integrated program's broodstock.

Results and Discussion:

For CESRF integrated program return years 2001-2020, PNI averaged 65% while pHOS averaged 53.6% (Table 28). As stated in the introduction to this report and in the final Environmental Impact Statement for the Yakima Fisheries Project (BPA 1996), one of the explicit purposes of the project is to test the assumption that new artificial propagation or hatchery reform techniques (Cuenco et al. 1993, Mobernd et al. 2005) can be used to increase natural production without causing significant impacts to existing natural populations. Therefore, it has always been the intent of this project to purposely allow integrated hatchery-origin fish to escape to the natural spawning grounds, i.e., we intentionally maintained a relatively high pHOS rate. Even with a high pHOS relative to recommendations, PNI for the CESRF integrated program remained in the "low hatchery influence for conservation of natural populations" category described by the HSRG (Paquet et al. 2011).

The project will continue to monitor PNI considering factors such as: policy input regarding controlling the number and types of fish allowed to escape to natural spawning areas, meeting overall production goals of the project, guidance from the literature relative to percentage of hatchery fish on the spawning grounds with fitness loss, considerations about what risk is acceptable in a project designed to evaluate impacts from that risk, and the numerous risk containment measures already in place in the project. The State of Washington is using mark-selective fisheries in the lower Columbia River and, when possible, in the lower Yakima River in part as a tool to manage escapement proportions. In 2011, the project implemented an effort to transfer some returning hatchery-origin CESRF adults from Roza Dam to Lake Cle Elum for the purpose of returning marine derived nutrients and salmon to the watersheds that feed the lake. These measures will also increase PNI in the major

spawning areas of the Upper Yakima Basin. Additional adaptive management measures will be considered when and if monitoring and evaluation indicates a need.

Table 28. Escapement (Roza Dam counts less brood-stock collection and harvest above Roza) of natural-(NoR) and hatchery-origin (HoR) spring Chinook to the upper Yakima subbasin, 1982 – present.

Year	Wild/Natural (NoR)			CESRF (HoR)			Total			pHOS ¹	PNI ¹
	Adults	Jacks	Total	Adults	Jacks	Total	Adults	Jacks	Total		
1982			1,146								
1983			1,007								
1984			1,535								
1985			2,331								
1986			3,251								
1987			1,734								
1988			1,340								
1989			2,331								
1990			2,016								
1991			1,583 ²								
1992			3,009								
1993			1,869								
1994			563								
1995			355								
1996			1,631								
1997	1,141	43	1,184								
1998	369	18	387								
1999	498	468	966								
2000	10,491	481	10,972		688	688	10,491	1,169	11,660	5.9%	
2001	4,454	297	4,751	6,065	982	7,047	10,519	1,279	11,798	59.7%	62.6%
2002	1,820	89	1,909	6,064	71	6,135	7,884	160	8,044	76.3%	56.7%
2003	394	723	1,117	1,036	1,105	2,141	1,430	1,828	3,258	65.7%	60.3%
2004	6,536	671	7,207	2,876	204	3,080	9,412	875	10,287	29.9%	77.0%
2005	4,401	175	4,576	627	482	1,109	5,028	657	5,685	19.5%	83.7%
2006	1,510	121	1,631	1,622	111	1,733	3,132	232	3,364	51.5%	66.0%
2007	683	161	844	734	731	1,465	1,417	892	2,309	63.4%	61.2%
2008	988	232	1,220	2,157	957	3,114	3,145	1,189	4,334	71.9%	58.2%
2009	1,843	701	2,544	2,234	2,260	4,494	4,077	2,961	7,038	63.9%	61.0%
2010	2,436	413	2,849	4,524	1,001	5,525	6,960	1,414	8,374	66.0%	60.2%
2011	3,092	926	4,018	3,162	1,404	4,566	6,254	2,330	8,584	53.2%	65.3%
2012	2,359	191	2,550	2,661	265	2,926	5,020	456	5,476	53.4%	65.2%
2013	1,708	678	2,386	1,587	840	2,427	3,295	1,518	4,813	50.4%	66.5%
2014	3,099	685	3,784	2,150	794	2,944	5,249	1,479	6,728	43.8%	69.6%
2015	3,357	163	3,520	1,779	167	1,946	5,136	330	5,466	35.6%	73.7%
2016	2,070	266	2,336	1,198	705	1,903	3,268	971	4,239	44.9%	69.0%
2017	1,135	194	1,329	1,328	660	1,988	2,463	854	3,317	59.9%	62.5%
2018	500	33	533	1,033	233	1,266	1,533	266	1,799	70.4%	58.7%
2019	316	81	397	828	266	1,094	1,144	347	1,491	73.4%	57.7%
2020	497	56	553	746	341	1,087	1,243	397	1,640	66.3%	60.1%
Mean ³	2,321	328	2,648	2,221	679	2,794	4,171	922	5,093	53.6%	64.8%

1. Proportionate Natural Influence equals Proportion Natural-Origin Brood-stock (PNOB; 1.0 as only NoR fish are used for supplementation line brood-stock) divided by PNOB plus Proportion Hatchery-Origin Spawners (PHOS).
2. This is a rough estimate since Roza counts are not available for 1991.
3. For NoR columns, mean of 1997-present values. For all other columns, mean of 2001-present values.

Both the CESRF integrated and segregated programs have now proceeded for several generations and we can evaluate actual outcomes relative to the hypothetical outcomes given in Figure 26 above. Results were presented in Waters et al. (2015) and empirically demonstrate that using managed gene flow (i.e, using only natural-origin fish for brood stock) reduced genetic divergence over time in the CESRF integrated (S-line) fish compared to the segregated (HC-line; hatchery-origin parents) fish (Figure 27). The actual results are remarkably consistent with the projected outcomes in Figure 25 demonstrating that there is considerable merit to the concepts behind hatchery reform. While some detractors of hatchery supplementation choose to highlight the differences the CESRF program has found between hatchery and natural-origin fish such as those documented in Knudsen et al. (2006 and 2008), it is important to note that integrated hatchery-origin fish were never expected to be identical to wild fish (Figure 26), but rather similar enough to increase demographic abundance of natural spawners while minimizing risk, which is exactly what the results to date for this project demonstrate (Fast et al. 2015; Koch et al. 2017). Additional evaluation is required before definitive answers to key biological cost and benefit questions relative to using this type of management over the long-term will be known with scientific certainty (Fraser 2008). The YKFP is continuing its collaboration with University of Washington and NOAA scientists to further evaluate and associate genetic divergence results from Waters et al. (2015) with the phenotypic trait analyses in Knudsen et al. (2006 and 2008).




Figure 27. Estimated genetic divergence (variation) for integrated (INT blue), segregated (SEG red), and wild founder (black) spring Chinook in the CESRF program after 4 parental-generations of the hatchery program (P1=1998, F1=2002, F2=2006, F3=2010, F4=2014; updated from Figure 4 in Waters et al. 2015).

Additional information and results from the CESRF program are provided in Appendix B and in Fast et al. (2015).

Predation Management and Predator Control

Avian Predation Index

Avian predators are capable of significantly depressing smolt production. The loss of wild spring Chinook salmon juveniles to various types of avian predators has long been suspected as a significant constraint on production and could limit the success of supplementation. Therefore, a long-standing objective of the YKFP has been to monitor, evaluate, and index the impact of avian predation on annual salmon and steelhead smolt production in the Yakima Subbasin. Accurate methods of indexing avian predation across years have been developed.

Methods:

River Reach Surveys

The spring river surveys included six river reaches (Table 29) and were generally consistent with avian point count methods described in monitoringmethods.org method [1151](#). The survey accounts for coverage of approximately 70 miles of the lower portion of the Yakima River.

Table 29. Avian predation river reach survey start and end locations and total reach length.

<i>Survey Name</i>	<i>River Mile Start</i>	<i>River Mile End</i>	<i>Survey Distance</i>
Parker	107.0	93.8	13.2
Granger-Emerald	85.3	66.5	18.8
Mabton- Prosser	60.6	48.5	12.1
Below Prosser	46.4	36.6	9.7
Chandler Power Plant -Benton	36.6	30.2	6.5
Below Horn Rapids-Van Giesen	16.8	9.4	7.4

All river reach surveys were conducted by a two-person team from a 16-foot drift boat or 12-foot raft. Surveys began between 8:00 am and 9:00 am and lasted between 2 to 6 hours depending upon the length of the reach and the water level. All surveys were conducted while actively rowing the drift boat or raft downstream to decrease

the interval of time required to traverse the reach. One person rowed the boat while the other person recorded piscivorous birds encountered.

Table 30. Yakima River Avian Predators.

Common Name	Scientific Name	Acronym
Common Merganser	<i>Mergus merganser</i>	COME
American White Pelican	<i>Pelecanus erythrorhynchos</i>	AWPE
California Gull	<i>Larus californicus</i>	GULL
Ring-billed Gull	<i>Larus delawarensis</i>	GULL
Belted Kingfisher	<i>Ceryle alcyon</i>	BEKI
Great Blue Heron	<i>Ardea herodias</i>	GBHE
Double-crested Cormorant	<i>Phalacrocorax auritus</i>	DCCO
Black-crowned Night-Heron	<i>Nycticorax nycticorax</i>	BCHE
Forster's Tern	<i>Sterna forsteri</i>	FOTE
Great Egret	<i>Ardea alba</i>	GREG
Hooded Merganser	<i>Lophodytes cucullatus</i>	HOME
Bald Eagle	<i>Haliaeetus leucocephalus</i>	BAEA
Osprey	<i>Pandion haliaetus</i>	OSPR
Caspian Tern	<i>Sterna caspia</i>	CATE

All birds detected visually or aurally were recorded, including time of observation, species, and sex and age if distinguishable. Leica 10x42 binoculars were used to help observe birds. All piscivorous birds encountered on the river were recorded at the point of initial observation. Most birds observed were only mildly disturbed by the presence of the survey boat and were quickly passed. Navigation of the survey boat to the opposite side of the river away from encountered birds minimized escape behaviors. If the bird attempted to escape from the survey boat by moving down river a note was made that the bird was being pushed. Birds being pushed were usually kept in sight until passed by the survey boat. If the bird being pushed down river moved out of sight of the survey personnel, a note was made, and the next bird of the same species/age/sex to be encountered within the next 1000 meters of river was assumed to be the pushed bird. If a bird of the same species/age/sex was not encountered in the subsequent 1000 meters, the bird was assumed to have departed the river or passed the survey boat without detection, and the next identification of a bird of the same species/age/sex was recorded as a new observation.

Avian Predator Hotspot Surveys

Two “hotspots” of avian predators have been identified within the Lower Yakima River (Figure 28). These “hotspots” consist of an area below the Chandler fish bypass outfall pipe and below Wanawish Dam. To include data about these hotspots weekly bird counts will be conducted at each of these “hotspots” by YN personnel and BOR personnel. Data will be single day counts of piscivorous birds during the early morning.

Acclimation Site Surveys

Three Spring Chinook acclimation sites in upper Yakima River (Clark Flat, Jack Creek, and Easton) were surveyed for piscivorous birds from 2004 through 2018 (Figure 1). Surveys were conducted between January 23 and June 10, though dates varied for each site. Three surveys were conducted at the Spring Chinook sites each day, at 8:00 am, 12:00 noon, and 4:00 pm. Surveys were conducted on foot. All piscivorous birds within the acclimation facility, along the length of the artificial acclimation stream, and 50 meters above and 150 meters below the acclimation stream outlet, into the main stem of the Yakima River or its tributaries, were recorded.




Figure 28. Avian Predator Survey Locations.

Results and Discussion:

River Reach Surveys

Thirteen different piscivorous bird species were observed on the Yakima River. These included: American White Pelican, Bald Eagle, Black-crowned Night Heron, Belted Kingfisher, Caspian Tern, Common Merganser, Double-crested Cormorant, Forster's Tern, Great Egret, Great Blue Heron, Gull species (California and Ringbill), Hooded Merganser, and Osprey. With the exception of the Forster's Tern, 12 of the species have been observed in most survey years. Graph Data (Figure 29) for river reach surveys represents Avian Predator totals by reach of the lower Yakima River (surveys below Wapato Dam). The total avian predators in the Parker Reach by week are represented in (Figure 30) and numbers increased as river flows decreased. The avian predator counts within the Parker, Granger, Below Prosser, Benton, and Lower Yakima reaches are represented in the bar graphs by their survey acronyms (Figures 31-35).

The Osprey, Great Blue Heron, Common Merganser, and Belted Kingfisher were observed within all six reaches in 2019 while American White Pelicans and Double Crested Cormorants have also been observed in these six reaches in prior years. Common Mergansers were the most abundant Avian Predators in the upper surveyed reaches of the river. The abundance of the Common Merganser in the upper Yakima River in 2019 and all previous years monitored suggest they are the top avian predator for the upper river while American White Pelicans are dominant at Parker and Granger (Figures 31-32).

Gull numbers in the lower Yakima River decreased in 2016 and this trend continued into 2018. In 2019, gulls were again abundant showing increased numbers below Prosser Dam and in the Benton reach at the end of May and in June. Double Crested Cormorants numbers remained consistent in 2019. DCCO numbers remain a concern due to nest takeover of Great Blue Heron Rookeries in various areas along the Yakima River along their high capacity for consuming salmon smolts. Monitoring of the Double Crested Cormorant on the river and in rookeries will be a priority in upcoming years as the Army Corp of Engineers culls and removes breeding habitat at the estuary of the Columbia River in efforts to reduce juvenile salmon predation. These actions may result in displacement and searching out of new habitat for the Cormorants and lead to impacts on salmon in other rivers and basins. The American White Pelican numbers remain consistently high in the lower Yakima River. In the Yakima River pelicans can be seen in groups of over 100 in the Wapato Reach of the river along the borders of the Yakama Indian Reservation.




Figure 29. Avian Predator Totals by Reach.




Figure 30. Parker Reach Total Avian Predators by Week.




Figure 31. Parker Reach Avian Predator Species Counts.




Figure 32. Granger Reach Avian Predator Species Counts.




Figure 33. Below Prosser Avian Predator Species Counts.




Figure 34. Benton Reach Avian Predator Species Counts.




Figure 35. Lower Yakima Reach Avian Predator Species Counts.

Hotspot Surveys

Avian predator surveys were conducted at the Chandler fish bypass pipe (river mile ~46; Figure 36) and Wanawish Dam (river mile ~18.5; Figure 37) hotspots. In 2019 there was an increase in avian predators at both hotspot locations. At Chandler the species diversity stayed the same, where at Wanawish dam there was a decrease in diversity. Only three species were observed at Wanawish in 2019, American White Pelican, Double Crested Cormorant, and Gulls.




Figure 36. Avian Predator Counts at Chandler “hotspot”.




Figure 37. Avian Predator Counts at Wanawish Dam “hotspot”.

Acclimation Sites Surveys

At the three Spring Chinook salmon acclimation sites in the upper Yakima River and its tributaries piscivorous bird surveys were conducted over a 3-5 month period in the winter and spring of 2019. The most common species of birds observed at acclimation sites were Bald Eagle, Belted Kingfisher, Common Merganser, Great Blue Heron, Great Egret, and Osprey. Using the assumption that birds frequenting

acclimation ponds are only consuming acclimating juvenile salmon, an average consumption rate can be determined. The average consumption rate can be calculated using the average number of birds at each site, daily energy requirements of the birds and the average size of juvenile salmon.

It was estimated that these bird species together consumed 786 juvenile Chinook at Clark Flat (Table 31). Great Blue Herons had the highest consumption rate, consuming 545 juvenile Chinook. At Easton, it was estimated that 375 juvenile Chinook were consumed. Great Blue Herons and Bald Eagles had the highest consumption rates. Great Blue Herons consumed 122 juvenile Chinook and Bald Eagles consumed 188 juvenile Chinook. Only Belted Kingfishers and Common Mergansers were observed at Jack Creek. It was estimated that they consumed 151 juvenile Chinook. Common Mergansers consumed 137 juvenile Chinook. In 2018, these bird species together consumed 950 juvenile Chinook at Clark Flat, 339 juvenile Chinook at Easton and 961 juvenile Chinook at Jack Creek.

Table 31. Estimated consumption in 2019 by Avian species at three spring Chinook Salmon acclimation sites.

2019 SPRING CHINOOK ACCLIMATION SITES				
CLARK FLAT				
	AVG. # OF BIRDS	# FISH EATEN BY SPECIES	% OF FISH EATING BY SPECIES	% OF TOTAL FISH EATEN BY SITE
BAEA	0.015224359	57	7	0.023337223
BEKI	0.303685897	111	14	0.045446171
COME	0.004807692	14	2	0.005731949
GBHE	0.212339744	545	69	0.223136605
GREG	0.008012821	7	1	0.002865975
OSPR	0.024038462	52	7	0.021290098
TOTAL	0.568108974	786	100	0.321808021
EASTON				
	AVG. # OF BIRDS	# FISH EATEN BY SPECIES	% OF FISH EATING BY SPECIES	% OF TOTAL FISH EATEN BY SITE
BAEA	0.050595238	188	50	0.080529093
COME	0.007936508	22	6	0.009423617
GBHE	0.047619048	122	33	0.052258241
OSPR	0.01984127	43	11	0.018418888
TOTAL	0.125992063	375	100	0.16062984
JACK CREEK				
	AVG. # OF BIRDS	# FISH EATEN BY SPECIES	% OF FISH EATING BY SPECIES	% OF TOTAL FISH EATEN BY SITE
BEKI	0.056818182	14	9	0.006006418
COME	0.071969697	137	91	0.058777093
TOTAL	0.128787879	151	100	0.064783512

Fish Predation Index and Predator Control

Fish predators are also capable of significantly depressing smolt production. Thus the YKFP has a long-established objective to monitor, evaluate, and manage the impact of piscivorous fish on annual smolt production of Yakima Subbasin salmon and

steelhead. By indexing the mortality rate of upper Yakima spring Chinook attributable to piscivorous fish in the lower Yakima River, the contribution of in-basin predation to variations in hatchery- and natural-origin spring Chinook smolt-to-adult survival rate can be deduced.

Based on YKFP and WDFW studies of piscivorous fish in the Yakima River Basin (Fritts and Pearsons 2004, 2006, 2008), it was determined that management of the piscivorous fish populations in the area is necessary to improve survival of juvenile salmonids. Initial steps were taken in 2009 to identify locations that would be suitable for a multi-pass removal population study. In early 2010, the YKFP began initial study checks to determine management and study goals for piscivorous fish. Presence and absence of piscivorous fish was determined through electro-fishing various sections of the Yakima River to determine temporal and spatial trends of each species of piscivorous fish. On March 1, 2013, the [Washington Fish and Wildlife Commission](#) adopted numerous changes to sport fishing rules, including the elimination of catch restrictions for non-native predators.

Methods:

Data was collected on piscivorous fish from six electrofishing sites within the Yakima River (Figure 38). Sites were sampled via boat electrofishing through time to assess spatial and temporal patterns of fish abundance and distribution. Each sampling segment was defined by river features of dams and boat launches. The partitioned sample locations consist of four ten mile surveys, one four-mile survey, and one six mile survey (Table 32). Total river mile distance of the combined Yakima River surveys is 50 miles. Survey locations were marked by GPS unit (Garmin GPSmap 78; Garmin International, Olathe, Kansas). After marking sampling reaches, we sample weekly beginning April 2nd and ending June 22nd (dates may vary depending on river stage). (Fish Predators Schei, monitoring methods [47](#)), (Predator Reduction Mclellan, monitoring methods [438](#)).

Sampling was conducted using three different types of vessels and electrofisher; 1. For five of the Yakima River surveys sampling were conducted using a Smith Root SR-16H Electrofishing boat equipped with the 7.5 GPP electrofishing unit powered by a 6,000-W Kohler boat generator in; 2. For the Yakima River survey below Prosser sampling was conducted with a 13 foot raft equipped with a smith root 1.5-KVA electrofisher powered by Honda EU2000i generator; 3. For the survey in the McNary pool sampling was conducted with a 16 foot aluminum jet boat equipped with a Smith Root VVP-15B electrofisher powered by a Honda EM3500S generator. Electro-fishing settings were adjusted to continuous DC for an output of approximately 700 V and 9–12 A. Invasive species monitoring for the Yakima River will be used as an aid

for tracking changes in fish populations and abundance as the area experiences global climate change.




Figure 38. Fish Predator Survey Locations.

Table 32. Fish Predator Survey River Miles and Distances.

<i>Survey Name</i>	<i>River Mile Start</i>	<i>River Mile End</i>	<i>Survey Distance Miles</i>
Parker	106.1	96.1	10
Granger	85.3	75.3	10
Above Prosser	52.4	48.4	4
Below Prosser	46.4	40.4	6
Benton	31.1	21.1	10
Lower Yakima	13.8	3.8	10

Sampling was conducted continuously along river margins when possible. As river stage changes, limiting access to areas within survey segments, continuous electrofishing was not always possible. The start and endpoints of shocker operation within the segment at low river stages was marked, resulting in discontinuous, marked sub-segments of electrofisher operation within each survey area.

Data collected during each sampling event consisted of:

- Water Temperature, Dissolved Oxygen, Specific Conductivity gathered by a HACH 30qd water multi-meter
- Water Turbidity gathered by a HACH TSS Handheld Instrument
- River CFS gathered from Bureau of Reclamation gaging stations
- Electrode start and end times
- Numbers and species (Table 33) of all fish observed and their size class greater than or less than 100mm

At the start of each sampling event a small group of fish were caught and examined to insure that electro-fishing settings were not causing visible injuries. To further insure injuries to fish were minimized, sampling procedures by the National Marine Fisheries Service, “Guidelines for Electrofishing Waters Containing Salmonids Listed under the Endangered Species Act,” were followed.

Table 33. Yakima River Fish Species (Note: Spring Chinook and Coho total counts are combined in results as SP+CO).

Family	Common Name	Scientific Name	Acronym
Salmonidae:	Steelhead/Rainbow trout	<i>Oncorhynchus mykiss</i>	STH
	Coho Salmon	<i>Oncorhynchus kisutch</i>	COHO*
	Chinook Salmon	<i>Oncorhynchus tshawytscha</i>	SPCK/FAK*
	Mountain Whitefish	<i>Prosopium williamsoni</i>	WT
Cyprinidae:	Chiselmouth	<i>Acrocheilus alutaceus</i>	CH
	Carp	<i>Cyprinus carpio</i>	CP
	Peamouth	<i>Mylocheilus caurinus</i>	PEA
	Speckled Dace	<i>Rhinichthys osculus</i>	SPDA
	Northern Pikeminnow	<i>Ptychocheilus oregonensis</i>	NPM
	Redside Shiner	<i>Richardsonius balteatus</i>	SH
Catostomidae:	Sucker	<i>Catostomus columbianus</i> <i>Catostomus catostomus</i>	SK
Ictaluridae:	Brown Bullhead	<i>Ameiurus nebulosus</i>	BRCT
	Channel Catfish	<i>Ictalurus punctatus</i>	CHCT
Centrarchidae:	Pumpkin Seed	<i>Lepomis gibbosus</i>	PKSC
	Blue Gill	<i>Lepomis macrochirus</i>	BG
	Smallmouth Bass	<i>Micropterus dolomieu</i>	SMB
	Large Mouth Bass	<i>Micropterus salmoides</i>	LMB
	Black Crappie	<i>Pomoxis nigromaculatus</i>	CRAP
Percidae:	Walleye	<i>Stizostedion vitreum vitreum</i>	WALLEYE
	Yellow Perch	<i>Perca flavescens</i>	YP
Cottidae:	Sculpin	<i>Cottus bairdi</i>	SC
Clupeidae:	Shad	<i>Alosa sapidissima</i>	SHAD

Results and Discussion:

During surveys of 2018 to 2019 the highest abundance of non-native fish predators were found in the lower reaches of the Yakima River (Figure 39). Piscivorous fish were identified in all 6 survey reaches of the Yakima River. Smallmouth Bass and Channel Catfish were the fish predators found in the highest abundance. These two predators are often considered to be the top salmon predators in the lower Yakima River.

Northern Pike Minnow are the dominant piscivorous fish in the upper portion of the 2019 surveyed reaches of Yakima River (reaches above Prosser Dam). They were the fish predator found in the highest abundance in this area during electro-fishing surveys of 2019. Fish counts for all species observed during the 2019 surveys are given for all reaches in figures 40 through 45.




Figure 39. Fish Predator Counts by Reach and Species.




Figure 40. Parker Reach Fish Counts by Species.




Figure 41. Granger Reach Fish Counts by Species.




Figure 42. Above Prosser Dam Fish Counts by Species.




Figure 43. Below Prosser Dam Fish Counts by Species.




Figure 44. Benton Reach Fish Counts by Species.




Figure 45. Lower Yakima Reach Fish Counts by Species.

Large amounts of introduced fish predators inhabit the Lower Yakima River. Predator numbers tend to increase as time progresses in the spring and summer. Increases in predator abundance in 2019 showed significant correlation with increasing date (Figure 46). These increases also correspond with increasing water temperatures and decreasing river flows.




Figure 46. Total Count of Fish Predators below Prosser Dam.

Smallmouth Bass (SMB) have been found to exhibit a spike in abundance during their spawning periods in the Lower Yakima River. Spawning for Smallmouth Bass is typically between April 1 and July 1. This time period coincides with juvenile salmonid outmigration. This timing provides a readily available prey source for the adult spawning bass and their young recruits. Catch and catch per unit effort for adult Smallmouth Bass begins to rise in the May and June survey periods (Figure 47) as Smallmouth Bass migrate from the Columbia River into the Yakima River to spawn. A rise in catch in adults also correlates with a rise in Yakima River water temperature (Figure 48).

The rise and fall of SMB relative abundance may correlate with the water year of 2015 which produced extremely low flows and high water temperatures and the subsequent high water year in 2016, 2017, and 2018. It is the increase in water temperature in the lower Yakima River which is thought to create productive habitat for SMB. Overall

years there is increased catch success during the late summer and fall months and electro-fishing efforts are increased to maximize catch for managing numbers of SMB in the lower Yakima River. Current efforts to increase salmon populations target SMB populations for management in hopes to increase survival of juvenile salmon outmigration.




Figure 47. Adult Smallmouth Bass Totals by Reach.




Figure 48. Adult and Juvenile Smallmouth Bass Total below Prosser Dam.

Adaptive Management and Lessons Learned

As noted extensively throughout this report, this project is a collaborative effort involving many agencies, boards, and individuals. As such, project coordination and review of project standards and protocols occurs continually amongst tribal, state, federal, and local entities during normal day-to-day operations of the project. Project results are communicated broadly through the annual [science and management conference](#), technical reports and peer-reviewed journal publications (see references and project-related publications), and via several related web sites described in Appendix A.

We support the principles established in Mobrand et al. (2005) and Paquet et al. (2011) that hatchery programs should be well-defined, scientifically defensible, and use informed decision making tools including adaptive management. Many of these principles were initially published in Cuenco et al. (1993) including specific recommended decision criteria, management protocols, release strategies, and risk management strategies for hatchery programs. We designed a number of these protocols and strategies into the CESRF program and they are clearly contributing to

the results documented here for the Upper Yakima River Basin spring Chinook populations.

Results to date from Yakama Nation supplementation and research efforts in the Yakima River Basin indicate several lessons that may be of broader application on the regional scale.

1. We need to be realistic. Can or should we expect to see “self-sustaining natural populations” in river systems that have been highly altered from their historical state due to ever-increasing human demands on shared resources? In the highly altered systems we live and work in today, hatchery programs provide a necessary means to ameliorate some of the effects of human population growth and development.
2. We need to be honest. Hatchery programs are not the cause of poor productivity. The historical record is replete with documentation (see Dompier 2005) that the region knew exactly what it was doing to natural salmon productivity when development of the region began to intensify with implementation of the Federal Columbia River Power System as early as the 1930s.
3. We need to be patient. Hatchery reform is a relatively new concept and results for longer term 20-25 year efforts such as the Idaho Supplementation Studies (ISS; Venditti et al. 2017) and CESRF program (Fast et al. 2015) are only now becoming available. These programs empirically support the idea that hatchery reform principles can provide the expected benefits.
4. While hatchery supplementation has demonstrated increases in natural production (increased redd and juvenile abundance), supplementation by itself cannot and was never intended to increase natural productivity. To accommodate expanding human population growth and resource demand, it is imperative that we continue and even increase habitat restoration actions to ensure that sufficient spawning and rearing habitat remains available to all naturally spawning fish.
5. Every subbasin, species, and study is unique, so we should not be surprised to see differing results from the many studies of hatchery effects that are ongoing. Researchers need to continue efforts to better understand the root causes of poor natural productivity and the extent to which hatchery programs effect productivity.
6. Evaluation of hatchery programs should include evaluation of environmental and other factors so that hatchery effects are properly reported.

7. Hatchery programs should be regularly evaluated at the local level using expertise across disciplines to collaboratively and iteratively develop appropriate solutions that address the unique problems and limiting factors encountered in each subbasin or tributary that hosts a hatchery program. In the Yakima Basin, this is achieved with the annual [Yakima Basin Aquatic Science and Management Conference](#), and we use the results to evaluate existing goals, objectives, and strategies and to adaptively manage projects in response to new information.

References and Project-related Publications

- Beckman, B.R., D.A. Larsen, B. Lee-Pawlak, and W.W. Dickhoff. 1998. Relation of Fish Size and Growth Rate to Migration of Spring Chinook Salmon Smolts. *North American Journal of Fisheries Management* 18:537-546.
- Beckman, B.R., D.A. Larsen, C. Sharpe, B. Lee-Pawlak, C.B. Schreck, and W.W. Dickhoff. 2000. Physiological Status of Naturally Reared Juvenile Spring Chinook Salmon in the Yakima River: Seasonal Dynamics and Changes Associated with Smolting. *Transactions of the American Fisheries Society* 129:727-753.
- Beckman, B.R. and D.A. Larsen. 2005. Upstream Migration of Minijack (Age-2) Chinook Salmon in the Columbia River: Behavior, Abundance, Distribution, and Origin. *Transactions of the American Fisheries Society* 134:1520-1541.
- Beckman, B.R., B. Gadberry, P. Parkins, and D.A. Larsen. 2008. The Effect of Yakima River Spring Chinook Salmon Sire Life History Type on Emergence Timing and Size of Progeny. *Transactions of the American Fisheries Society* 137:1285-1291.
- Bosch, W.J. 2004. The promise of hatchery-reared fish and hatchery methodologies as tools for rebuilding Columbia Basin salmon runs: Yakima Basin overview. *American Fisheries Society Symposium* 44:151-160.
- Bosch, W.J., T.H. Newsome, J.L. Dunnigan, J.D. Hubble, D. Neeley, D.T. Lind, D.E. Fast, L.L. Lamebull, and J.W. Blodgett. 2007. Evaluating the Feasibility of Reestablishing a Coho Salmon Population in the Yakima River, Washington. *North American Journal of Fisheries Management* 27:198-214.
- BPA (Bonneville Power Administration). 1996. Yakima Fisheries Project. Final Environmental Impact Statement. Bonneville Power Administration. Washington Department of Fish and Wildlife. Yakama Indian Nation. January, 1996. DOE/EIS-0169. DOE/BP-2784. Portland, OR.
- Busack, C., T. Pearsons, C. Knudsen, S. Phelps, Washington Department of Fish and Wildlife, B. Watson, M. Johnston, Yakama Nation, U.S. Department of Energy, Bonneville Power Administration, Division of Fish and Wildlife. 1997. Yakima

Fisheries Project spring Chinook supplementation monitoring plan. Project Number 195-065, Contract Number DE-BI79-1996 BPA64878. <https://pisces.bpa.gov/release/documents/documentviewer.aspx?doc=64878-1>

- Busack, C. and C.M. Knudsen. 2007. Using factorial mating designs to increase the effective number of breeders in fish hatcheries. *Aquaculture* 273:24-32.
- Busack, C., C.M. Knudsen, G. Hart, and P. Huffman. 2007. Morphological Differences Between Adult Wild and First-Generation Hatchery Upper Yakima River Spring Chinook Salmon. *Transactions of the American Fisheries Society* 136:1076-1087.
- Busack, C, P. Hulett, T. Pearsons, J. Tipping and J. B. Scott, Jr. 2008. Chapter 4 Artificial production. in J. B. Scott, Jr. and W. T. Gill, editors. *Oncorhynchus mykiss*: Assessment of Washington State's Steelhead populations and programs. Washington Department of Fish and Wildlife, Olympia, Washington. (<https://wdfw.wa.gov/publications/00150/wdfw00150.pdf>).
- Chapman, D., and eight co-authors. 1994. Status of summer/fall Chinook salmon in the Mid-Columbia Region. Don Chapman Consultants, Boise, ID. 412 pp.
- Clune, T. and D. Dauble. 1991. The Yakima/Klickitat Fisheries Project: A Strategy for Supplementation of Anadromous Salmonids. *Fisheries* 16: 28-34.
- CRITFC (Columbia River Inter-Tribal Fish Commission). 1995. Wy-Kan-Ush-Mi Wa-Ksih-Wit (Spirit of the Salmon). Columbia River Anadromous Fish Restoration Plan of the Nez Perce, Umatilla, Warm Springs and Yakama Tribes. Portland, OR.
- Cuenca, M.L., T.W.H. Backman, and P.R. Mundy. 1993. The use of supplementation to aid in natural stock restoration. Pages 269-293 in J.G. Cloud and G.H. Thorgaard, editors. *Genetic conservation of salmonid fishes*. Plenum Press, New York.
- Currens, K.P., and C.A. Busack. 1995. A framework for assessing genetic vulnerability. *Fisheries* 20:24-31.
- Dittman, A.H., D. May, D.A. Larsen, M.L. Moser, M. Johnston, and D. Fast. 2010. Homing and spawning site selection by supplemented hatchery- and natural-origin Yakima River spring Chinook salmon. *Transactions of the American Fisheries Society* 139:1014-1028.
- Dompier, D. W. 2005. *The Fight of the Salmon People: Blending Tribal Tradition with Modern Science to Save Sacred Fish*. Xlibris Corporation, www.Xlibris.com.
- Evans, A.F. and N.J. Hostetter. 2012. Systemwide evaluation of avian predation on juvenile salmonids from the Columbia River based on recoveries of Passive Integrated Transponder tags. *Transactions of the American Fisheries Society* 141:975-989.

- Fast, D.E., D. Neeley, D.T. Lind, M.V. Johnston, C.R. Strom, W.J. Bosch, C.M. Knudsen, S.L. Schroder, and B.D. Watson. 2008. Survival Comparison of Spring Chinook Salmon Reared in a Production Hatchery under Optimum Conventional and Seminatural Conditions. *Transactions of the American Fisheries Society* 137:1507–1518.
- Fast, D.E., W.J. Bosch, M.V. Johnston, C.R. Strom, C.M. Knudsen, A.L. Fritts, G.M. Temple, T.N. Pearsons, D.A. Larsen, A.H. Dittman, and D. May. 2015. A Synthesis of Findings from an Integrated Hatchery Program after Three Generations of Spawning in the Natural Environment. *North American Journal of Aquaculture* 77:377-395.
- Fiander, W., D.E. Fast, and W.J. Bosch (editors). 2019. Yakima-Klickitat Fisheries Project Monitoring and Evaluation – Yakima Subbasin, [Final Report for the performance period May/2018-April/2019](#), Project number 1995-063-25, 275 electronic pages.
- Fraser, D. J. 2008. How well can captive breeding programs conserve biodiversity? A review of salmonids. *Evolutionary Applications*, 1:535-586.
- Fritts, A.L., and T.N. Pearsons. 2004. Smallmouth bass predation on hatchery and wild salmonids in the Yakima River, Washington. *Transactions of the American Fisheries Society* 133:880-895.
- Fritts, A.L. and T.N. Pearsons. 2006. Effects of Predation by Nonnative Smallmouth Bass on Native Salmonid Prey: the Role of Predator and Prey Size. *Transactions of the American Fisheries Society* 135:853-860.
- Fritts, A.L., J.L. Scott, and T.N. Pearsons. 2007. The effects of domestication on the relative vulnerability of hatchery and wild spring Chinook salmon to predation. *Canadian Journal of Fisheries and Aquatic Sciences* 64:813-818.
- Fritts, A.L., and T.N. Pearsons. 2008. Can nonnative smallmouth bass, *Micropterus dolomieu*, be swamped by hatchery fish releases to increase juvenile Chinook salmon, *Oncorhynchus tshawytscha*, survival? *Environmental Biology of Fishes* 83:485–494.
- Greene, C.H., B.A. Block, D. Welch, G. Jackson, G.L. Lawson, E.L. Rechisky. 2009. Advances in conservation oceanography: New tagging and tracking technologies and their potential for transforming the science underlying fisheries management. *Oceanography*. Vol. 22, no. 1, pp 210-223.
- Ham, K.D., and T.N. Pearsons. 2000. Can reduced salmonid population abundance be detected in time to limit management impacts? *Canadian Journal of Fisheries and Aquatic Sciences* 57:17-24.
- Ham, K.D., and T.N. Pearsons. 2001. A practical approach for containing ecological risks associated with fish stocking programs. *Fisheries* 25(4):15-23.

- Hiebert, S., L.A. Helfrich, D.L. Weigmann, and C. Liston. 2000. Anadromous Salmonid Passage and Video Image Quality under Infrared and Visible Light at Prosser Dam, Yakima River, Washington. *North American Journal of Fisheries Management* 20:827-832.
- Hubble J., T. Newsome, and J. Woodward. 2004. [Yakima Coho Master Plan](#). Prepared by Yakama Nation in cooperation with Washington State Department of Fish and Wildlife. September 2004. Yakima Klickitat Fisheries Project, Toppenish, WA.
- Independent Scientific Advisory Board (ISAB). 2015. Density Dependence and its Implications for Fish Management and Restoration Programs in the Columbia River Basin. Northwest Power and Conservation Council, Portland, OR. Available at: <http://www.nwcouncil.org/media/7148891/isab2015-1.pdf>.
- Independent Scientific Review Panel (ISRP). 2011. Retrospective Report 2011. Northwest Power and Conservation Council, Portland, OR. Available at: <http://www.nwcouncil.org/library/isrp/isrp2011-25.pdf>.
- Johnson, C.L., G.M. Temple, T.N. Pearsons, and T.D. Webster. 2009. An Evaluation of Data Entry Error and Proofing Methods for Fisheries Data. *Transactions of the American Fisheries Society* 138:593-601.
- Kiffney, P.M., E.R. Buhle, S.M. Naman, G.R. Pess, and R.S. Klett. 2014. Linking resource availability and habitat structure to stream organisms: an experimental and observational assessment. *Ecosphere*, 5(4):39. Available at: <http://www.esajournals.org/doi/pdf/10.1890/ES13-00269.1>.
- Knudsen, C.M., S.L. Schroder, C.A. Busack, M.V. Johnston, T.N. Pearsons, W.J. Bosch, and D.E. Fast. 2006. Comparison of Life History Traits between First-Generation Hatchery and Wild Upper Yakima River Spring Chinook Salmon. *Transactions of the American Fisheries Society* 135:1130–1144.
- Knudsen, C.M., S.L. Schroder, C. Busack, M.V. Johnston, T.N. Pearsons, and C.R. Strom. 2008. Comparison of Female Reproductive Traits and Progeny of First-Generation Hatchery and Wild Upper Yakima River Spring Chinook Salmon. *Transactions of the American Fisheries Society* 137:1433-1445.
- Knudsen, C.M., M.V. Johnston, S.L. Schroder, W.J. Bosch, D.E. Fast, and C.R. Strom. 2009. Effects of passive integrated transponder tags on smolt-to-adult recruit survival, growth, and behavior of hatchery spring Chinook salmon. *North American Journal of Fisheries Management* 29:658-669.
- Larsen, D.A., B.R. Beckman, K.A. Cooper, D. Barrett, M. Johnston, P. Swanson, and W.W. Dickhoff. 2004. Assessment of High Rates of Precocious Male Maturation in a Spring Chinook Salmon Supplementation Hatchery Program. *Transactions of the American Fisheries Society* 133:98-120.

- Larsen, D.A., B.R. Beckman, C.R. Strom, P.J. Parkins, K.A. Cooper, D.E. Fast, and W.W. Dickhoff. 2006. Growth Modulation Alters the Incidence of Early Male Maturation and Physiological Development of Hatchery-Reared Spring Chinook Salmon: A Comparison with Wild Fish. *Transactions of the American Fisheries Society* 135:1017-1032.
- Larsen, D.A., B.R. Beckman, and K.A. Cooper. 2010. Examining the Conflict between Smolting and Precocious Male Maturation in Spring (Stream-Type) Chinook Salmon. *Transactions of the American Fisheries Society* 139:564-578.
- Larsen, D.A., D.L. Harstad, C.R. Strom, M.V. Johnston, C.M. Knudsen, D.E. Fast, T.N. Pearsons, and B.R. Beckman. 2013. Early Life History Variation in Hatchery- and Natural-Origin Spring Chinook Salmon in the Yakima River, Washington. *Transactions of the American Fisheries Society* 142:2, 540-555.
- Lichatowich, J., L. Mobrand, L. Lestelle, and T. Vogel. 1995. An approach to the diagnosis and treatment of depleted Pacific Salmon populations in Pacific Northwest watersheds. *Fisheries* 20:10-18.
- Lichatowich, J.A. 1999. *Salmon Without Rivers: A History of the Pacific Salmon Crisis*. Washington D.C. Island Press.
- Loeffel, R. E., and H. O. Wendler. 1968. Review of the Pacific coast chinook and coho salmon resources with special emphasis on the troll fishery. Prepared by the U.S. working group of the Informal Committee on Chinook and Coho, 107 p. (Available from Oregon Department of Fish and Wildlife, P.O. Box 59, Portland, OR 97207.)
- Major, W.W. III, J.M. Grassley, K.E. Ryding, C.E. Grue, T.N. Pearsons, D.A. Tipton, and A.E. Stephenson. 2005. Abundance and consumption of fish by California gulls and ring-billed gulls at water and fish management structures within the Yakima River, Washington. *Waterbirds* 28:366-377.
- Martin, S.W., J.A. Long, and T.N. Pearsons. 1995. Comparison of survival, gonad development, and growth between rainbow trout with and without surgically implanted dummy radio transmitters. *North American Journal of Fisheries Management* 15:494-498.
- McElhany, P., M. H. Ruckelhaus, M. J. Ford, T. C. Wainwright and E. P. Bjorkstedt. 2000. Viable salmonid populations and the recovery of evolutionarily significant units. U. S. Department of Commerce, NOAA Technical Memorandum NMFS-NWFSC-42, 156 pp. Available at: http://docs.lib.noaa.gov/noaa_documents/NMFS/NWFSC/TM_NMFS_NWFSC/TM_NMFS_NWFSC_42.pdf
- McMichael, G.A. 1993. Examination of electrofishing injury and short-term mortality in hatchery rainbow trout. *North American Journal of Fisheries Management* 13:229-233.

- McMichael, G.A., C.S. Sharpe, and T.N. Pearsons. 1997. Effects of residual hatchery-reared steelhead on growth of wild rainbow trout and spring chinook salmon. *Transactions of the American Fisheries Society* 126:230-239.
- McMichael, G.A., and T.N. Pearsons. 1998. Effects of wild juvenile spring chinook salmon on growth and abundance of wild rainbow trout. *Transactions of the American Fisheries Society* 127:261-274.
- McMichael, G.A., A.L. Fritts, and T.N. Pearsons. 1998. Electrofishing injury to stream salmonids: injury assessment at the sample, reach, and stream scales. *North American Journal of Fisheries Management* 18:894-904.
- McMichael, G.A., T.N. Pearsons, and S.A. Leider. 1999. Behavioral interactions among hatchery-reared steelhead smolts and wild *Oncorhynchus mykiss* in natural streams. *North American Journal of Fisheries Management* 19:948-956.
- McMichael, G.A., T.N. Pearsons, and S.A. Leider. 1999. Minimizing ecological impacts of hatchery-reared juvenile steelhead trout on wild salmonids in a Yakima Basin watershed. Pages 365-380 in E.E. Knudson, C.R. Steward, D.D. MacDonald, J.E. Williams, and D.W. Reiser editors. *Sustainable fisheries management: Pacific salmon*. CRC Press, Boca Raton, FL.
- McMichael, G.A. and T.N. Pearsons. 2001. Upstream movement of residual hatchery steelhead into areas containing bull trout and cutthroat trout. *North American Journal of Fisheries Management* 21:517-520.
- Milbrink, G., T. Vrede, L.J. Tranvik, and E. Rydin. 2011. Large-scale and long-term decrease in fish growth following the construction of hydroelectric reservoirs. *Canadian Journal of Fisheries and Aquatic Sciences*, 68:2167-2173.
- Mobrand, L.E., J. Barr, L. Blankenship, D.E. Campton, T.T.P. Evelyn, T.A. Flagg, C.V.W. Mahnken, L.W. Seeb, P.R. Seidel, and W.W. Smoker. 2005. Hatchery Reform in Washington State: Principles and Emerging Issues. *Fisheries* 30:11-23.
- Murdoch, A.R., P.W. James, and T.N. Pearsons. 2005. Interactions between rainbow trout and bridgelip suckers spawning in a small Washington stream. *Northwest Science* 79: 120-130.
- Neeley, D. 2010. 2009 Annual Report: Chandler Certification for Yearling Outmigrating Spring Chinook Smolt. Appendix D in Sampson, Fast, and Bosch, [Yakima/Klickitat Fisheries Project Monitoring and Evaluation](#), Final Report for the Performance Period May 1, 2009 through April 30, 2010. Yakama Fisheries, Toppenish, WA.
- Neeley, D. 2012a. Prosser-Passage Estimation Issues. Appendix F in Sampson, Fast, and Bosch, [Yakima/Klickitat Fisheries Project Monitoring and Evaluation](#), Final Report for the Performance Period May 1, 2011 through April 30, 2012. Yakama Fisheries, Toppenish, WA.

- Neeley, D. 2012b. 2011 Annual Report: Smolt-to-smolt survival to McNary Dam of Yakima fall and summer Chinook. Appendix G in Sampson, Fast, and Bosch, [Yakima/Klickitat Fisheries Project Monitoring and Evaluation](#), Final Report for the Performance Period May 1, 2011 through April 30, 2012. Yakama Fisheries, Toppenish, WA.
- NMFS. 1999a. Endangered Species Act - Section 7 Consultation - Supplemental Biological Opinion and Incidental Take Statement. The Pacific Coast Salmon Plan and Amendment 13 to the Plan. NMFS, Protected Resources Division. April 28, 1999. 39 pp. + attachment.
- NMFS. 1999b. Endangered Species Act – Reinitiation of Section 7 Consultation – Biological Opinion and Incidental Take Statement. The Fishery Management Plan for Commercial and Recreational Fisheries off the Coasts of Washington, Oregon, and California of the Pacific Fishery Management Council. NMFS. Sustainable Fisheries Division. April 30, 1999. 46 pp.
- NMFS 1999c. Endangered Species Act - Reinitiated Section 7 Consultation - Approval of the Pacific Salmon Treaty by the U.S. Department of State and Management of the Southeast Alaska Salmon Fisheries Subject to the Pacific Salmon Treaty. NMFS, Protected Resources Division. November 9, 1999. 90 p. + figures.
- NMFS. 1999d. Endangered and threatened species; threatened status for three Chinook salmon Evolutionarily Significant Units (ESUs) in Washington and Oregon, and endangered status for one Chinook salmon ESU in Washington. Federal Register 64: 56 (March 24, 1999) 14308-14328. Available at: <http://www.nwr.noaa.gov/ESA-Salmon-Listings/Salmon-Populations/Federal-Register-Notices.cfm>.
- NMFS. 2000a. Endangered Species Act - Reinitiated Section 7 Consultation - Effects of Pacific coast ocean and Puget Sound salmon fisheries during the 2000-2001 annual regulatory cycle. NMFS, Protected Resources Division. April 28, 2000. 99 pp.
- NMFS. 2000b. Endangered Species Act - Reinitiated Section 7 Consultation – Biological Opinion and Incidental Take Statement. Effects of Pacific Coast Salmon Plan on California Central Valley spring-run Chinook, and California coastal Chinook salmon. NMFS, Protected Resources Division. April 28, 2000. 31 pp.
- NMFS. 2000c. RAP – A risk assessment procedure for evaluating harvest mortality on Pacific Salmonids. Sustainable Fisheries Division, NMFS, Northwest Region and Resource Utilization and Technology Division, NMFS, Northwest Fisheries Science Center. May 23, 2000. 33 p.

- Northwest Power and Conservation Council. Columbia River Basin Fish and Wildlife Program 2017 Research Plan, Pre-publication version. Available online: <https://www.nwcouncil.org/media/7491163/2017-4.pdf>. Portland, Oregon.
- NRC (National Research Council). 1996. Upstream: Salmon and society in the Pacific Northwest. National Academy Press, Washington D.C.
- Paquet, P. J., T. Flagg, A. Appleby, J. Barr, L. Blankenship, D. Campton, M. Delarm, T. Evelyn, D. Fast, J. Gislason, P. Kline, D. Maynard, L. Mobrand, G. Nandor, P. Seidel, and S. Smith. 2011. Hatcheries, Conservation, and Sustainable Fisheries—Achieving Multiple Goals: Results of the Hatchery Scientific Review Group's Columbia River Basin Review. *Fisheries* 36:11, 547-561.
- Pearsons, T.N., and A.L. Fritts. 1999. Maximum size of chinook salmon consumed by juvenile coho salmon. *North American Journal of Fisheries Management* 19:165-170.
- Pearsons, T.N., and C.W. Hopley. 1999. A practical approach for assessing ecological risks associated with fish stocking programs. *Fisheries* 24(9):16-23.
- Pearsons, T.N. 2002. Chronology of ecological interactions associated with the life-span of salmon supplementation programs. *Fisheries* 27(12):10-15.
- Pearsons, T.N., S.R. Phelps, S.W. Martin, E.L. Bartrand, and G.A. McMichael. 2007. Gene flow between resident and anadromous rainbow trout in the Yakima Basin: Ecological and genetic evidence. Pages 56-64 in R. K. Schroeder and J. D. Hall, editors. Redband trout: resilience and challenge in a changing landscape. Oregon Chapter, American Fisheries Society, Corvallis, Oregon.
- Pearsons, T.N. and G.M. Temple. 2007. Impacts of Early Stages of Salmon Supplementation and Reintroduction Programs on Three Trout Species. *North American Journal of Fisheries Management* 27:1-20.
- Pearsons, T.N., A.L. Fritts, and J.L. Scott. 2007. The effects of hatchery domestication on competitive dominance of juvenile spring Chinook salmon. *Canadian Journal of Fisheries and Aquatic Sciences* 64:803-812.
- Pearsons, T.N., D.D. Roley, and C.L. Johnson. 2007. Development of a carcass analog for nutrient restoration in streams. *Fisheries* 32:114-124.
- Pearsons, T.N. 2008. Misconception, reality, and uncertainty about ecological interactions and risks between hatchery and wild salmonids. *Fisheries* 33:278-290.
- Pearsons, T.N., C.L. Johnson, B.B. James, and G.M. Temple. 2009. Abundance and Distribution of Precociously Mature Male Spring Chinook Salmon of Hatchery and Natural Origin in the Yakima River. *North American Journal of Fisheries Management* 29:778-790.

- Pearsons, T.N. and G.M. Temple. 2010. Changes to Rainbow Trout Abundance and Salmonid Biomass in a Washington Watershed as Related to Hatchery Salmon Supplementation. *Transactions of the American Fisheries Society* 139:502-520.
- Pearsons, T.N. 2010. Operating Hatcheries within an Ecosystem Context Using the Adaptive Stocking Concept. *Fisheries* 35:23-31.
- Pacific Salmon Commission (PSC). 1994. Pacific Salmon Commission Joint Chinook Technical Committee 1993 annual report. Pacific Salmon Commission. Report Chinook (94)-1, 121 p. + app. (Available from Pacific Salmon Commission, 600-1155 Robson St., Vancouver, B.C. V6E 1B5.)
- RASP (Regional Assessment of Supplementation Planning). 1992. Supplementation in the Columbia River Basin, Parts 1-5. Report DOE/[BP 01830-11](#), Bonneville Power Administration.
- Rechisky, E.L., D.W. Welch, A.D. Porter, M.C. Jacobs, A. Ladouceur. 2009. Experimental measurement of hydrosystem-induced delayed mortality in juvenile Columbia River spring Chinook salmon using a large-scale acoustic array. *Canadian Journal of Fisheries and Aquatic Sciences* 66: 1019-1024.
- Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. *Bulletin of the Fisheries Research Board of Canada* 191:382.
- Salafsky, N., R. Margoluis, and K. Redford. 2001. Adaptive management: A tool for conservation practitioners. Washington, D.C. Biodiversity Support Program. Available at: <http://www.fosonline.org/wordpress/wp-content/uploads/2010/06/AdaptiveManagementTool.pdf>
- Sandford, B.P. and S.G. Smith. 2002. Estimation of smolt-to-adult return percentages for Snake River Basin anadromous salmonids, 1990-1997. *Journal of Agricultural, Biological, and Environmental Statistics* 7:243-263.
- Schroder, S.L., C.M. Knudsen, T.N. Pearsons, T.W. Kassler, S.F. Young, C.A. Busack, and D.E. Fast. 2008. Breeding Success of Wild and First-Generation Hatchery Female Spring Chinook Salmon Spawning in an Artificial Stream. *Transactions of the American Fisheries Society*, 137:1475-1489.
- Schroder, S.L., C.M. Knudsen, T.N. Pearsons, T.W. Kassler, S.F. Young, E.P. Beall, and D.E. Fast. 2010. Behavior and Breeding Success of Wild and First-Generation Hatchery Male Spring Chinook Salmon Spawning in an Artificial Stream. *Transactions of the American Fisheries Society*, 139:989-1003.
- Schroder, S.L., C.M. Knudsen, T.N. Pearsons, T.W. Kassler, E.P. Beall, S.F. Young, and D.E. Fast. 2012. Breeding Success of four male life history types of spring Chinook Salmon spawning in an artificial stream. *Environmental Biology of Fishes*, 94:231-248.

- Smith, E.P., D.R. Orvos, and J. Cairns, Jr. 1993. Impact assessment using the before-after-control-impact (BACI) model: concerns and comments. *Canadian Journal of Fisheries and Aquatic Sciences* 50:627-637.
- Stewart-Oaten, A., W.W. Murdoch, and K.R. Parker. 1986. Environmental impact assessment: pseudoreplication in time? *Ecology*: 67:929-940.
- TAC (*United States versus Oregon* Technical Advisory Committee). 1997. 1996 All Species Review, Columbia River Fish Management Plan. August 4, 1997. Columbia River Inter-Tribal Fish Commission, Portland, Oregon.
- Temple, G.M. and T. N. Pearsons. 2006. Evaluation of the recovery period in mark-recapture population estimates of rainbow trout in small streams. *North American Journal of Fisheries Management* 26:941-948.
- Temple, G.M., and T.N. Pearsons. 2007. Electrofishing: Backpack and Driftboat. Pages 95-132 in D. L. Johnson and 6 editors. *Salmonid Field Protocol Handbook*. American Fisheries Society, Bethesda, Maryland. (Protocols Handbook Chapter 3).
- Temple, G.M., T. Newsome, T.D. Webster, and S.W. Coil. 2012. Interactions between rainbow trout and reintroduced coho salmon in Taneum Creek, Washington. Chapter 2 in *Ecological interactions between non-target taxa of concern and hatchery supplemented salmon*, Annual Report to BPA. Available at: <https://pisces.bpa.gov/release/documents/documentviewer.aspx?doc=P128686>
- Temple, G.M., T. Newsome, T.D. Webster, and S.W. Coil. 2017. Evaluation of Rainbow Trout Abundance, Biomass, and Condition Following Coho Salmon Reintroduction in Taneum Creek, Washington. *Northwest Science* 91:54-68.
- Tiffan, K.F., P.G. Wagner, K.S. Wolf, and P.A. Hoffarth. 2009. Application of the SHOALS survey system to fisheries investigations in the Columbia River. In J.M. Bayer and J.L. Schei, (eds.). *PNAMP Special Publication: Remote Sensing Applications for Aquatic Resource Monitoring*, Pacific Northwest Aquatic Monitoring Partnership, Cook, WA. Chapter 5, p. 35-42.
- USACE (United States Army Corps of Engineers). 2014. Double-crested Cormorant Management Plan to Reduce Predation of Juvenile Salmonids in the Columbia river Estuary, Draft Environmental Impact Statement. United States Army Corps of Engineers, Portland, Oregon. Available online at: http://www.nwp.usace.army.mil/Portals/24/docs/announcements/EIS/DRAFT_Double-Crested_Cormorant_Plan_Reduce_Predation_Columbia_River_Estuary_EIS.pdf
- Venditti, D.A., R.N. Kinzer, K.A. Apperson, B. Barnett, M. Belnap, T. Copeland, M.P. Corsi, and K. Tardy. 2017. Effects of hatchery supplementation on abundance and productivity of natural-origin Chinook salmon: two decades of evaluation and implications for conservation programs. *Canadian Journal of Fisheries and Aquatic*

Sciences, <https://doi.org/10.1139/cjfas-2016-0344>. See also: <https://nwcouncil.app.box.com/s/gsolcxk9nv1w3897am4th5nl7xe6108g>.

- Waters, C.D., J.J. Hard, M.S.O. Briec, D.E. Fast, K.I. Warheit, R. Waples, C.M. Knudsen, W.J. Bosch, and K.A. Naish. 2015. Effectiveness of managed gene flow in reducing genetic divergence associated with captive breeding. *Evolutionary Applications* 8:956-971. DOI: [10.1111/eva.12331](https://doi.org/10.1111/eva.12331).
- Welch, D.W., E.L. Rechisky, M.C. Melnychuk, A.D. Porter, C.J. Walters, S. Clements, B.J. Clemens, R.S. McKinley, C. Schreck. 2008. Survival of migrating salmon smolts in large rivers with and without dams. *PLoS Biology* Vol. 6, Issue 10, p e265, doi:10.1371/journal.pbio.0060265.
- Wise, D.R., M.L. Zuroske, K.D. Carpenter, and R.L. Kiesling. 2009. Assessment of Eutrophication in the Lower Yakima River Basin, Washington, 2004-07: U.S. Geological Survey Scientific Investigations Report [2009-5078](https://pubs.usgs.gov/ofr/2009/5078/), 108 p.
- Wright, S. G. 1970. Size, age, and maturity of coho salmon in Washington's ocean troll fishery. *Wash. Dep. Fish., Fish. Res. Papers* 3(2):63-71.
- Yakama Nation. 2019. Revised Master Plan for Yakima Subbasin Summer-and Fall-Run Chinook, Coho Salmon and Steelhead. Confederated Tribes and Bands of the Yakama Nation. Toppenish, WA. September 2019.
- Yakama Subbasin Fish and Wildlife Planning Board (YSFWPB). 2004. Final Draft Yakima Subbasin Plan, May 28, 2004 and Management Plan Supplement Yakima Subbasin Plan, November 26, 2004. Yakima, Wa. Available at: <http://www.ybfwrb.org/recovery-planning/subbasin-plan/>

APPENDICES

- A. Use of Data and Products
- B. Summary of Data Collected by the Yakama Nation relative to Yakima River Spring Chinook Salmon and the Cle Elum Spring Chinook Supplementation and Research Facility
- C. 2020 Annual Chandler Certification for Out-migrating Spring (Yearling) Chinook Smolts
- D. Survival to McNary Dam for PIT-tagged Spring Chinook Salmon smolts released at Roza Dam from 1999--2020
- E. Juvenile Coho outmigration survival and adult Coho returns to the Yakima Basin, 1999-2020
- F. Juvenile Outmigration Survival of Yakima Basin Summer Chinook Smolts to Prosser and McNary Dams, 2009-2020

Appendix A: Use of Data & Products

All data and findings should be considered preliminary until results are published in the peer-reviewed literature.

Where will you post or publish the data your project generates?

[Fish Passage Center](#)

[Yakama Nation Fisheries website](#)

[RMIS - Regional Mark Information System](#)

[Columbia River DART](#)

[StreamNet Database](#)

[cbfish.org](#) (see projects [1995-063-25](#) and [1988-120-25](#))

[PTAGIS Website](#)

[Washington State SaSI](#)

A system has been developed that serves Yakima Basin adult abundance and trap sampling (requires login) data for the Prosser and Roza data sets. This system can be accessed at: <https://www.yakamafish-nsn.gov/fish-data>.

Describe the accessibility of the data and what the requirements are to access them?

- Prosser and Roza dam daily count and trap sample (requires login) data <https://www.yakamafish-nsn.gov/fish-data>.
- Integration of PIT and CWT release and recovery data with [PTAGIS](#), [RMIS](#), and [Fish Passage Center](#) databases (available to the public)
- BPA quarterly and annual reports (e.g., PISCES, available to the public via [CBfish.org](#))
- NPCC project proposals (available to the public via [nwcouncil.org](#))
- Yakima Basin [conference presentations](#) and [project technical reports](#) (available to the public)
- Yakima Basin [Status and Trends Annual Reports](#) (available to the public)

Additional data is available in the main body and other appendices of this report and by email contact through the data managers (Yakima Basin, contact Bill Bosch, bill_bosch@yakama.com; Klickitat Basin, contact Michael Babcock, mbabcock@ykfp.org). Project data managers continue to participate in the Coordinated Assessments process to develop pilot exchange templates for adult and juvenile abundance and productivity parameters. However, we continue to believe that the best way to prioritize our data management work load is to develop databases to store the status and trend data we have been collecting over many years as well as the web tools necessary to access these data in downloadable format. The system we have developed to share Prosser and Roza dam daily count and trap sample data is an example of the progress we are making towards this end.

Appendix B

Summary of Data Collected by the Yakama Nation
relative to
Yakima River Spring Chinook Salmon and the
Cle Elum Spring Chinook Supplementation and Research Facility

2020 Annual Report

July 16, 2021

Prepared by:

Bill Bosch
Yakima/Klickitat Fisheries Project
Yakama Nation Fisheries
760 Pence Road
Yakima, WA 98908

Prepared for:

Bonneville Power Administration
P.O. Box 3621
Portland, OR 97208
Project Numbers: 1995-063-25
Contract Numbers: 56662 REL 214

Acknowledgments

Monitoring and evaluation efforts for the Cle Elum Supplementation and Research Facility (CESRF) and Yakima River spring Chinook salmon are the result of a cooperative effort by many individuals from a variety of agencies including the Yakama Nation Fisheries Program (YN), the Washington Department of Fish and Wildlife (WDFW), the United States Fish and Wildlife Service (USFWS), the National Oceanic and Atmospheric Administration Fisheries department (NOAA Fisheries) as well as some consultants and contractors.

The core project team includes the following individuals: Mark Johnston, Bill Bosch, Shubha Pandit, Andrew Matala, Daylen Isaac, Chris Frederiksen, Michael Porter, Joe Hoptowit, and a number of technicians from the YN; Charles Strom and a number of assistants from the CESRF; Anthony Fritts, Gabe Temple, Christopher Johnson, and a number of assistants from the WDFW; the USFWS for fish health related analyses; and Don Larsen, Andy Dittman, and assistants from NOAA Fisheries. The technicians and assistants are too numerous and varied to mention each by name (and risk leaving some out). However, their hard work in the field is the source of much of the raw data needed to complete this report. We sincerely appreciate their hard work and dedication to this project.

We would especially like to thank former members of the Yakima/Klickitat Fisheries Project, Dave Fast, David Lind, Paul Huffman, Bruce Watson, Joel Hubble, Bill Hopley, Todd Pearsons, Steve Schroder, Curt Knudsen, Doug Neeley and Craig Busack. These individuals put in countless hours of hard work during the planning, design, and implementation of this project. Their contributions helped to lay a solid foundation for this project and our monitoring and evaluation efforts. Dan Barrett (retired) served as the manager of the CESRF from 1997-2002. He helped to lay a solid foundation for the critical work done day in and day out at the Cle Elum facility.

We also need to recognize and thank the Yakama Nation and WDFW for their continued support, and the Columbia River Inter-Tribal Fish Commission, the University of Idaho, the Pacific States Marine Fisheries Commission, Moberg, Jones, and Stokes, and Central Washington University for their many contributions to this project including both recommendations and data services.

This work is funded by the Bonneville Power Administration (BPA) through the Northwest Power and Conservation Council's (NPCC) Fish and Wildlife Program. Michelle O'Malley is BPA's contracting officer and technical representative (COTR) for this project. David Byrnes and Patricia Smith preceded Michelle in this position and contributed substantially to the project over the years.

Abstract

Historically, the return of spring Chinook salmon (*Oncorhynchus tshawytscha*) to the Yakima River numbered about 200,000 fish annually (BPA, 1990). Spring Chinook returns to the Yakima River averaged fewer than 3,500 fish per year through most of the 1980s and 1990s (less than 2% of the historical run size).

In an attempt to reverse this trend the Northwest Power and Conservation Council (formerly the Northwest Power Planning Council, NPPC) in 1982 first encouraged Bonneville Power Administration (BPA) to “fund the design, construction, operation, and maintenance of a hatchery to enhance the fishery for the Yakima Indian Nation as well as all other harvesters” (NPPC 1982). After years of planning and design, an Environmental Impact Statement (EIS) was completed in 1996 and the CESRF was authorized under the NPCC’s Fish and Wildlife Program with the stated purpose being “to test the assumption that new artificial production can be used to increase harvest and natural production while maintaining the long-term genetic fitness of the fish population being supplemented and keeping adverse genetic and ecological interactions with non-target species or stocks within acceptable limits”. The CESRF became operational in 1997. This project is co-managed by the Yakama Nation and the Washington Department of Fish and Wildlife (WDFW) with the Yakama Nation as the lead entity.

This report documents data collected from Yakama Nation tasks related to monitoring and evaluation of the CESRF and its effect on natural populations of spring Chinook in the Yakima Basin through 2020. This report is not intended to be a scientific evaluation of spring Chinook supplementation efforts in the Yakima Basin. Rather, it is a summary of methods and data (additional information about methods used to collect these data may be found in the main section of this annual report) relating to Yakima River spring Chinook collected by Yakama Nation biologists and technicians from 1982 (when the Yakama Nation fisheries program was implemented) to present. Data summarized in this report include:

- Adult-to-adult returns
- Annual run size and escapement
- Adult traits (e.g., age composition, size-at-age, sex ratios, migration timing, etc.)
- CESRF reproductive statistics (including fecundity and fish health profiles)
- CESRF juvenile survival (egg-to-fry, fry-to-smolt, smolt-to-smolt, and smolt-to-adult)
- CESRF juvenile traits (e.g., length-weight relationships, migration timing, etc.)
- Harvest impacts

The data presented here are, for the most part, “raw” data and should not be used without paying attention to caveats associated with these data and/or consultation with project biologists. No attempt is made to explain the significance of these data in this report as this is left to more comprehensive reports and publications produced by the project. Data in this report should be considered preliminary until published in the peer reviewed literature.

Table of Contents

Abstract	i
List of Tables	iii
List of Figures	v
List of Appendices	v
Introduction	1
Program Objectives	1
Facility Descriptions	1
Yakima River Basin Overview	2
Adult Salmon Evaluation	3
Broodstock Collection and Representation	3
Natural- and Hatchery-Origin Escapement	5
Adult-to-adult Returns	7
Age Composition	14
Sex Composition	20
Size at Age	25
Migration Timing	33
Spawning Timing	35
Redd Counts and Distribution	36
Homing	37
Straying	38
CESRF Spawning and Survival	39
Female BKD Profiles	42
Fecundity	43
Juvenile Salmon Evaluation	44
Food Conversion Efficiency	44
Length and Weight Growth Profiles	45
Juvenile Fish Health Profile	46
Incidence of Precocialism	46
Smolt Outmigration Timing	49
Smolt-to-Smolt Survival	50
Smolt-to-Adult Survival	52
Harvest Monitoring	61
Yakima Basin Fisheries	61
Marine Fisheries	64
Literature Cited	66

List of Tables

Table 1. Counts of wild/natural spring Chinook (including jacks), brood collection, and brood representation of wild/natural run at Roza Dam, 1997 – present.	4
Table 2. Escapement (Roza Dam counts less brood stock collection and harvest above Roza) of natural- (NoR) and hatchery-origin (HoR) spring Chinook to the upper Yakima subbasin, 1982 – present.	6
Table 3. Yakima River spring Chinook run (CESRF and wild, adults and jacks combined) reconstruction, 1991-present.....	8
Table 4. Adult-to-adult productivity indices for upper Yakima wild/natural stock.	9
Table 5. Adult-to-adult productivity indices for Naches River wild/natural stock.....	10
Table 6. Adult-to-adult productivity indices for American River wild/natural stock.....	11
Table 7. Adult-to-adult productivity indices for Naches/American aggregate (wild/natural) population.	12
Table 8. Adult-to-adult productivity for Cle Elum SRF spring Chinook.	13
Table 9. Percentage by sex and age of American River wild/natural spring Chinook carcasses sampled on the spawning grounds and sample size (n), 1986-present.	15
Table 10. Percentage by sex and age of Naches River wild/natural spring Chinook carcasses sampled on the spawning grounds and sample size (n), 1986-present.	16
Table 11. Percentage by sex and age of upper Yakima River wild/natural spring Chinook carcasses sampled on the spawning grounds and sample size (n), 1986-present.	17
Table 12. Percentage by sex and age of upper Yakima River CESRF spring Chinook carcasses sampled on the spawning grounds and sample size (n), 2001-present.	18
Table 13. Percentage by sex and age of upper Yakima River wild/natural spring Chinook collected for brood stock at Roza Dam and sample size (n), 1997-present.....	19
Table 14. Percentage by sex and age of upper Yakima River CESRF spring Chinook collected for research or brood stock at Roza Dam and sample size (n), 2001-present.	19
Table 15. Percent of American River wild/natural spring Chinook carcasses sampled on the spawning grounds by age and sex, 1986-present.....	21
Table 16. Percent of Naches River wild/natural spring Chinook carcasses sampled on the spawning grounds by age and sex, 1986-present.....	22
Table 17. Percent of Upper Yakima River wild/natural spring Chinook carcasses sampled on the spawning grounds by age and sex, 1986-present.....	23
Table 18. Percent of upper Yakima River CESRF spring Chinook carcasses sampled on the spawning grounds by age and sex, 2001-present.....	24
Table 19. Percent of upper Yakima River wild/natural spring Chinook collected for brood stock at Roza Dam by age and sex, 1997-present.	24
Table 20. Percent of Upper Yakima River CESRF spring Chinook collected for research or brood stock at Roza Dam by age and sex, 2001-present.	25
Table 21. Counts and mean mid-eye (MEHP) or post-orbital (POHP) to hypural plate lengths (cm) of American River wild/natural spring Chinook from carcasses sampled on the spawning grounds by sex and age, 1987-present.....	26
Table 22. Counts and mean mid-eye (MEHP) or post-orbital (POHP) to hypural plate lengths (cm) of Naches River wild/natural spring Chinook from carcasses sampled on the spawning grounds by sex and age, 1987-present.	27

Table 23. Counts and mean mid-eye (MEHP) or post-orbital (POHP) to hypural plate lengths (cm) of upper Yakima River wild / natural spring Chinook from carcasses sampled on the spawning grounds by sex and age, 1986-present.....	28
Table 24. Counts and mean post-orbital to hypural plate (POHP) lengths (cm) of upper Yakima River CESRF spring Chinook from carcasses sampled on the spawning grounds by sex and age, 2001-present.	29
Table 25. Counts and mean post-orbital to hypural plate (POHP) lengths (cm) of upper Yakima River wild/natural spring Chinook from carcasses sampled at the CESRF prior to spawning by sex and age, 1997-present.....	29
Table 26. Counts and mean post-orbital to hypural plate (POHP) lengths (cm) of upper Yakima River CESRF spring Chinook from carcasses sampled at the CESRF prior to spawning by sex and age, 2001-present.....	30
¹ Few length samples were collected since these fish were not spawned in 2006. Table 27. Counts and mean post-orbital to hypural plate (POHP) lengths (cm) of upper Yakima River wild/natural spring Chinook from fish sampled at Roza Dam by sex ¹ and age, 1997-present.	30
¹ Sex determined by visual observation prior to 2010 and by ultrasound from 2010 to present. Table 28. Counts and mean post-orbital to hypural plate (POHP) lengths (cm) of upper Yakima River CESRF spring Chinook from fish sampled at Roza Dam by sex ¹ and age, 2001-present.	31
Table 29. Comparison of 5%, median (50%), and 95% passage dates of wild/natural and CESRF adult spring Chinook (including jacks) at Roza Dam, 1997-Present.....	34
Table 30. Median spawn ¹ dates for spring Chinook in the Yakima Basin.	35
Table 31. Yakima Basin spring Chinook redd count summary, 1981 – present.	36
Table 32. Estimated number of PIT- and CWT-tagged CESRF fish not returning to the Yakima River Basin (strays), and marked fish sampled during spawner surveys in the Naches Basin, per number of returning fish, brood years 1997-present.....	38
Table 33. Cle Elum Supplementation and Research Facility spawning and survival statistics (NoR brood only), 1997 - present.	40
Table 34. Cle Elum Supplementation and Research Facility spawning and survival statistics (HoR brood only), 2002 - present.	41
Table 35. Mean fecundity by age of adult females (BKD rank < 6) spawned at CESRF, 1997-present.....	43
Table 36. Mean food conversion (lbs fed/lbs gained) of CESRF juveniles by brood year and growth month, 1997 – present.....	44
Table 37. CESRF total releases by brood year, treatment, and acclimation site.	48
Table 38. CESRF average pond densities at release by brood year, treatment, and acclimation site.	49
Table 39. Estimated smolt passage at Chandler and smolt-to-adult return indices (Chandler smolt to Yakima R. mouth adult) for Yakima Basin wild/natural and CESRF-origin spring Chinook.....	54
Table 40. Estimated wild/natural smolt-to-adult return rates (SAR) based on adult detections of PIT tagged fish. Roza tagged smolts to Bonneville Dam adult returns. Footnotes follow Table 42.	55
Table 41. Estimated CESRF smolt-to-adult return rates (SAR) based on adult detections of PIT tagged fish. Roza tagged smolts to Bonneville Dam adult returns.....	56

Table 42. Overall McNary Dam smolt to Bonneville Dam adult return rates (SAR) based on juvenile and adult detections of wild/natural Yakima R. spring Chinook PIT-tagged and released at Roza Dam (Table B.74 in McCann et al. 2020).	57
Table 43. Overall McNary Dam smolt to Bonneville Dam adult return rates (SAR) based on juvenile and adult detections of CESRF PIT-tagged spring Chinook (Table B.80 in McCann et al. 2020).	58
Table 44. Estimated release-to-adult survival of PIT-tagged CESRF fish (CESRF tagged smolts to Bonneville and Roza Dam adult returns).	59
Table 45. Estimated release-to-adult survival of non-PIT-tagged CESRF fish (CESRF tagged smolts to Roza Dam adult returns).	60
Table 46. Spring Chinook harvest in the Yakima River Basin, 1985-present.	62
Table 47. Estimated run size, harvest, and harvest rates of Yakima Basin spring Chinook in Columbia River mainstem and terminal area fisheries, 1986-present.	63
Table 48. Marine and freshwater recoveries of CWTs from brood year 1997-2015 releases of spring Chinook from the CESRF as reported to the Regional Mark Information System (RMIS) Jan. 5, 2021.	64

List of Figures

Figure 1. Yakima River Basin.	2
Figure 2. Mean spring Chinook run timing and broodstock collection at Roza Dam, 2011-2020.	3
Figure 3. Proportionate passage timing at Roza Dam of wild/natural and CESRF adult spring Chinook (including jacks), 2011-2020.	33
Figure 4. Proportion of wild/natural females spawned at CESRF by BKD rank, 1997 – present.	42
Figure 5. Mean fork length (cm) of CESRF juveniles by brood year and growth month, 1997 - present.	45
Figure 6. Mean Weight (fish/lb) of CESRF juveniles by brood year and growth month, 1997 - present.	45
Figure 7. ELISA-risk profile of CESRF juveniles by brood year, 1997 – present (data source: USFWS).	46
Figure 8. Mean flow approaching Prosser Dam versus mean estimated smolt passage at Prosser of aggregate wild/natural and CESRF spring Chinook for outmigration years 1999-2020.	50
Figure 9. Marine recovery locations of coded-wire-tagged CESRF spring Chinook, recovery years 2008-2012.	65

List of Appendices

<i>Appendix A . Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.</i>	67
----------------------------------------------------------------------------------------------------	-----------

Introduction

Program Objectives

The CESRF was authorized in 1996 under the NPCC's Fish and Wildlife Program with the stated purpose being "to test the assumption that new artificial production can be used to increase harvest and natural production while maintaining the long-term genetic fitness of the fish population being supplemented and keeping adverse genetic and ecological interactions with non-target species or stocks within acceptable limits". The CESRF became operational in 1997. The experimental design calls for a total release of 810,000 smolts annually from each of three acclimation sites associated with the facility (see facility descriptions). To minimize risk of over-collecting brood stock and to maintain lower pond rearing densities, the YKFP policy group took action in 2011 to create a release target range of 720,000-810,000 smolts for brood collection purposes. Female percentage, fecundity and survival rates are expected to result in releases between 720,000 and 810,000 smolts in most years. The first program cycle (brood years 1997 through 2001) also included testing new Semi-Natural rearing Treatments (SNT) against the Optimum Conventional Treatments (OCT) of existing successful hatcheries in the Pacific Northwest. The second program cycle (brood years 2002-2004) tested whether a slower, more natural growth regime could be used to reduce the incidence of precocialism that may occur in hatchery releases without adversely impacting overall survival to adult returns. Brood years 2005-2007 tested survival using different types of feed treatment. Subsequent broods have used a standard treatment in all raceways. With guidance and input from the NPCC and the Independent Scientific Review Panel (ISRP) in 2001, the Naches subbasin population of spring Chinook was established as a wild/natural control. A hatchery control line at the CESRF was also established with the first brood production for this line collected in 2002. Please refer to the project's "[Supplementation Monitoring Plan](#)" (Chapter 7 in 2005 annual report on project genetic studies) for additional information regarding these control lines.

Facility Descriptions

Returning adult spring Chinook are monitored at the Roza adult trapping facility located on the Yakima River (Rkm 205.8). This facility provides the means to monitor every fish returning to the upper Yakima Basin and to collect adults for the CESRF program. All returning CESRF fish (adipose-clipped fish) are sampled for biological characteristics and marks and returned to the river with the exception of fish collected for broodstock, experimental sampling, and all hatchery control line fish. Through 2006, all wild/natural fish passing through the Roza trap were returned directly to the river with the exception of fish collected for broodstock or fish with metal tag detections which were sampled for marks and biological characteristics. Beginning in 2007, all wild/natural fish were sampled (as described above) and tissue samples were collected for a "Whole Population" Pedigree Study of Upper Yakima Spring Chinook (see related project [2009-009-00](#)).

The CESRF is located on the Yakima River just south of the town of Cle Elum (rkm 295.5). It is used for adult broodstock holding and spawning, and early life incubation and rearing. Fish are spawned in September and October of a given brood year (BY). Fish are typically ponded in March or April of BY+1. The juveniles are reared at Cle Elum, marked in October through

December of BY+1, and moved to one of three acclimation sites for final rearing in January to February of BY+2. Acclimation sites are located at Easton (ESJ, rkm 317.8), Clark Flats near the town of Thorp (CFJ, rkm 266.6), and Jack Creek (JCJ, approximately 32.5 km north of Cle Elum) on the North Fork Teanaway River (rkm 10.2). Fish are volitionally released from the acclimation sites beginning on March 15 of BY+2, with any remaining fish “flushed out” of the acclimation sites by May 15 of BY+2. The annual production goal for the CESRF program is 720,000 to 810,000 fish for release as yearlings at 30 g/fish or 15 fish per pound (fpp) although size-at-release may vary depending on experimental protocols (see Program Objectives).

Yakima River Basin Overview

The Yakima River Basin is located in south central Washington. From its headwaters near the crest of the Cascade Range, the Yakima River flows 344 km (214 miles) southeastward to its confluence with the Columbia River (Rkm 539.5; Figure 1).




Figure 1. Yakima River Basin.

Three genetically distinguishable populations of spring Chinook salmon exist in the Yakima basin: the American River, the Naches, and the Upper Yakima Stocks (Figure 1). The upper Yakima was selected as the population best suited for supplementation and associated evaluation and research efforts.

Local habitat problems related to irrigation, logging, road building, recreation, agriculture, and livestock grazing have limited the production potential of spring Chinook in the Yakima River basin. It is hoped that recent initiatives to improve habitat within the Yakima Basin, such as those being funded through the NPCC's fish and wildlife program, the Pacific Coastal Salmon Recovery Fund, and the Washington State salmon recovery fund, will: 1) restore and maintain natural stream stability; 2) reduce water temperatures; 3) reduce upland erosion and sediment delivery rates; 4) improve and re-establish riparian vegetation; and 5) re-connect critical habitats throughout the basin. These habitat restoration efforts should permit increased utilization of habitat by spring Chinook salmon in the Yakima basin thereby increasing fish survival and productivity.

Adult Salmon Evaluation

Broodstock Collection and Representation

One of the program's goals is to collect broodstock from a representative portion of the population throughout the run. If the total run size could be known in advance, collecting brood stock on a daily basis in exact proportion to total brood need as a proportion of total run size would result in ideal run representation. Since it is not possible to know the run size in advance, the CESRF program uses a brood collection schedule that is based on average run timing once the first fish arrive at Roza Dam. We have found that, while river conditions dictate run timing (i.e., fish may arrive earlier or later depending on flow and temperature), once fish begin to move at Roza, the pattern in terms of relative run strength over time is very similar from year to year. Thus a brood collection schedule matching normal run timing patterns was developed to assure that fish are collected from all portions of the run (Figure 2).




Figure 2. Mean spring Chinook run timing and broodstock collection at Roza Dam, 2011-2020.

Another program goal is to take no more than 50% of the wild/natural adult return to Roza Dam for broodstock. Given this goal and with a set brood collection schedule at Roza Dam, the project imposed a rule that no more than 50% of the fish arriving on any given day be taken for broodstock. Under-collection relative to the schedule is “carried over” to subsequent days and weeks. This allows brood collection to adjust relative to actual run timing and run strength. Performance across years with respect to these brood collection goals is given in Table 1. Since 2015, the spring Chinook return has been impeded by thermal barriers in the lower Yakima River as warmer air temperatures combined with reduced summer and fall flows have increased water temperatures. Mean daily water temperatures near Prosser (rkm 76 from the mouth of the Yakima R.) have exceeded 68° F on several days between June and September during these years (source [U.S. BOR hydromet](#) database). This may have caused a large number of fish to stray or be delayed in their migration above Roza Dam.

Table 1. Counts of wild/natural spring Chinook (including jacks), brood collection, and brood representation of wild/natural run at Roza Dam, 1997 – present.

Year	Trap Count	Brood Take	Brood %	Portion of run collected: ¹			Portion of collection from: ²		
				Early ³	Middle ³	Late ³	Early ³	Middle ³	Late ³
1997	1,445	261	18.1%	26.4%	17.6%	17.7%	7.3%	83.1%	9.6%
1998	795	408	51.3%	51.1%	51.3%	51.9%	5.6%	84.3%	10.0%
1999	1,704	738	43.3%	44.6%	44.1%	35.9%	5.6%	86.3%	8.1%
2000	11,639	567	4.9%	10.7%	4.5%	4.4%	12.5%	77.8%	9.7%
2001	5,346	595	11.1%	6.9%	11.4%	10.7%	3.0%	87.7%	9.2%
2002	2,538	629	24.8%	15.7%	25.2%	26.1%	3.2%	86.3%	10.5%
2003	1,558	441	28.3%	52.5%	25.9%	36.4%	9.5%	77.8%	12.7%
2004	7,804	597	7.6%	2.6%	7.4%	12.8%	2.0%	81.6%	16.4%
2005	5,086	510	10.0%	2.2%	9.5%	21.9%	1.3%	77.0%	21.7%
2006	2,050	419	20.4%	48.5%	22.2%	41.0%	9.1%	75.1%	15.8%
2007	1,293	449	34.7%	25.0%	34.4%	60.6%	3.2%	80.0%	16.9%
2008	1,677	457	27.3%	57.7%	26.7%	32.4%	9.3%	79.0%	11.6%
2009	3,030	486	16.0%	10.0%	14.1%	35.9%	3.5%	73.9%	22.6%
2010	3,185	336	10.5%	6.4%	15.0%	22.5%	2.0%	82.6%	15.3%
2011	4,395	377	8.6%	11.3%	9.2%	21.3%	5.6%	73.2%	21.2%
2012	2,924	374	12.8%	1.9%	12.3%	27.4%	1.1%	79.9%	19.0%
2013	2,784	398	14.3%	18.5%	13.0%	22.0%	9.5%	75.1%	15.3%
2014	4,168	384	9.2%	4.8%	8.6%	16.9%	2.3%	80.5%	17.1%
2015	3,962	442	11.2%	3.1%	8.2%	40.6%	2.0%	59.9%	38.1%
2016	2,712	376	13.9%	5.3%	14.8%	18.6%	2.5%	84.7%	12.9%
2017	1,711	382	22.3%	53.6%	19.0%	45.4%	11.4%	69.9%	18.7%
2018	827	294	35.6%	3.0%	33.7%	87.6%	0.3%	75.1%	24.6%
2019	703	312	44.4%	48.1%	46.3%	29.1%	8.3%	84.3%	7.3%
2020	958	427	44.6%	47.7%	48.1%	15.9%	4.9%	91.1%	4.0%

1. This is the proportion of the earliest, middle, and latest running components of the entire wild/natural run which were taken for broodstock. Ideally, this collection percentage would be equal throughout the run and would match the “Brood %”.
2. This is the proportion of the total broodstock collection taken from the earliest, middle, and latest components of the entire wild/natural run. Ideally, these proportions would match the definitions for early, middle, and late given in 3.
3. Early is defined as the first 5% of the run, middle is defined as the middle 85%, and late as the final 10% of the run.

Natural- and Hatchery-Origin Escapement

While the project does not actively manage for a specific spawning escapement proportion (natural- to hatchery-origin adults), we are monitoring the proportion of natural influence (PNI; Table 2). The project will adaptively manage this parameter considering factors such as: policy input regarding surplusing of fish, meeting overall production goals of the project, guidance from the literature relative to percentage of hatchery fish on the spawning grounds with fitness loss, considerations about what risk is acceptable in a project designed to evaluate impacts from that risk, and the numerous risk containment measures already in place in the project. The State of Washington is using mark-selective fisheries in the lower Columbia River and, when possible, in the lower Yakima River in part as a tool to manage escapement proportions. In 2011, the project initiated an effort to transfer some returning hatchery-origin CESRF adults from Roza Dam to Lake Cle Elum for the purpose of returning marine derived nutrients and salmon to the watersheds that feed the lake. This effort will also increase PNI in the major spawning areas of the Upper Yakima Basin. Natural- and hatchery-origin escapement to the upper Yakima Basin is given in Table 2. Wild/natural escapement to the Naches subbasin is given in Table 3.

Table 2. Escapement (Roza Dam counts less brood stock collection and harvest above Roza) of natural- (NoR) and hatchery-origin (HoR) spring Chinook to the upper Yakima subbasin, 1982 – present.

Year	Wild/Natural (NoR)			CESRF (HoR)			Total			pHOS ¹	PNI ¹
	Adults	Jacks	Total	Adults	Jacks	Total	Adults	Jacks	Total		
1982			1,146								
1983			1,007								
1984			1,535								
1985			2,331								
1986			3,251								
1987			1,734								
1988			1,340								
1989			2,331								
1990			2,016								
1991			1,583 ²								
1992			3,009								
1993			1,869								
1994			563								
1995			355								
1996			1,631								
1997	1,141	43	1,184								
1998	369	18	387								
1999	498	468	966								
2000	10,491	481	10,972		688	688	10,491	1,169	11,660	5.9%	
2001	4,454	297	4,751	6,065	982	7,047	10,519	1,279	11,798	59.7%	62.6%
2002	1,820	89	1,909	6,064	71	6,135	7,884	160	8,044	76.3%	56.7%
2003	394	723	1,117	1,036	1,105	2,141	1,430	1,828	3,258	65.7%	60.3%
2004	6,536	671	7,207	2,876	204	3,080	9,412	875	10,287	29.9%	77.0%
2005	4,401	175	4,576	627	482	1,109	5,028	657	5,685	19.5%	83.7%
2006	1,510	121	1,631	1,622	111	1,733	3,132	232	3,364	51.5%	66.0%
2007	683	161	844	734	731	1,465	1,417	892	2,309	63.4%	61.2%
2008	988	232	1,220	2,157	957	3,114	3,145	1,189	4,334	71.9%	58.2%
2009	1,843	701	2,544	2,234	2,260	4,494	4,077	2,961	7,038	63.9%	61.0%
2010	2,436	413	2,849	4,524	1,001	5,525	6,960	1,414	8,374	66.0%	60.2%
2011	3,092	926	4,018	3,162	1,404	4,566	6,254	2,330	8,584	53.2%	65.3%
2012	2,359	191	2,550	2,661	265	2,926	5,020	456	5,476	53.4%	65.2%
2013	1,708	678	2,386	1,587	840	2,427	3,295	1,518	4,813	50.4%	66.5%
2014	3,099	685	3,784	2,150	794	2,944	5,249	1,479	6,728	43.8%	69.6%
2015	3,357	163	3,520	1,779	167	1,946	5,136	330	5,466	35.6%	73.7%
2016	2,070	266	2,336	1,198	705	1,903	3,268	971	4,239	44.9%	69.0%
2017	1,135	194	1,329	1,328	660	1,988	2,463	854	3,317	59.9%	62.5%
2018	500	33	533	1,033	233	1,266	1,533	266	1,799	70.4%	58.7%
2019	316	81	397	828	266	1,094	1,144	347	1,491	73.4%	57.7%
2020	497	56	553	746	341	1,087	1,243	397	1,640	66.3%	60.1%
Mean ³	2,321	328	2,648	2,221	679	2,794	4,171	922	5,093	53.6%	64.8%

1. Proportion Natural Influence (including jacks) equals Proportion Natural-Origin Broodstock (pNOB; 1.0 as only NoR fish are used for supplementation line brood stock) divided by pNOB plus Proportion Hatchery-Origin Spawners (pHOS).
2. This is a rough estimate since Roza counts are not available for 1991.
3. For NoR columns, mean of 1997-present values. For all other columns, mean of 2001-present values.

Adult-to-adult Returns

The overall status of Yakima Basin spring Chinook is summarized in Table 3. Adult-to-adult return and productivity data for the various populations are given in Tables 4-8 (Means are for 1988 to present).

Table 3. Yakima River spring Chinook run (CESRF and wild, adults and jacks combined) reconstruction, 1991-present.

Year	River Mouth Run Size ¹			Harvest	Prosser	Harvest	Spawners	Roza	Roza	Est. Escapement		Redd Counts	
	Adults	Jacks	Total	Below Prosser	Count	Above Prosser	Below Roza ²	Count	Removals ³	Upper Y.R. ⁴	Naches ⁵	Upper Y.R.	Naches
1991	2,802	104	2,906	27	2,879	5	131		40	1,583	1,121	582	460
1992	4,492	107	4,599	184	4,415	161	39	3,027	18	3,009	1,188	1,230	425
1993	3,800	119	3,919	44	3,875	85	56	1,869	0	1,869	1,865	637	554
1994	1,282	20	1,302	0	1,302	25	10	563	0	563	704	285	272
1995	526	140	666	0	666	79	9	355	0	355	223	114	104
1996	3,060	119	3,179	100	3,079	375	26	1,631	0	1,631	1,047	801	184
1997	3,092	81	3,173	0	3,173	575	20	1,445	261	1,184	1,133	413	339
1998	1,771	132	1,903	0	1,903	188	3	795	408	387	917	147	330
1999	1,513	1,268	2,781	8	2,773	596	55	1,704	738	966	418	212	186
2000	17,519	1,582	19,101	90	19,011	2,368	204	12,327	667	11,660	4,112	3,770	888
2001	21,225	2,040	23,265	1,793	21,472	2,838	286	12,516	718	11,798	5,829	3,226	1,192
2002	14,616	483	15,099	328	14,771	2,780	29	8,922	878	8,044	3,041	2,816	943
2003	4,868	2,089	6,957	59	6,898	381	83	3,842	584	3,258	2,592	868	935
2004	13,974	1,315	15,289	135	15,154	1,544	90	11,005	718	10,287	2,515	3,414	719
2005	8,059	699	8,758	34	8,724	440	28	6,352	667	5,685	1,904	2,009	574
2006	5,951	363	6,314	0	6,314	600	14	4,028	664	3,364	1,672	1,245	447
2007	2,968	1,335	4,303	10	4,293	269	13	3,025	716	2,309	986	722	313
2008	6,615	1,983	8,598	539	8,059	993	9	5,478	1,144	4,334	1,578	1,372	495
2009	7,441	4,679	12,120	1,517	10,603	836	18	8,633	1,595	7,038	1,117	1,575	482
2010	11,027	2,114	13,142	156	12,986	1,585	9	9,900	1,526	8,374	1,491	2,668	552
2011	13,398	4,561	17,960	909	17,051	3,471	0	10,520	1,936	8,584	3,060	1,898	580
2012	11,083	970	12,053	1,331	10,722	1,989	7	6,826	1,350	5,476	1,900	1,468	811
2013	7,101	3,144	10,245	1,191	9,054	1,462	171	6,053	1,240	4,813	1,369	648	376
2014	8,850	2,472	11,322	221	11,101	1,950	23	7,997	1,269	6,728	1,130	1,149	379
2015	8,795	556	9,351	83	9,268	732	0	6,433	967	5,466	2,103	1,321	614
2016	5,517	1,399	6,916	24	6,892	420	42	5,098	859	4,239	1,332	611	366
2017	5,462	1,701	7,163	122	7,041	1,150	25	4,193	876	3,317	1,673	539	293
2018	3,156	448	3,605	251	3,353	297	18	2,404	605	1,799	634	348	128
2019	1,756	466	2,222	0	2,222	40	17	2,007	516	1,491	158	235	31
2020	2,833	529	3,362	24	3,338	44	24	2,211	571	1,640	1,059	237	146
Mean ⁶	6,795	1,624	8,420	416	8,004	1,155	33	5,374	1,019	4,355	1,442	845	372

1. River Mouth run size is the greater of the Prosser count plus lower river harvest or estimated escapement plus all known harvest and removals.
2. Estimated as the average number of fish per redd in the upper Yakima times the number of redds between the Naches confluence and Roza Dam.
3. Roza removals include harvest above Roza, hatchery removals, and/or wild broodstock removals.
4. Estimated escapement into the upper Yakima River is the Roza count, less harvest or broodstock removals above Roza Dam except in 1991 when Upper Yakima River escapement is estimated as the (Prosser count - harvest above Prosser - Roza subtractions) times the proportion of redds counted in the upper Yakima.
5. Naches River escapement was estimated as the Prosser count, less harvest above Prosser and the Roza counts.
6. Recent 10-year average (2011-2020).

Estimated spawners for the Upper Yakima River are calculated as the estimated escapement to the Upper Yakima plus the estimated number of spawners in the Upper Yakima between the confluence with the Naches River and Roza Dam (Table 3). Total returns are based on the information compiled in Table 3. Age composition for Upper Yakima returns is estimated from spawning ground carcass scale samples for the years 1982-1996 (Table 11) and from Roza Dam brood stock collection samples for the years 1997 to present (Table 13). Since age-3 fish (jacks) are not collected for brood stock in proportion to the jack run size, the proportion of age-3 fish in the upper Yakima for 1997 to present is estimated using the proportion of jacks (based on visual observation) counted at Roza Dam relative to the total run size.

Table 4. Adult-to-adult productivity indices for upper Yakima wild/natural stock.

Brood Year	Estimated Spawners	Estimated Yakima R. Mouth Returns				Returns/ Spawner
		Age-3	Age-4	Age-5	Total	
1985	2,578	114	2,746	105	2,965	1.15
1986	3,960	171	2,574	149	2,893	0.73
1987	2,003	53	1,571	109	1,733	0.87
1988	1,400	53	3,138	132	3,323	2.37
1989	2,466	68	1,779	9	1,856	0.75
1990	2,298	79	566	0	645	0.28
1991	1,713	9	326	22	358	0.21
1992	3,048	87	1,861	95	2,043	0.67
1993	1,925	66	1,606	57	1,729	0.90
1994	573	60	737	92	890	1.55
1995	364	59	1,036	129	1,224	3.36
1996	1,657	1,059	12,882	630	14,571	8.79
1997	1,204	621	5,837	155	6,613	5.49
1998	390	434	2,803	145	3,381	8.68
1999	1,021 ¹	164	722	45	930	0.91
2000	11,864	856	7,689	127	8,672	0.73
2001	12,087	775	5,074	222	6,071	0.50
2002	8,073	224	1,875	148	2,247	0.28
2003	3,341	158	1,036	63	1,257	0.38
2004	10,377	207	1,547	75	1,828	0.18
2005	5,713	293	2,630	14	2,936	0.51
2006	3,378	868	2,887	133	3,888	1.15
2007	2,322	456	3,976	65	4,498	1.94
2008	4,343	1,135	3,410	123	4,668	1.07
2009	7,056	283	2,572	109	2,964	0.42
2010	8,383	923	3,854	59	4,836	0.58
2011	8,584	832	3,908	144	4,883	0.57
2012	5,483	197	2,445	20	2,662	0.49
2013	4,984	299	1,622	36	1,957	0.39
2014	6,751	241	814	12	1,067	0.16
2015	5,466	66	620	14 ²	701 ²	0.13 ²
2016	4,281	99	905 ²			
2017	3,342	75 ²				
2018	1,817					
2019	1,508					
2020	1,664 ²					
Mean	4,031	329	2,679	106	3,117	1.43

1. The geometric mean jack (age-3) proportion of spawning escapement from 1999-2020 was 0.17.

2. Preliminary.

Estimated spawners for the Naches/American aggregate population (Table 7) are calculated as the estimated escapement to the Naches Basin (Table 3). Estimated spawners for the individual Naches and American populations are calculated using the proportion of redds counted in the Naches Basin (excluding the American River) and the American River, respectively (see Table 31). Total returns are based on the information compiled in Table 3. Age composition for Naches Basin age-4 and age-5 returns are estimated from spawning ground carcass scale samples (see Tables 9-12). The proportion of age-3 fish is estimated after reviewing jack count (based on visual observations) data at Prosser and Roza dams. Since sample sizes for carcass surveys in the American and Naches Rivers can be very low in some years (Tables 9 and 10), it is recommended that the data in Tables 5 and 6 be used as indices only. Table 7 likely provides the most accurate view of overall productivity rates in the Naches River Subbasin.

Table 5. Adult-to-adult productivity indices for Naches River wild/natural stock.

Brood Year	Estimated Spawners	Estimated Yakima R. Mouth Returns					Returns/ Spawner
		Age-3	Age-4	Age-5	Age-6	Total	
1985	683	132	574	396	0	1,102	1.61
1986	2,666	68	712	499	15	1,294	0.49
1987	1,162	27	183	197	0	407	0.35
1988	1,340	32	682	828	0	1,542	1.15
1989	992	28	331	306	0	665	0.67
1990	954	24	170	74	0	269	0.28
1991	706	7	37	121	57	222	0.31
1992	852	29	877	285	0	1,191	1.40
1993	1,145	45	593	372	0	1,010	0.88
1994	474	14	164	164	0	343	0.72
1995	124	40	164	251	0	455	3.66
1996	887	179	3,983	1,620	0	5,782	6.52
1997	762	207	3,081	708	0	3,996	5.24
1998	503	245	1,460	1,128	0	2,833	5.63
1999	358 ¹	113	322	190	0	626	1.75
2000	3,862	71	2,060	215	0	2,346	0.61
2001	3,912	126	1,254	471	0	1,850	0.47
2002	1,861	59	753	153	0	965	0.52
2003	1,400	52	237	175	0	464	0.33
2004	2,197	107	875	218	0	1,199	0.55
2005	1,439	167	653	116	0	936	0.65
2006	1,163	192	838	254	0	1,283	1.10
2007	463	125	1,649	514	0	2,288	4.94
2008	1,074	414	827	290	0	1,531	1.42
2009	903	84	448	65	0	597	0.66
2010	1,024	209	653	198	0	1,059	1.03
2011	1,942	137	1,088	305	0	1,530	0.79
2012	1,110	64	419	260	0	743	0.67
2013	750	110	660	148	0	919	1.23
2014	746	142	376	13	0	532	0.71
2015	1,285	26	34	206 ²		266 ²	0.21 ²
2016	790	6	523 ²				
2017	971	32 ²					
2018	500						
2019	51						
2020	740 ²						
Mean	1,140	101	830	353	3	1,301	1.57

1. The geometric mean jack (age-3) proportion of spawning escapement from 1999-2020 was 0.09.

2. Preliminary.

Table 6. Adult-to-adult productivity indices for American River wild/natural stock.

Brood Year	Estimated Spawners	Estimated Yakima R. Mouth Returns					Returns/ Spawner
		Age-3	Age-4	Age-5	Age-6	Total	
1984	187	54	301	458	0	813	4.36
1985	337	81	149	360	0	590	1.75
1986	1,457	36	134	329	11	509	0.35
1987	567	12	71	134	0	216	0.38
1988	827	19	208	661	5	892	1.08
1989	524	11	69	113	0	193	0.37
1990	425	15	113	84	0	213	0.50
1991	414	3	5	22	0	30	0.07
1992	335	23	157	237	0	417	1.24
1993	721	8	218	405	8	639	0.89
1994	230	7	36	16	0	59	0.26
1995	98	33	32	98	0	163	1.65
1996	159	30	176	760	0	967	6.07
1997	371	13	1,543	610	0	2,166	5.84
1998	414	120	766	1,136	0	2,022	4.88
1999	61	72	99	163	0	334	5.50
2000	250	60	163	110	0	333	1.33
2001	1,917	18	364	256	0	638	0.33
2002	1,180	19	279	257	0	555	0.47
2003	1,192	23	183	440	0	646	0.54
2004	318	121	52	33	0	206	0.65
2005	464	79	173	127	0	378	0.81
2006	509	45	308	451	0	805	1.58
2007	523	57	645	493	0	1,194	2.28
2008	504	239	461	465	0	1,165	2.31
2009	213	60	143	44	0	247	1.16
2010	467	172	326	173	0	671	1.44
2011	1,118	71	646	236	0	953	0.85
2012	789	41	261	253	0	555	0.70
2013	619	76	412	53	0	542	0.88
2014	385	103	87	37		227	0.59
2015	819	7	61	120 ¹		188 ¹	0.23 ¹
2016	542	12	195 ¹				
2017	703	14 ¹					
2018	134						
2019	107						
2020	319 ¹						
Mean	546	52	268	285	1	610	1.60

1. Preliminary.

Table 7. Adult-to-adult productivity indices for Naches/American aggregate (wild/natural) population.

Brood Year	Estimated Spawners	Estimated Yakima R. Mouth Returns				Total	Returns/ Spawner
		Age-3	Age-4	Age-5	Age-6		
1984	570	164	1,109	1,080	0	2,354	4.13
1985	1,020	213	667	931	0	1,811	1.77
1986	4,123	103	670	852	31	1,657	0.40
1987	1,729	39	231	400	0	669	0.39
1988	2,167	51	815	1,557	11	2,434	1.12
1989	1,517	39	332	371	0	741	0.49
1990	1,380	40	326	168	0	533	0.39
1991	1,121	10	32	144	127	314	0.28
1992	1,188	52	1,034	661	0	1,747	1.47
1993	1,865	53	603	817	17	1,489	0.80
1994	704	21	160	167	0	348	0.49
1995	223	73	201	498	0	771	3.46
1996	1,047	209	4,010	2,359	0	6,579	6.29
1997	1,133	220	4,644	1,377	0	6,241	5.51
1998	917	364	2,167	2,316	12	4,859	5.30
1999	418 ¹	185	369	279	0	833	1.99
2000	4,112	131	2,286	346	0	2,762	0.67
2001	5,829	144	1,598	785	0	2,526	0.43
2002	3,041	78	975	443	0	1,496	0.49
2003	2,592	75	387	1,028	0	1,489	0.57
2004	2,515	227	514	232	0	973	0.39
2005	1,904	246	845	268	0	1,359	0.71
2006	1,672	237	1,120	759	0	2,117	1.27
2007	986	182	2,239	1,033	0	3,454	3.50
2008	1,578	653	1,262	803	0	2,718	1.72
2009	1,117	144	542	116	0	802	0.72
2010	1,491	381	972	412	0	1,766	1.18
2011	3,060	208	1,693	559	0	2,459	0.80
2012	1,900	105	662	540	0	1,307	0.69
2013	1,369	186	1,046	226	0	1,459	1.07
2014	1,130	245	439	49		733	0.65
2015	2,103	33	96	355 ²		484 ²	0.23 ²
2016	1,332	18	688 ²				
2017	1,673	46 ²					
2018	634						
2019	158						
2020	1,059 ²						
Mean	1,686	152	1,052	685	6	1,915	1.54

1. The geometric mean jack (age-3) proportion of spawning escapement from 1999-2020 was 0.09.
2. Preliminary.

Estimated spawners at the CESRF are the total number of wild/natural fish collected at Roza Dam and taken to the CESRF for production brood stock. Total returns are based on the information compiled in Table 3 and at Roza dam sampling operations. Age composition for CESRF fish is estimated using scales and PIT tag detections from CESRF fish sampled passing upstream through the Roza Dam adult monitoring facility.

Table 8. Adult-to-adult productivity for Cle Elum SRF spring Chinook.

Brood Year	Estimated Spawners	Estimated Yakima R. Mouth Returns				Returns/ Spawner
		Age-3	Age-4	Age-5	Total	
1997	261	741	7,753	176	8,670	33.22
1998	408	1,242	7,939	602	9,782	23.98
1999	738 ¹	134	714	16	864	1.17
2000	567	1,103	3,647	70	4,819	8.50
2001	595	396	845	9	1,251	2.10
2002	629	345	1,886	69	2,300	3.66
2003	441	121	800	12	932	2.11
2004	597	805	3,101	116	4,022	6.74
2005	510	1,305	3,052	21	4,378	8.58
2006	419	3,038	5,812	264	9,114	21.75
2007	449	1,277	5,174	108	6,558	14.61
2008	457	2,344	4,567	65	6,976	15.27
2009	486	461	2,663	58	3,181	6.55
2010	336	1,495	3,183	30	4,707	14.01
2011	377	1,233	2,340	34	3,607	9.57
2012	374	221	1,492	10	1,723	4.61
2013	398	802	1,993	0	2,795	7.02
2014	384	1,008	1,447	7	2,463	6.41
2015	442	314	877	0 ²	1,191 ²	2.70 ²
2016	376	287	771 ²		1,058 ²	2.81 ²
2017	382	349 ²				
2018	294					
2019	306					
2020	405 ²					
Mean	443	906	3,003	88	4,020	6.84 ³

1. 357 or 48% of these fish were jacks.
2. Preliminary.
3. Geometric mean.

Age Composition

Comparisons of the age composition in the Roza adult monitoring facility (RAMF) samples and spawning ground carcass recovery samples show that older, larger fish are recovered as carcasses on the spawning grounds at significantly higher rates than younger, smaller fish (Knudsen et al. 2003 and Knudsen et al. 2004). Based on historical scale-sampled carcass recoveries between 1986 and 2016 (there were no carcass recoveries in 2017 or 2018), age composition of American River spring Chinook has averaged 1, 44, 54, and 1 percent age-3, -4, -5, and -6, respectively (Table 9). Naches system spring Chinook averaged 2, 61, 36 and 0.5 percent age-3, -4, -5 and -6, respectively (Table 10). The upper Yakima River natural origin fish averaged 8, 88, and 4 percent age-3, -4, and -5, respectively (Table 11). While these ages are biased toward the older age classes, we believe the bias is approximately equal across populations and is a good relative indicator of differences in age composition between populations. The data show distinct differences with the American River population having the oldest age of maturation, followed closely by the Naches system and then the upper Yakima River which has significantly more age-3's, fewer age-5's and no age-6 fish.

Table 9. Percentage by sex and age of American River wild/natural spring Chinook carcasses sampled on the spawning grounds and sample size (n), 1986-present.

Return Year	Males					Females					Total			
	3	4	5	6	n	3	4	5	6	n	3	4	5	6
1986		23.8	76.2		21		8.9	86.7	4.4	45		13.6	83.3	3.0
1987		70.8	25.0	4.2	24		42.9	57.1		21		57.8	40.0	2.2
1988			100.0		1		100.0			1		33.3	66.7	
1989		39.6	60.4		48		10.0	90.0		50		24.5	75.5	
1990	2.5	25.0	72.5		40		28.3	71.7		46	1.2	26.7	72.1	
1991		23.8	76.2		42		13.3	86.7		60		17.6	82.4	
1992		71.2	23.1	5.8	52		45.8	54.2		48		59.0	38.0	3.0
1993	4.8	14.3	81.0		21		8.0	92.0		75	1.0	9.4	89.6	
1994		44.4	55.6		18		50.0	46.7	3.3	30		49.0	49.0	2.0
1995	14.3	14.3	71.4		7			100.0		13	5.0	5.0	90.0	
1996		100.0			2		83.3	16.7		6		87.5	12.5	
1997		40.0	60.0		5		22.2	64.4	13.3	45		24.0	64.0	12.0
1998		12.1	87.9		33		6.6	93.4		76		8.3	91.7	
1999		100.0			2		40.0	40.0	20.0	5		57.1	28.6	14.3
2000		66.7	33.3		15		61.5	38.5		13		64.3	35.7	
2001		65.6	34.4		90		67.9	32.1		106		67.0	33.0	
2002	1.7	53.4	44.8		58		56.4	43.6		110	0.6	55.4	44.0	
2003		8.1	91.9		74		7.9	92.1		151		8.0	92.0	
2004		100.0			3		20.0	80.0		5		50.0	50.0	
2005		64.7	35.3		17		84.0	16.0		25		76.7	23.3	
2006		61.5	38.5		13		48.6	51.4		35		52.1	47.9	
2007	10.5	31.6	57.9		19		43.8	56.3		48	3.0	40.3	56.7	
2008		8.7	91.3		23		11.9	88.1		42		10.6	89.4	
2009	30.8	69.2			13		75.0	25.0		16	13.8	72.4	13.8	
2010	6.3	56.3	37.5		16		75.0	25.0		32	2.0	69.4	28.6	
2011		40.0	60.0		10		63.2	36.8		19		58.8	41.2	
2012		50.0	50.0		14		47.8	52.2		16		48.3	51.7	
2013	11.1	11.1	77.8		9		26.9	73.1		26	2.9	22.9	74.3	
2014	5.6	77.8	16.7		18		90.9	9.1		33	2.0	86.3	11.8	
2015	7.4	74.1	18.5		27		78.3	21.7		46	2.7	76.7	20.5	
2016		28.6	71.4		14		65.4	34.6		26		52.5	47.5	
2017							No carcasses were sampled							
2018							No carcasses were sampled							
2019							Only 1 carcass sampled due to low run size							
2020	50.0	50.0			2		100.0			3	20.0	80.0		
Mean	3.1	46.7	50.0	0.3			44.6	54.0	1.3		1.1	44.7	53.1	1.2

Table 10. Percentage by sex and age of Naches River wild/natural spring Chinook carcasses sampled on the spawning grounds and sample size (n), 1986-present.

Return Year	Males					Females					Total			
	3	4	5	6	n	3	4	5	6	n	3	4	5	6
1986	5.0	60.0	30.0	5.0	20		33.3	64.3	2.4	42	1.6	41.9	53.2	3.2
1987	5.9	76.5	11.8	5.9	17		69.0	31.0		42	1.7	71.7	25.0	1.7
1988		50.0	50.0		8	5.6	38.9	55.6		18	3.3	46.7	50.0	
1989		70.2	29.8		47		34.9	63.5	1.6	63		50.0	49.1	0.9
1990	9.1	60.6	30.3		33	10.7	57.1	32.1		28	11.1	57.1	31.7	
1991	4.3	52.2	43.5		23		13.3	86.7		45	1.5	26.5	72.1	
1992	4.0	80.0	12.0	4.0	25		70.6	29.4		34	1.7	75.0	21.7	1.7
1993		42.3	57.7		26		18.6	81.4		43		28.6	71.4	
1994		50.0	50.0		4		30.0	70.0		10		35.7	64.3	
1995		25.0	75.0		4		28.6	71.4		7		33.3	66.7	
1996		100.0			17		75.0	25.0		16		87.9	12.1	
1997	2.9	70.6	20.6	5.9	34		57.1	36.7	6.1	49	1.2	62.7	30.1	6.0
1998		29.4	70.6		17		27.9	72.1		43		30.6	69.4	
1999	12.5	62.5	25.0		8		33.3	66.7		9	5.9	47.1	47.1	
2000	1.7	94.9	3.4		59		92.2	7.8		77	0.7	93.4	5.9	
2001	1.7	72.9	25.4		59		61.0	39.0		118	0.6	65.2	34.3	
2002	2.1	78.7	19.1		47		63.3	36.7		98	0.7	66.9	32.4	
2003	7.8	25.0	67.2		64	1.1	18.9	80.0		95	3.8	21.4	74.8	
2004	7.5	87.5	5.0		40		91.3	8.7		92	2.3	89.5	8.3	
2005		81.8	18.2		11		83.8	16.2		37		83.7	16.3	
2006		61.5	38.5		13		61.5	38.5		13		61.5	38.5	
2007		75.0	25.0		4		57.9	42.1		19		60.9	39.1	
2008	36.4	45.5	18.2		11		87.0	13.0		23	11.8	73.5	14.7	
2009	7.1	71.4	21.4		14		76.9	23.1		26	2.4	73.2	24.4	
2010	4.5	90.9	4.5		22		83.3	16.7		42	2.9	85.3	11.8	
2011	11.5	80.8	7.7		26		78.9	21.1		19	6.3	81.3	12.5	
2012	11.8	41.2	47.1		17		64.4	33.3		45	4.8	58.7	36.5	
2013	15.4	53.8	30.8		13		56.3	43.8		16	6.7	56.7	36.7	
2014		86.7	13.3		15		92.3	7.7		26		90.9	9.1	
2015		100.0			10		75.0	25.0		16		84.6	15.4	
2016		25.0	75.0		4		64.3	35.7		14		57.9	42.1	
2017							No carcasses were sampled							
2018							No carcasses were sampled							
2019							No carcasses were sampled							
2020		100.0			1		100.0			1		100.0		
Mean	4.9	64.6	29.9	0.7		0.6	57.9	41.1	0.3		2.3	61.3	36.0	0.4

Table 11. Percentage by sex and age of upper Yakima River wild/natural spring Chinook carcasses sampled on the spawning grounds and sample size (n), 1986-present.

Return Year	Males				Females				Total		
	3	4	5	n	3	4	5	n	3	4	5
1986		100.0		12		94.1	5.9	51		95.2	4.8
1987	10.8	81.5	7.7	65		77.8	22.2	126	3.7	79.1	17.3
1988	22.5	70.0	7.5	40	10.4	75.0	14.6	48	15.6	73.3	11.1
1989	0.8	93.1	6.2	130	0.4	95.5	4.1	246	0.5	94.7	4.8
1990	6.3	88.4	5.3	95	2.1	94.8	3.1	194	3.4	92.8	3.8
1991	9.1	87.3	3.6	55		89.2	10.8	111	3.0	88.6	8.4
1992	2.4	91.6	6.0	167		98.1	1.9	315	0.8	95.9	3.3
1993	4.0	90.0	6.0	50	0.9	92.0	7.1	112	1.9	91.4	6.8
1994		100.0		16		98.0	2.0	50		98.5	1.5
1995	20.0	80.0		5		100.0		12	5.6	94.4	
1996	9.1	89.6	1.3	154	0.7	98.2	1.1	282	3.7	95.2	1.1
1997		96.7	3.3	61		96.3	3.7	136		96.4	3.6
1998	14.3	85.7		21	5.3	86.8	7.9	38	8.5	86.4	5.1
1999	61.8	38.2		34		94.4	5.6	36	31.0	66.2	2.8
2000	2.8	97.2		72		100.0		219	1.0	99.0	
2001	2.7	89.2	8.1	37		83.6	16.4	122	0.6	85.0	14.4
2002	2.4	58.5	39.0	41	3.6	87.5	8.9	56	5.1	73.7	21.2
2003	60.5	39.5		38	4.3	82.6	13.0	23	39.3	55.7	4.9
2004	6.5	93.5		108	0.0	99.5	0.5	198	2.3	97.4	0.3
2005	9.2	90.0		120	1.4	97.2	1.4	214	4.2	94.7	1.2
2006	23.7	74.6		59	2.3	96.5	1.2	86	11.0	87.6	1.4
2007	17.1	82.9		76	0.9	93.8	5.4	112	7.4	89.4	3.2
2008	11.8	88.2		34	0.0	95.8	4.2	24	6.9	91.4	1.7
2009	47.7	52.3		111	2.2	95.6	2.2	45	34.6	64.7	0.6
2010	27.7	72.3		47		100.0		71	11.0	89.0	
2011	37.5	62.5		16		100.0		27	13.6	86.4	
2012	25.0	75.0		8	7.7	92.3		13	14.3	85.7	
2013						100.0		8		100.0	
2014	3.3	96.7		30		100.0		59	1.1	98.9	
2015	carcass surveys discontinued as Roza samples deemed adequate										
Mean	15.7	80.9	3.4		1.5	93.6	4.9		7.9	87.8	4.3

Carcasses from upper Yakima River CESRF origin fish allowed to spawn naturally have also been sampled since age-4 adults began returning in 2001. These fish averaged 13, 85, and 1 percent age-3, -4, and -5, respectively (Table 12) from 2001-2014 compared to 8, 88, and 4.3 percent respectively for their wild/natural counterparts in the upper Yakima for the same years (Table 11). The observed difference in age distribution between wild/natural and CESRF sampled on the spawning grounds may be due in part to the carcass recovery bias described above. A better comparison of age distribution between upper Yakima wild/natural and CESRF fish is from samples collected at Roza Dam which are displayed in Tables 13 and 14. However, it must be noted that jacks (age-3 males) were collected at Roza in proportion to run size from 1997 to 1999, but from 2000-present we have attempted to collect them at their mean brood representation rate (approximately 7% of the spawning population). Age-3 females do occur rarely in the Upper Yakima population, but it is likely that the data in Table 13 slightly over-represent the proportion of age-3 females due to human error associated with scale collection, handling, processing, and management and entry of these data.

Table 12. Percentage by sex and age of upper Yakima River CESRF spring Chinook carcasses sampled on the spawning grounds and sample size (n), 2001-present.

Return Year	Males				Females				Total		
	3	4	5	n	3	4	5	n	3	4	5
2001	23.5	76.5		34	0.9	99.1		108	6.3	93.7	
2002	8.0	81.3	10.7	75		88.6	11.4	140	2.8	86.2	11.1
2003	100.0			1		100.0		1	50.0	50.0	
2004	9.5	90.5		21		98.0	2.0	51	2.8	95.8	1.4
2005	42.9	57.1		21		90.9	4.5	22	23.3	74.4	2.3
2006	26.7	73.3		15		100.0		43	6.9	93.1	
2007	66.7	33.3		6		100.0		11	23.5	76.5	
2008				0		100.0		1		100.0	
2009	60.0	40.0		5				0	60.0	40.0	
2010	28.6	71.4		7		100.0		11	11.1	88.9	
2011	37.5	62.5		16	4.5	95.5		22	18.4	81.6	
2012		100.0		4	5.3	94.7		19	4.3	95.7	
2013		100.0		1		100.0		7		100.0	
2014		100.0		20		100.0		62	1.2	98.8	
2015	carcass surveys discontinued as Roza samples deemed adequate										
Mean ¹	25.3	73.8	0.9		0.5	97.2	1.8		13.4	85.4	1.2

1. Excludes years where sample size < 5.

Table 13. Percentage by sex and age of upper Yakima River wild/natural spring Chinook collected for brood stock at Roza Dam and sample size (n), 1997-present.

Return Year	Males				Females				Total		
	3	4	5	n	3	4	5	n	3	4	5
1997	4.5	92.0	3.4	88		94.6	5.4	111	2.0	93.5	4.5
1998	22.4	73.1	4.5	134		91.6	8.4	179	9.6	83.7	6.7
1999	71.1	26.1	2.8	425		92.6	7.4	215	48.8	47.0	4.2
2000	17.8	81.7	0.4	230		98.7	1.3	313	7.5	91.5	0.9
2001	12.4	77.4	10.3	234	0.9	90.5	8.5	328	5.7	85.2	9.2
2002	16.4	78.3	5.3	226	0.6	94.8	4.7	343	6.9	88.2	4.9
2003	27.4	60.2	12.4	201		83.3	16.7	228	12.8	72.6	14.7
2004	15.1	84.5	0.4	239	0.3	99.0	0.7	305	6.8	92.6	0.6
2005	15.5	82.3	2.2	181	0.4	97.1	2.5	276	6.3	91.2	2.4
2006	11.1	77.4	11.5	226		89.4	10.6	255	5.2	83.8	11.0
2007	13.6	74.7	11.7	162		87.8	12.2	255	5.3	82.7	12.0
2008	20.0	77.4	2.6	190		95.6	4.4	252	8.6	87.8	3.6
2009	17.4	81.2	1.4	207	0.8	96.1	3.1	258	8.2	89.5	2.4
2010	20.0	79.4	0.6	155	0.4	99.3	0.4	285	7.3	92.3	0.5
2011	18.1	81.3	0.5	182	0.8	95.3	3.8	236	8.4	89.2	2.4
2012	12.5	86.5	1.0	104		97.4	2.6	189	4.4	93.5	2.0
2013	18.0	77.6	4.3	161	0.0	96.2	3.8	183	8.4	87.5	4.1
2014	20.9	76.3	2.8	177	0.0	97.8	2.2	184	10.2	87.3	2.5
2015	9.3	89.4	1.2	161	0.0	98.7	1.3	231	3.8	94.9	1.3
2016	12.5	81.6	5.9	152	0.5	95.2	4.3	210	5.5	89.5	5.0
2017	13.7	84.9	1.4	146	1.0	97.9	1.0	194	6.5	92.4	1.2
2018	17.6	79.4	2.9	102	0.0	95.8	4.2	144	7.3	89.0	3.7
2019	13.2	86.8	0.0	76	0.7	97.3	2.0	149	4.9	93.8	1.3
2020	9.6	89.6	0.8	125	0.0	97.8	2.2	183	3.9	94.5	1.6
Mean	17.9	78.3	3.8		0.3	95.0	4.7		8.5	87.2	4.3

Table 14. Percentage by sex and age of upper Yakima River CESRF spring Chinook collected for research or brood stock at Roza Dam and sample size (n), 2001-present.

Return Year	Males				Females				Total		
	3	4	5	n	3	4	5	n	3	4	5
2001	12.5	87.5		40		100.0		75	5.1	94.9	
2002	14.7	83.8	1.5	68		98.3	1.7	115	5.5	92.9	1.6
2003	36.1	34.7	29.2	72		61.2	38.8	67	18.7	47.5	33.8
2004	19.6	80.4		46		100.0		60	8.5	91.5	
2005	17.8	75.6	6.7	45		88.1	11.9	59	7.7	82.7	9.6
2006	18.3	80.0	1.7	60		100.0		65	8.8	90.4	0.8
2007	33.3	60.8	5.9	51		87.5	12.5	56	15.9	74.8	9.3
2008	50.0	50.0		40		100.0		56	20.8	79.2	
2009	25.4	71.2	3.4	59	1.2	97.6	1.2	84	11.2	86.7	2.1
2010	27.9	72.1		61		99.0	1.0	100	10.6	88.8	0.6
2011	21.2	72.7	6.1	66	0.9	97.2	1.9	107	8.7	87.9	3.5
2012	13.0	85.2	1.9	54		97.0	3.0	101	4.5	92.9	2.6
2013	17.9	80.6	1.5	67	1.1	96.7	2.2	92	8.2	89.9	1.9
2014	31.9	66.0	2.1	47	0.0	100.0	0.0	33	18.8	80.0	1.3
2015	33.3	66.7	0.0	27	0.0	97.9	2.1	48	12.0	86.7	1.3
2016	26.5	69.4	4.1	49	0.0	100.0	0.0	47	13.5	84.4	2.1
2017	43.6	56.4	0.0	39	0.0	100.0	0.0	66	16.2	83.8	
2018	28.9	71.1	0.0	38	0.0	100.0	0.0	38	14.5	85.5	
2019	26.3	73.7	0.0	19	3.5	96.5	0.0	57	9.2	90.8	
2020	12.5	87.5	0.0	8	0.0	100.0	0.0	14	4.5	95.5	
Mean	25.5	71.3	3.2		0.3	95.9	3.8		11.1	85.3	3.5

Sex Composition

In the American River, the mean proportion of males to females in wild/natural carcasses sampled on the spawning grounds from 1986-2016 was 41:59 for age-4 and 33:67 for age-5 spring Chinook (Table 15). In the Naches River, the mean proportion of males to females was 41:59 for age-4 and 27:73 for age-5 fish (Table 16). In the upper Yakima River, the mean proportion of males to females was 33:67 for age-4 and 23:77 for age-5 fish (Table 17). Collection of carcass samples from the spawning grounds throughout the Yakima Basin did not occur in 2017-2019 and very few carcasses were sampled in 2020.

For upper Yakima fish collected at Roza Dam for brood stock or research purposes from 1997-2020, the mean proportion of males to females was 38:62 and 35:65 for age-4 fish from the wild/natural and CESRF populations, respectively (Tables 19 and 20). For these same samples, the mean proportion of males to females was 35:65 and 41:59 for age-5 fish from the wild/natural and CESRF populations (excluding years with very small age-5 sample sizes), respectively (Tables 19 and 20). For adult fish, the mean proportion of males to females in spawning ground carcass recoveries was substantially lower than the ratio found at RAMF (Tables 17 and 19), indicating that sex ratios estimated from hatchery origin carcass recoveries were biased due to female carcasses being recovered at higher rates than male carcasses (Knudsen et al, 2003 and 2004). Again, despite these biases, we believe these data are good relative indicators of differences in sex composition between populations and between years.

Sample sizes for Tables 15-20 were given in Tables 9-14. As noted earlier, few age-6 fish are found in carcass surveys and those that have been found were located in the American and Naches systems. The data indicate that age-3 females may occasionally occur in the upper Yakima and, to a lesser extent, the Naches systems.

Table 15. Percent of American River wild/natural spring Chinook carcasses sampled on the spawning grounds by age and sex, 1986-present.

Return Year	Age-3		Age-4		Age-5		Age-6	
	M	F	M	F	M	F	M	F
1986			55.6	44.4	29.1	70.9		100.0
1987			65.4	34.6	33.3	66.7	100.0	
1988			0.0	100.0	100.0	0.0		
1989			79.2	20.8	39.2	60.8		
1990	100.0		43.5	56.5	46.8	53.2		
1991			55.6	44.4	38.1	61.9		
1992			62.7	37.3	31.6	68.4	100.0	
1993	100.0		33.3	66.7	19.8	80.2		
1994			34.8	65.2	41.7	58.3		100.0
1995	100.0		100.0	0.0	27.8	72.2		
1996			28.6	71.4	0.0	100.0		
1997			16.7	83.3	9.4	90.6		100.0
1998			44.4	55.6	29.0	71.0		
1999			50.0	50.0	0.0	100.0		100.0
2000			55.6	44.4	50.0	50.0		
2001			45.0	55.0	47.7	52.3		
2002	100.0		33.3	66.7	35.1	64.9		
2003			33.3	66.7	32.9	67.1		
2004			75.0	25.0	0.0	100.0		
2005			34.4	65.6	60.0	40.0		
2006			32.0	68.0	21.7	78.3		
2007	100.0		22.2	77.8	28.9	71.1		
2008			28.6	71.4	36.2	63.8		
2009			42.9	57.1	0.0	100.0		
2010			27.3	72.7	42.9	57.1		
2011			25.0	75.0	46.2	53.8		
2012			24.1	75.9	22.6	77.4		
2013			12.5	87.5	26.9	73.1		
2014			31.8	68.2	50.0	50.0		
2015			35.7	64.3	33.3	66.7		
2016			19.0	81.0	52.6	47.4		
2017			No carcasses were sampled					
2018			No carcasses were sampled					
2019			Only 1 carcass sampled; low return					
2020	100.0		25.0	75.0				
mean			40.2	59.8	33.3	66.7		

Table 16. Percent of Naches River wild/natural spring Chinook carcasses sampled on the spawning grounds by age and sex, 1986-present.

Return Year	Age-3		Age-4		Age-5		Age-6			
	M	F	M	F	M	F	M	F		
1986	100.0		46.2	53.8	18.2	81.8	50.0	50.0		
1987	100.0		31.0	69.0	13.3	86.7	100.0			
1988		100.0	36.4	63.6	28.6	71.4				
1989			60.0	40.0	25.9	74.1		100.0		
1990	50.0	50.0	55.6	44.4	52.6	47.4				
1991	100.0		66.7	33.3	20.4	79.6				
1992	100.0		45.5	54.5	23.1	76.9	100.0			
1993			57.9	42.1	30.0	70.0				
1994			40.0	60.0	22.2	77.8				
1995			33.3	66.7	37.5	62.5				
1996			58.6	41.4		100.0				
1997	100.0		46.2	53.8	28.0	72.0	40.0	60.0		
1998			29.4	70.6	27.9	72.1				
1999	100.0		62.5	37.5	25.0	75.0				
2000	100.0		44.1	55.9	25.0	75.0				
2001	100.0		37.4	62.6	24.6	75.4				
2002	100.0		37.4	62.6	20.0	80.0				
2003	83.3	16.7	47.1	52.9	36.1	63.9				
2004	100.0		29.4	70.6	20.0	80.0				
2005			22.5	77.5	25.0	75.0				
2006			50.0	50.0	50.0	50.0				
2007			21.4	78.6	11.1	88.9				
2008	100.0		20.0	80.0	40.0	60.0				
2009	100.0		33.3	66.7	33.3	66.7				
2010	100.0		36.4	63.6	12.5	87.5				
2011	100.0		58.3	41.7	33.3	66.7				
2012	66.7	33.3	19.4	80.6	34.8	65.2				
2013	100.0		43.8	56.3	36.4	63.6				
2014			35.1	64.9	50.0	50.0				
2015			45.5	54.5		100.0				
2016			10.0	90.0	37.5	62.5				
2017			No carcasses were sampled							
2018			No carcasses were sampled							
2019			No carcasses were sampled							
2020			50.0	50.0						
mean			40.6	59.4	27.2	72.8				

Table 17. Percent of Upper Yakima River wild/natural spring Chinook carcasses sampled on the spawning grounds by age and sex, 1986-present.

Return Year	Age-3		Age-4		Age-5	
	M	F	M	F	M	F
1986			20.0	80.0		100.0
1987	100.0		35.1	64.9	15.2	84.8
1988	64.3	35.7	43.8	56.3	30.0	70.0
1989	50.0	50.0	34.0	66.0	44.4	55.6
1990	60.0	40.0	31.3	68.7	45.5	54.5
1991	100.0		32.7	67.3	14.3	85.7
1992	100.0		33.1	66.9	62.5	37.5
1993	66.7	33.3	30.4	69.6	27.3	72.7
1994			24.6	75.4		100.0
1995	100.0		25.0	75.0		
1996	87.5	12.5	33.3	66.7	40.0	60.0
1997			31.1	68.9	28.6	71.4
1998	60.0	40.0	35.3	64.7		100.0
1999	100.0		27.7	72.3		100.0
2000	100.0		24.2	75.8		
2001	100.0		24.4	75.6	13.0	87.0
2002	33.3	66.7	32.9	67.1	76.2	23.8
2003	95.8	4.2	44.1	55.9		100.0
2004	100.0		33.9	66.1		100.0
2005	78.6	21.4	34.2	65.8	25.0	75.0
2006	87.5	12.5	34.6	65.4	50.0	50.0
2007	92.9	7.1	37.5	62.5		100.0
2008	100.0		56.6	43.4		100.0
2009	98.1	1.9	57.4	42.6		100.0
2010	100.0		32.4	67.6		
2011	100.0		27.0	73.0		
2012	66.7	33.3	33.3	66.7		
2013				100.0		
2014	100.0	0.0	33.0	67.0		
2015	carcass surveys discontinued as Roza samples deemed adequate					
mean	85.7	14.3	33.0	67.0	22.5	77.5

Table 18. Percent of upper Yakima River CESRF spring Chinook carcasses sampled on the spawning grounds by age and sex, 2001-present.

Return Year	Age-3		Age-4		Age-5	
	M	F	M	F	M	F
2001	88.9	11.1	19.5	80.5		
2002	100.0		33.0	67.0	33.3	66.7
2003	100.0			100.0		
2004	100.0		27.5	72.5		100.0
2005	90.0	10.0	37.5	62.5		100.0
2006	100.0		20.4	79.6		
2007	100.0		15.4	84.6		
2008				100.0		
2009	100.0		100.0			
2010	100.0		31.3	68.8		
2011	85.7	14.3	32.3	67.7		
2012			18.2	81.8		
2013			12.5	87.5		
2014			24.4	75.6		
2015	carcass surveys discontinued as Roza samples deemed adequate					
mean	96.5	3.5	26.6	73.4		

Table 19. Percent of upper Yakima River wild/natural spring Chinook collected for brood stock at Roza Dam by age and sex, 1997-present.

Return Year	Age-3		Age-4		Age-5	
	M	F	M	F	M	F
1997	100.0		43.5	56.5	33.3	66.7
1998	100.0		37.4	62.6	28.6	71.4
1999	100.0		35.8	64.2	42.9	57.1
2000	100.0		37.8	62.2	20.0	80.0
2001	90.6	9.4	37.9	62.1	46.2	53.8
2002	94.9	5.1	35.3	64.7	42.9	57.1
2003	100.0		38.9	61.1	39.7	60.3
2004	97.3	2.7	40.1	59.9	33.3	66.7
2005	96.6	3.4	35.7	64.3	36.4	63.6
2006	100.0		43.4	56.6	49.1	50.9
2007	100.0		35.1	64.9	38.0	62.0
2008	100.0		37.9	62.1	31.3	68.8
2009	94.7	5.3	40.4	59.6	27.3	72.7
2010	96.9	3.1	30.3	69.7	50.0	50.0
2011	94.3	5.7	39.7	60.3	10.0	90.0
2012	100.0		32.8	67.2	16.7	83.3
2013	100.0		41.5	58.5	50.0	50.0
2014	100.0		42.9	57.1	55.6	44.4
2015	100.0		38.7	61.3	40.0	60.0
2016	95.0	5.0	38.3	61.7	50.0	50.0
2017	90.9	9.1	39.5	60.5	50.0	50.0
2018	100.0		37.0	63.0	33.3	66.7
2019	90.9	9.1	31.3	68.7	0.0	100.0
2020	100.0		38.5	61.5	20.0	80.0
mean	97.5	2.5	37.9	62.1	35.8	64.2

Table 20. Percent of Upper Yakima River CESRF spring Chinook collected for research or brood stock at Roza Dam by age and sex, 2001-present.

Return Year	Age-3		Age-4		Age-5	
	M	F	M	F	M	F
2001	100.0	0.0	31.8	68.2		
2002	100.0	0.0	33.5	66.5	33.3	66.7
2003	100.0	0.0	37.9	62.1	44.7	55.3
2004	100.0	0.0	38.1	61.9		
2005	100.0	0.0	39.5	60.5	30.0	70.0
2006	100.0	0.0	42.5	57.5	100.0	
2007	100.0	0.0	38.8	61.3	30.0	70.0
2008	100.0	0.0	26.3	73.7		
2009	93.8	6.3	33.9	66.1	66.7	33.3
2010	100.0	0.0	30.8	69.2		100.0
2011	93.3	6.7	31.6	68.4	66.7	33.3
2012	100.0		31.9	68.1	25.0	75.0
2013	92.3	7.7	37.8	62.2	33.3	66.7
2014	100.0	0.0	48.4	51.6	100.0	0.0
2015	100.0	0.0	27.7	72.3		
2016	100.0	0.0	42.0	58.0	100.0	0.0
2017	100.0	0.0	25.0	75.0		
2018	100.0	0.0	41.5	58.5		
2019	71.4	28.6	20.3	79.7		
2020	100.0	0.0	33.3	66.7		
mean	97.5	2.5	34.6	65.4	41.2	58.8

Size at Age

Prior to 1996, samplers were instructed to collect mid-eye to hypural plate (MEHP) lengths from carcasses surveyed on the spawning grounds. From 1996 to present the method was changed and post-eye to hypural plate (POHP) lengths have been recorded. Mean POHP lengths averaged 39, 62, and 76 cm for age-3, -4, and -5 males, and averaged 63 and 72 cm for age-4 and -5 females, respectively, from carcasses sampled on the spawning grounds in the American River from 1996-2016 (Table 21). In the Naches River, mean POHP lengths averaged 42, 60, and 76 cm for age-3, -4, and -5 males, and averaged 61 and 72 cm for age-4 and -5 females, respectively (Table 22). For wild/natural spring Chinook sampled on the spawning grounds in the upper Yakima River, mean POHP lengths averaged 44, 60, and 72 cm for age-3, -4, and -5 males, and averaged 59 and 69 cm for age-4 and -5 females, respectively (Table 23). Beginning in 2012, carcass sampling in the Upper Yakima was scaled back considerably as large numbers of escaping fish are sampled at Roza Dam (Tables 27-28). From 2001-2020, CESRF fish returning to the upper Yakima have been generally smaller in size-at-age than their wild/natural counterparts (Tables 25-28).

Table 21. Counts and mean mid-eye (MEHP) or post-orbital (POHP) to hypural plate lengths (cm) of American River wild/natural spring Chinook from carcasses sampled on the spawning grounds by sex and age, 1987-present.

Return Year	Males								Females					
	Age 3		Age 4		Age 5		Age 6		Age 4		Age 5		Age 6	
	Count	MEHP	Count	MEHP	Count	MEHP	Count	MEHP	Count	MEHP	Count	MEHP	Count	MEHP
1987			17	58.0	6	80.8	1.0	86.0	9	64.5	12	76.9		
1988					1	79.0			1	63.0				
1989			19	61.1	29	77.4			5	63.0	45	73.5		
1990	1	41.0	10	63.6	29	77.3			13	62.5	33	73.6		
1991			10	59.5	32	77.1			8	65.1	52	73.4		
1992			37	60.6	12	76.2	3.0	86.7	22	64.1	26	76.4		
1993	1	47.0	3	64.0	17	80.2			6	63.7	69	75.5		
1994			8	67.3	10	83.0			15	70.8	14	76.4	1	85.0
1995	1	44.4	1	70.0	4	83.5					12	76.4		
		POHP		POHP		POHP		POHP		POHP		POHP		POHP
1996			2	56.3					5	59.0	1	67.0		
1997 ¹			2	62.0	1	63.0			4	62.8	14	64.4	5	71.0
1998			4	58.3	29	79.1			5	64.0	71	73.4		
1999			2	50.5					2	61.0	2	73.0	1	77.0
2000			10	57.9	5	83.2			8	63.9	5	76.2		
2001			59	65.9	31	77.6			72	63.6	34	73.0		
2002	1	40.0	31	63.0	26	77.3			62	64.4	48	74.7		
2003			6	63.0	68	79.4			12	64.3	139	76.7		
2004			3	56.0					1	58.0	4	77.5		
2005			11	60.6	6	80.2			21	62.6	4	74.8		
2006			8	60.8	5	75.4			17	61.8	18	71.7		
2007	2	37.0	6	62.8	11	76.5			21	60.0	27	73.3		
2008			2	67.5	21	83.1			5	67.4	37	78.9		
2009	4	44.0	9	68.3					12	62.6	4	69.8		
2010	1	38.0	9	70.1	6	75.7			24	65.1	8	73.0		
2011			4	65.5	6	82.8			12	65.8	7	75.9		
2012			7	64.1	7	77.3			22	63.7	24	74.3		
2013	1	34.0	1	56.0	7	70.1			7	65.7	18	70.3		
2014	1	36.0	14	61.1	3	66.7			30	61.2	3	63.3		
2015	2	42.0	20	63.4	5	77.4			36	61.3	10	71.2		
2016			4	65.0	10	71.5			17	59.7	9	67.6		
2017-19			No samples						No samples					
2020	1	38.0	1	52.0					3	65.7				
Mean ²		38.7		61.8		76.2				62.8		72.4		74.0

¹ Carcasses sampled in 1997 had a mix of MEHP and POHP lengths taken. Only POHP samples are given here.

² Mean of mean values for 1996-2016 post-eye to hypural plate lengths.

Table 22. Counts and mean mid-eye (MEHP) or post-orbital (POHP) to hypural plate lengths (cm) of Naches River wild/natural spring Chinook from carcasses sampled on the spawning grounds by sex and age, 1987-present.

Return Year	Males								Females							
	Age 3		Age 4		Age 5		Age 6		Age 3		Age 4		Age 5		Age 6	
	Count	MEHP	Count	MEHP	Count	MEHP	Count	MEHP	Count	MEHP	Count	MEHP	Count	MEHP	Count	MEHP
1987	1	37.0	12	64.2	2	80.5	1.0	94.0			29	67.9	13	75.7		
1988			4	62.0	4	74.6			1	45.0	7	69.1	10	73.6		
1989			33	58.4	14	77.5					22	61.7	40	73.2	1	75.0
1990	3	53.0	20	59.4	10	75.9			3	51.7	16	60.9	9	73.7		
1991	1	31.0	12	56.3	10	72.8					6	62.5	39	71.1		
1992	1	42.0	20	58.8	3	72.3	1.0	83.0			24	62.4	10	71.7		
1993			11	60.0	15	77.7					8	63.3	35	72.5		
1994			2	62.5	2	77.0					3	63.7	7	73.1		
1995			1	59.0	3	73.0					2	64.0	5	73.8		
		POHP		POHP		POHP		POHP		POHP		POHP		POHP		POHP
1996			17	58.1							12	60.3	4	69.6		
1997 ¹	1	39.0	24	59.8	4	71.5	2.0	78.0			28	60.0	15	68.6	1	75.0
1998			5	57.8	12	75.0					12	61.1	31	71.6		
1999	1	40.0	5	61.2	2	73.0					3	58.7	6	75.0		
2000	1	35.0	56	58.2	2	84.0					71	59.5	6	72.8		
2001	1	45.0	43	61.4	15	73.4					72	62.2	46	74.5		
2002	1	40.0	37	63.6	9	77.3					62	62.4	36	71.8		
2003	5	41.4	16	62.2	43	79.4			1	41.0	18	62.8	76	75.6		
2004	3	46.0	35	59.8	2	74.5					84	61.5	8	75.8		
2005			9	60.1	2	78.0					31	61.7	6	71.7		
2006			8	56.9	5	76.0					8	63.8	5	71.2		
2007			3	61.3	1	67.0					11	56.9	8	72.1		
2008	4	42.0	5	59.6	2	81.5					20	62.0	3	78.7		
2009	1	43.0	10	67.9	3	76.3					20	63.9	6	73.2		
2010	1	40.0	20	60.5	1	77.0					35	61.7	7	71.4		
2011	3	44.3	21	61.9	2	78.0					15	60.4	4	76.8		
2012	2	51.5	7	67.3	8	75.8			1	41.0	29	61.6	15	71.1		
2013	2	37.0	7	56.1	4	75.0					9	58.7	7	71.3		
2014			13	61.8	2	71.0					24	56.7	2	67.5		
2015			10	59.3							12	60.4	4	65.8		
2016			1	47.0	3	77.0					9	53.9	5	68.8		
2017-19			No samples									No samples				
2020			1	50.0							1	53.0				
Mean ²		41.9		60.1		75.8		78.0		41.0		60.5		72.1		75.0

¹ Carcasses sampled in 1997 had a mix of MEHP and POHP lengths taken. Only POHP samples are given here.

² Mean of mean values for 1996-2016 post-eye to hypural plate lengths.

Table 23. Counts and mean mid-eye (MEHP) or post-orbital (POHP) to hypural plate lengths (cm) of upper Yakima River wild / natural spring Chinook from carcasses sampled on the spawning grounds by sex and age, 1986-present.

Return Year	Males						Females					
	Age 3		Age 4		Age 5		Age 3		Age 4		Age 5	
	Count	MEHP	Count	MEHP	Count	MEHP	Count	MEHP	Count	MEHP	Count	MEHP
1986			12	60.8					48	58.7	3	70.3
1987	7	45.3	53	58.5	5	73.0			96	59.3	28	70.6
1988	9	40.0	28	59.0	3	79.0	5	52.6	36	59.2	7	70.3
1989	1	50.0	121	59.7	8	70.6	1	40.0	235	58.6	10	67.2
1990	6	47.0	84	58.0	5	77.0	4	51.5	184	59.3	6	72.5
1991	5	39.6	48	56.2	2	67.5			99	57.6	12	68.8
1992	4	43.0	153	58.4	10	71.2			309	58.2	6	69.5
1993	2	44.0	45	60.7	3	75.0	1	56.0	101	59.5	8	70.3
1994			15	62.9					49	61.3	1	72.0
1995	1	43.0	4	62.0					12	61.4	0	
		POHP		POHP		POHP		POHP		POHP		POHP
1996	14	40.9	138	59.1	2	66.5	2	41.0	277	58.6	3	68.0
1997			59	59.3	2	74.0			131	58.6	5	69.4
1998	3	38.7	18	56.4			2	47.0	33	57.5	3	66.7
1999	21	38.8	13	57.4					34	58.9	2	69.8
2000	2	41.0	70	60.3					219	58.3	0	
2001	1	43.0	33	60.7	3	74.7			102	60.6	20	69.8
2002	1	44.0	24	64.9	16	69.3	2	46.0	49	62.5	5	70.2
2003	23	44.4	15	59.8					19	62.4	3	67.8
2004	7	47.3	101	59.9					197	58.7	1	67.0
2005	11	49.2	108	60.6	1	75.0	3	48.7	207	59.5	3	67.3
2006	14	41.8	44	59.4	1	72.0	2	39.5	82	58.3	1	71.0
2007	13	44.2	61	61.7					101	60.6	6	66.0
2008	3	48.3	29	60.5					22	59.7	1	77.0
2009	53	46.8	58	57.6			1	51.0	43	60.2	1	68.0
2010	13	47.7	34	60.5					70	59.5		
2011	6	47.0	10	58.9					27	59.3		
2012	2	44.5	6	58.0			1	47.0	12	57.5		
2013			No samples						8	56.6		
2014	1	45.0	29	61.2					59	61.3		
2015			carcass surveys discontinued as Roza samples deemed adequate									
Mean ¹		44.3		59.8		71.9		45.7		59.4		69.1

¹ Mean of mean values for 1996-2014 post-eye to hypural plate lengths.

Table 24. Counts and mean post-orbital to hypural plate (POHP) lengths (cm) of upper Yakima River CESRF spring Chinook from carcasses sampled on the spawning grounds by sex and age, 2001-present.

Return Year	Males						Females					
	Age 3		Age 4		Age 5		Age 3		Age 4		Age 5	
	Count	POHP	Count	POHP	Count	POHP	Count	POHP	Count	POHP	Count	POHP
2001	8	40.5	25	59.0	1	69.5	1	41.0	107	59.0		
2002	6	47.7	61	61.2	8	68.9			124	60.6	16	71.2
2003	1	42.0							1	69.0		
2004	2	52.0	19	60.8					50	57.9	1	68.0
2005	8	41.8	12	59.9			1	46.0	20	59.6	1	72.0
2006	4	42.3	11	54.0					43	57.0		
2007	4	44.3	2	58.5					11	60.1		
2008	0		0						1	58.0		
2009	3	47.7	2	---								
2010	2	44.0	5	61.8					11	55.5		
2011	6	40.7	10	59.1			1	46.0	21	59.0		
2012			4	63.0			1	50.0	18	57.3		
2013			1	---					7	53.6		
2014			20	60.8					62	59.0		
2015	carcass surveys discontinued as Roza samples deemed adequate											
Mean		44.3		59.8		69.2				58.9		70.4

Table 25. Counts and mean post-orbital to hypural plate (POHP) lengths (cm) of upper Yakima River wild/natural spring Chinook from carcasses sampled at the CESRF prior to spawning by sex and age, 1997-present.

Return Year	Males						Females					
	Age 3		Age 4		Age 5		Age 3		Age 4		Age 5	
	Count	POHP	Count	POHP	Count	POHP	Count	POHP	Count	POHP	Count	POHP
1997	4	39.7	81	59.7	3	73.3			105	60.5	6	68.9
1998	28	43.0	95	57.3	6	67.0			161	59.2	15	65.6
1999	124	41.4	75	59.5	10	64.6			199	60.4	16	67.4
2000	19	42.0	145	59.0	1	77.0			263	59.4	3	69.4
2001	17	42.9	115	59.6	14	74.1			196	60.5	19	69.8
2002	23	42.1	113	60.6	5	72.9	1	36.6	233	61.2	9	70.9
2003	37	42.7	92	60.4	19	73.7			164	61.4	31	69.4
2004	18	42.4	108	58.9	1	67.8			225	58.3	2	66.5
2005	19	42.1	113	60.0	2	67.3	1	42.6	223	59.8	5	67.8
2006	17	41.0	82	56.7	20	70.4			197	57.8	24	68.1
2007	20	44.6	108	58.8	17	67.6			181	59.4	24	67.2
2008	17	45.5	121	59.6	4	71.1			209	59.7	11	68.4
2009	16	44.4	122	61.5	3	69.3	1	50.4	206	60.3	6	68.0
2010	9	45.0	88	61.5	1	71.2			192	60.9		
2011	11	47.5	91	60.3	1	75.3	1	52.5	182	60.2	4	72.9
2012	13	43.7	83	59.8	1	62.4			178	59.3	5	66.6
2013	18	45.8	112	59.6	7	70.0			161	58.9	6	69.7
2014	27	43.3	112	61.3	5	70.0			173	59.9	4	63.1
2015	8	41.2	110	59.6	2	71.7			167	59.9	2	70.5
2016	16	45.9	110	61.4	8	68.9			159	60.4	7	68.0
2017	18	43.2	115	61.0	2	66.0	2	47.7	167	62.1	2	64.9
2018	17	40.5	77	59.2	3	66.0			132	58.9	6	62.9
2019	6	39.8	55	55.2			1	39.5	120	56.2	1	63.5
2020	12	39.7	105	55.9	1	71.1			173	55.9	4	62.3
Mean		42.9		59.4		69.9				59.6		67.5

Table 26. Counts and mean post-orbital to hypural plate (POHP) lengths (cm) of upper Yakima River CESRF spring Chinook from carcasses sampled at the CESRF prior to spawning by sex and age, 2001-present.

Return Year	Males						Females					
	Age 3		Age 4		Age 5		Age 3		Age 4		Age 5	
	Count	POHP	Count	POHP	Count	POHP	Count	POHP	Count	POHP	Count	POHP
2001			4	61.3					33	60.4		
2002	2	40.2	25	59.6					63	59.4	2	66.1
2003	17	42.6	16	57.8	15	74.0			31	59.7	19	70.4
2004	6	39.4	9	57.1					42	59.3		
2005	6	37.9	21	58.4	2	68.7			38	58.6	5	68.0
2006 ¹			3	57.2					3	56.3		
2007	8	40.4	18	59.3	1	71.4			35	58.2	5	67.6
2008	17	43.8	9	59.1					28	59.4		
2009	5	43.8	11	61.1					32	60.1	1	67.5
2010	11	41.8	18	59.2					40	61.0		
2011	4	43.4	10	62.7	1	79.2			32	60.4	2	71.7
2012	3	39.0	23	59.3	1	73.7			43	59.4	1	67.2
2013	2	45.7	24	60.3					32	57.3		
2014	7	39.2	21	61.8	1	70.2			32	60.5		
2015	7	38.9	17	58.5					42	59.2	1	66.7
2016	2	42.8	22	61.4	2	75.0			34	60.8		
2017	11	44.1	20	59.9					36	61.9		
2018	8	38.4	22	59.5					34	59.4		
2019	3	37.3	14	56.2					25	55.8		
2020	1	37.4	7	54.9					13	54.6		
Mean		40.9		59.2		73.2				59.1		68.2

¹ Few length samples were collected since these fish were not spawned in 2006.

Table 27. Counts and mean post-orbital to hypural plate (POHP) lengths (cm) of upper Yakima River wild/natural spring Chinook from fish sampled at Roza Dam by sex¹ and age, 1997-present.

Return Year	Males						Females					
	Age 3		Age 4		Age 5		Age 3		Age 4		Age 5	
	Count	POHP	Count	POHP	Count	POHP	Count	POHP	Count	POHP	Count	POHP
1997	4	39.6	81	60.6	2	73.3			121	60.5	10	70.6
1998	36	42.4	108	58.3	11	67.7	1	58.5	201	59.4	13	67.0
1999	350	40.7	80	59.4	11	67.5	2	46.8	256	60.3	19	68.3
2000	40	41.3	145	60.5	1	77.0	1	46.0	354	60.2	4	72.1
2001	32	42.9	111	61.9	28	73.8			371	61.2	24	70.7
2002	43	41.6	146	61.2	21	71.4	2	52.5	379	60.7	8	70.3
2003	54	43.3	52	64.6	18	75.3	1	51.0	262	61.9	45	71.2
2004	41	43.4	121	61.1	1	69.0			394	59.4	2	69.5
2005	35	43.2	134	61.1	5	74.2			307	60.8	6	68.3
2006	27	41.3	77	59.1	22	72.6	1	47.0	336	58.8	27	69.5
2007	31	42.9	83	60.8	18	69.8	1	50.0	280	60.5	34	69.7
2008	38	45.8	101	61.7	8	72.4			293	60.7	8	69.1
2009	36	45.3	125	63.4	4	71.5	3	52.7	297	61.9	8	69.9
2010	39	43.7	129	62.6	1	74.0	1	51.0	298	62.8	1	70.0
2011	42	46.7	154	61.2	3	77.3	2	53.0	235	61.9	10	75.3
2012	27	43.6	113	60.5	1	63.0			202	60.3	5	68.0
2013	31	45.4	132	59.9	8	70.6			181	59.8	7	70.6
2014	38	44.7	138	62.2	5	72.2			181	61.2	4	65.5
2015	16	44.0	150	61.2	3	72.0			245	61.2	3	71.7
2016	21	46.0	130	62.3	10	71.4			210	61.6	10	69.8
2017	21	43.3	128	61.3	2	66.5	2	48.0	195	62.5	2	66.0
2018	21	40.9	86	59.3	3	67.3			140	59.2	7	64.4
2019	11	40.9	67	57.7			1	42.0	148	58.6	4	70.3
2020	13	41.7	127	58.5	1	75.0			192	58.3	4	66.3
Mean		43.1		60.9		71.5		49.9		60.6		69.3

¹ Sex determined by visual observation prior to 2010 and by ultrasound from 2010 to present.

Table 28. Counts and mean post-orbital to hypural plate (POHP) lengths (cm) of upper Yakima River CESRF spring Chinook from fish sampled at Roza Dam by sex¹ and age, 2001-present.

Return Year	Males						Females					
	Age 3		Age 4		Age 5		Age 3		Age 4		Age 5	
	Count	POHP	Count	POHP	Count	POHP	Count	POHP	Count	POHP	Count	POHP
2001	473	39.9	548	59.5			1	58.0	1795	59.2		
2002	26	38.7	383	59.5	19	67.7			1152	59.1	15	66.1
2003	392	41.8	48	61.8	61	73.0	2	47.0	207	60.3	154	70.8
2004	48	40.3	100	60.5			1	44.0	351	59.2	2	71.0
2005	98	40.4	58	60.1	6	73.0			160	59.1	12	68.7
2006	26	40.4	89	58.0					318	57.4	2	70.5
2007	174	41.4	46	60.7	6	71.7	1	47.0	185	59.0	13	69.8
2008	93	44.8	60	60.7			2	54.5	191	60.1	1	67.0
2009	254	43.6	78	62.8	5	65.0	1	50.0	212	61.8	6	69.5
2010	106	42.5	196	61.0	1	67.0	1	60.0	361	61.8	1	72.0
2011	155	42.9	146	60.9	8	73.5	2	57.5	265	61.5	13	73.4
2012	45	40.6	131	59.3	3	65.7	1	45.0	250	59.9	6	69.2
2013	92	44.4	122	59.0	3	70.0			163	58.8	4	69.3
2014	78	42.8	111	61.0	2	71.0			163	60.5	3	71.7
2015	19	41.2	90	59.5					146	60.3	3	72.0
2016	86	44.5	73	61.1	3	77.3	2	48.0	102	61.2	1	65.0
2017	83	43.9	47	61.6					160	62.3	1	67.0
2018	24	39.3	56	58.4			1	41.0	86	59.4		
2019	18	41.4	35	57.5			1	46.0	84	57.7	1	76.0
2020	35	41.7	25	57.4					52	57.7		
Mean		41.7		60.0		70.4		49.8		59.8		69.9

¹Sex determined by visual observation prior to 2010 and by ultrasound from 2010 to present.

Migration Timing

Wild/natural spring Chinook adults returning to the upper Yakima River have generally shown earlier passage timing at Roza Dam than CESRF spring Chinook (Figures 2 and 3).




Figure 3. Proportionate passage timing at Roza Dam of wild/natural and CESRF adult spring Chinook (including jacks), 2011-2020.

Table 29. Comparison of 5%, median (50%), and 95% passage dates of wild/natural and CESRF adult spring Chinook (including jacks) at Roza Dam, 1997-Present.

Year	Wild/Natural Passage			CESRF Passage		
	5%	Median	95%	5%	Median	95%
1997	10-Jun	17-Jun	21-Jul			
1998	22-May	10-Jun	10-Jul			
1999	31-May	24-Jun	4-Aug			
2000	12-May	24-May	12-Jul	21-May ¹	15-Jun ¹	27-Jul ¹
2001	4-May	23-May	11-Jul	8-May	28-May	15-Jul
2002	16-May	10-Jun	6-Aug	20-May	13-Jun	12-Aug
2003	13-May	11-Jun	19-Aug	13-May	10-Jun	24-Aug
2004	4-May	20-May	24-Jun	5-May	22-May	26-Jun
2005	9-May	22-May	23-Jun	15-May	31-May	2-Jul
2006	1-Jun	14-Jun	18-Jul	3-Jun	18-Jun	19-Jul
2007	16-May	5-Jun	9-Jul	24-May	14-Jun	19-Jul
2008	27-May	9-Jun	9-Jul	31-May	17-Jun	14-Jul
2009	31-May	14-Jun	17-Jul	2-Jun	19-Jun	17-Jul
2010	11-May	30-May	5-Jul	12-May	2-Jun	9-Jul
2011	6-Jun	23-Jun	16-Jul	9-Jun	24-Jun	15-Jul
2012	30-May	14-Jun	9-Jul	30-May	13-Jun	8-Jul
2013	22-May	4-Jun	3-Jul	24-May	8-Jun	8-Jul
2014	15-May	1-Jun	2-Jul	18-May	5-Jun	8-Jul
2015 ²	4-May	16-May	31-Aug	5-May	18-May	31-Aug
2016	17-May	29-May	28-Jun	21-May	4-Jun	20-Jul
2017	1-Jun	14-Jun	3-Jul	6-Jun	20-Jun	14-Jul
2018	1-Jun	8-Jun	18-Jul	2-Jun	14-Jun	16-Jul
2019	22-May	31-May	29-Jul	25-May	5-Jun	20-Aug
2020	21-May	11-Jun	9-Aug	27-May	23-Jun	23-Aug

1. In 2000 all returning CESRF fish were age-3 (jacks).
2. Mean daily water temperatures at Kiona (rkm 40 from the mouth of the Yakima R.) exceeded 70° F every day from May 21 to August 29, 2015 (source U.S. BOR hydromet database) causing delayed passage for late migrating fish.

Spawning Timing

Median spawn timing for CESRF spring Chinook is earlier than that observed for wild/natural fish in the Upper Yakima River. These differences are due in part to environmental conditions and spawning procedures at the hatchery. It must also be noted that spawning dates in the wild are only a coarse approximation, derived from weekly redd counts not actual dates of redd deposition. A clear delineation of wild/natural spawn timing between subbasins is apparent, with American River fish spawning about 1 month earlier than Naches Basin fish which spawn about 2 weeks earlier than Upper Yakima fish.

Table 30. Median spawn¹ dates for spring Chinook in the Yakima Basin.

Year	American	Naches	Upper Yakima	CESRF
1988	14-Aug	7-Sep	3-Oct	
1989	14-Aug	7-Sep	19-Sep	
1990	14-Aug	12-Sep	25-Sep	
1991	12-Aug	12-Sep	24-Sep	
1992	11-Aug	10-Sep	22-Sep	
1993	9-Aug	8-Sep	27-Sep	
1994	16-Aug	14-Sep	26-Sep	
1995	14-Aug	7-Sep	1-Oct	
1996	20-Aug	18-Sep	23-Sep	
1997	12-Aug	11-Sep	23-Sep	23-Sep
1998	11-Aug	15-Sep	30-Sep	22-Sep
1999	24-Aug	8-Sep	27-Sep	21-Sep
2000	7-Aug	20-Sep	19-Sep	19-Sep
2001	14-Aug	13-Sep	25-Sep	18-Sep
2002	12-Aug	11-Sep	23-Sep	24-Sep
2003	11-Aug	14-Sep	28-Sep	23-Sep
2004	17-Aug	12-Sep	27-Sep	21-Sep
2005	15-Aug	15-Sep	27-Sep	20-Sep
2006	15-Aug	14-Sep	26-Sep	19-Sep
2007	14-Aug	12-Sep	25-Sep	25-Sep
2008	11-Aug	12-Sep	23-Sep	23-Sep
2009	17-Aug	10-Sep	23-Sep	28-Sep
2010	17-Aug	12-Sep	21-Sep	21-Sep
2011	23-Aug	8-Sep	21-Sep	20-Sep
2012	21-Aug	11-Sep	24-Sep	25-Sep
2013	19-Aug	11-Sep	25-Sep	23-Sep
2014	19-Aug	18-Sep	29-Sep	24-Sep
2015	20-Aug	17-Sep	28-Sep	23-Sep
2016	16-Aug	16-Sep	27-Sep	20-Sep
2017 ²	16-Aug		26-Sep	19-Sep
2018	15-Aug	20-Sep	1-Oct	25-Sep
2019	15-Aug	9-Sep	1-Oct	24-Sep
2020	31-Aug	23-Sep	29-Sep	22-Sep
Mean	15-Aug	12-Sep	25-Sep	22-Sep

1. Approximately one-half of the redds in the system were counted by this date and one-half were counted after this date. For the CESRF, approximately one-half of the total broodstock were spawned by this date and one-half were spawned after this date.
2. Spawner surveys impacted by fires; especially in the Naches system.

Redd Counts and Distribution

Table 31. Yakima Basin spring Chinook redd count summary, 1981 – present.

Year	Upper Yakima River System				Naches River System				
	Mainstem ¹	Elum	Teaway	Total	American	Naches ¹	Bumping	Little Naches	Total
1981	237	57	0	294	72	64	20	16	172
1982	610	30	0	640	11	25	6	12	54
1983	387	15	0	402	36	27	11	9	83
1984	677	31	0	708	72	81	26	41	220
1985	795	153	3	951	141	168	74	44	427
1986	1,716	77	0	1,793	464	543	196	110	1,313
1987	968	75	0	1,043	222	281	133	41	677
1988	369	74	0	443	187	145	111	47	490
1989	770	192	6	968	187	200	101	53	541
1990	727	46	0	773	143	159	111	51	464
1991	568	62	0	630	170	161	84	45	460
1992	1,082	164	0	1,246	120	155	99	51	425
1993	550	105	1	656	214	189	88	63	554
1994	226	64	0	290	89	93	70	20	272
1995	105	12	0	117	46	25	27	6	104
1996	711	100	3	814	28	102	29	25	184
1997	364	56	0	420	111	108	72	48	339
1998	123	24	1	148	149	104	54	23	330
1999	199	24	1	224	27	95	39	25	186
2000	3,349	466	21	3,836	54	483	278	73	888
2001	2,910	374	21	3,305	392	436	257	107	1,192
2002	2,441	275	110	2,826	366	226	262	89	943
2003	772	87	31	890	430	228	216	61	935
2004	2,985	330	129	3,444	91	348	205	75	719
2005	1,717	287	15	2,019	140	203	163	68	574
2006	1,092	100	58	1,250	136	163	115	33	447
2007	665	51	10	726	166	60	60	27	313
2008	1,191	137	47	1,375	158	165	102	70	495
2009	1,349	197	33	1,579	92	159	163	68	482
2010	2,199	219	253	2,671	173	171	168	40	552
2011	1,663	171	64	1,898	212	145	175	48	580
2012	1,276	125	69	1,470	337	196	189	89	811
2013	552	85	34	671	170	66	85	55	376
2014	962	138	53	1,153	129	65	158	27	379
2015	1,258	39	24	1,321	239	177	152	46	614
2016	512	83	22	617	149	106	74	37	366
2017	402	118	23	543	123	84	56	30	293
2018	339	13	0	352	27	56	44	1	128
2019	185	44	9	238	21	1	2	7	31
2020	189	44	8	241	44	25	71	6	146
Mean	980	119	26	1,125	153	157	109	45	464

¹ Including minor tributaries.

Homing

A team from NOAA fisheries conducted studies to determine the spatial and temporal patterns of homing and spawning by wild and hatchery-reared salmon released from CESRF facilities from 2001 to 2010. These studies collected GPS information on each redd and carcass recovered within a survey reach. Carcass surveys were conducted annually in late-September to early October by NOAA personnel in cooperation with Yakama Nation survey crews over five different reaches of the upper Yakima River and recorded the location of each redd flagged and carcass recovered. For each carcass sex, hatchery/wild, male status (full adult, jack, mini-jack), and CWT location was recorded. Data collected on the body location of CWTs allowed the identification of the release site of some fish. While these studies were not designed to comprehensively map carcasses and redds in all spawning reaches in the upper watershed, preliminary data indicate that fish from the Easton, Jack Creek, and Clark Flat acclimation facilities had distinct spawner distributions. A more complete description of this project is available from NOAA fisheries and in this publication:

Dittman, A. H., D. May, D. A. Larsen, M. L. Moser, M. Johnston, and D. Fast. 2010. Homing and spawning site selection by supplemented hatchery- and natural-origin Yakima River spring Chinook salmon. *Transactions of the American Fisheries Society* 139:1014-1028.

Straying

The regional PTAGIS (PIT tag) and RMIS (CWT) databases were queried in January 2021 to determine the number of CESRF releases not returning to the Yakima River Basin (RMIS CWT data are incomplete for the most recent years). For adult (age-3, -4, or -5) PIT tagged fish, a stray is defined as detection at an out-of-basin facility in the Snake (Ice Harbor or Lower Granite) or Upper Columbia (Priest Rapids, Rock Island, or Wells) without a subsequent detection at Prosser or Roza Dam. For coded-wire tagged fish, a stray is generally defined as a tag recovery in tributaries of the Columbia River upstream (and including the Snake River Basin) of its' confluence with the Yakima River. Marked (adipose fin clipped) fish are occasionally found during carcass surveys in the Naches River system. All marked fish observed in spawning ground carcass surveys in the Naches Basin are assumed to be CESRF fish and are used to estimate in-basin stray rates.

Table 32. Estimated number of PIT- and CWT-tagged CESRF fish not returning to the Yakima River Basin (strays), and marked fish sampled during spawner surveys in the Naches Basin, per number of returning fish, brood years 1997-present.

Brood Year	CESRF PIT-Tagged Fish			All CESRF Fish			CESRF Age-4 Fish		
	Roza Adult Returns	Adult Strays	Stray Rate	Yakima River Mth Return	CWT Strays	Stray Rate	Yak R. MthRtn	In-Basin Strays ¹	Stray Rate
1997	598	2	0.33%	8,670	1	0.01%	7,753		
1998	398	0	0.00%	9,782			7,939	1	0.01%
1999	23	0	0.00%	864			714		
2000	150	4	2.60%	4,819	2	0.04%	3,647	4	0.11%
2001	80	3	3.61%	1,251			845	2	0.24%
2002	97	5	4.90%	2,300			1,886	1	0.05%
2003	31	0	0.00%	932			800		
2004	125	1	0.79%	4,022	4	0.10%	3,101		
2005	142	0	0.00%	4,378			3,052		
2006	462	3	0.65%	9,114			5,812		
2007	240	1	0.41%	6,558	5	0.08%	5,174	1	0.02%
2008	215	0	0.00%	6,976			4,567	1	0.02%
2009	110	0	0.00%	3,181			2,663	1	0.04%
2010	207	5	2.36%	4,707	2	0.04%	3,183		
2011 ²	181	28	13.40%	3,607	16	0.44%	2,340		
2012 ²	69	13	15.85%	1,723	20	1.16%	1,492		
2013	152	4	2.56%	2,795	6	0.21%	1,993		
2014 ²	131	13	9.03%	2,463	4	0.16%	1,447		
2015 ²	57	2	3.39%	1,191	1	0.08%	877		
2016 ²	62	10	13.89%	1,058	6	0.57%	771		

¹ All marked fish observed in spawning ground carcass surveys in the Naches Basin are assumed to be CESRF fish.

² Water temperatures in the lower Yakima River were greater than 68° F for much of the late spring/summer migration since 2015 which likely caused many fish returning in recent years to seek cooler water in other parts of the Columbia Basin.

CESRF Spawning and Survival

As described earlier, a portion of natural- and hatchery-origin (NoR and HoR, respectively) returning adults are captured at Roza Dam during the adult migration and taken to the CESRF for broodstock and/or research purposes. Fish are held in adult holding ponds at the CESRF from capture in the spring and summer until spawning in September through early October. All mortalities during the holding period are documented by sex and origin. During the spawning period data are kept on the number of males and females of each origin used for spawning or other purposes. All females have samples taken that are later evaluated for presence of BKD-causative agents. Eggs from females with high BKD-presence indicators are generally excluded (see Female BKD Profiles). Once fertilized, eggs are placed in holding troughs until shock time. Dead eggs are then sorted and hand-counted. All live eggs are machine counted, sorted into two lots per female (treatment and control) and placed into incubation (heath) trays. Using hand counts of egg samples from a subsample of female egg lots, WDFW staff determined that machine counts are biased and that the best approximation of live egg counts is given by the following equation:

$$\left(\left(\frac{\text{no. eggs in subsample}}{\text{wt. of subsample}} * \text{total egg mass wt} \right) * 0.945 \right) - \text{dead eggs}$$

where

the first 3 parameters are from egg samples taken from females at spawn time, dead eggs are the number of dead or unfertilized eggs counted at shock time, and the 0.945 value is a correction factor from 1997 and 2000 WDFW studies.

Total egg take is calculated as the total number of live eggs, dead eggs, and all documented egg loss (e.g. spilled at spawn time, etc.). Heath trays are periodically sampled during incubation and dead fry are culled and counted. The number of live eggs less documented fry loss is the estimate of the number of fry ponded. Once fry are ponded, mortalities are counted and recorded daily during the rearing period. Fish are hand counted in the fall prior to their release as they are 100-percent marked. This hand-count less documented mortalities from marking through release is the estimate of smolts released. Survival statistics by origin and life-stage are given in Tables 33 and 34.

Table 33. Cle Elum Supplementation and Research Facility spawning and survival statistics (NoR brood only), 1997 - present.

Brood Year	Total Collected	Total Morts.	PreSpawn Survival	No. Fish Spawned ¹			% BKD Loss	Total Egg Take	Live Eggs	% Egg Loss ³	Fry Ponedged ⁴	Live-Egg-Fry Survival	Smolts Released	Fry-Smolt Survival	Live-Egg-Smolt Survival
				Males ²	Females	%									
1997	261	23	91.2%	106	132	2.6%	500,750	463,948	7.3%	413,211	98.5%	386,048	93.4%	91.9%	
1998	408	70	82.8%	140	198	1.4%	739,802	664,125	10.2%	627,481	98.7%	589,648	94.0%	92.7%	
1999	738 ⁵	24	96.7%	213	222	2.7%	818,816	777,984	5.0%	781,872	97.3%	758,789	97.0%	94.5%	
2000	567	61	89.2%	170	278	9.2%	916,292	851,128	7.1%	870,328	97.3%	834,285	95.9%	93.4%	
2001	595	171	71.3%	145	223	53.2%	341,648	316,254	7.4%	380,880	98.6%	370,236	97.2%	96.1%	
2002	629	89	85.9%	125	261	10.0%	919,776	817,841	11.1%	783,343	98.0%	749,067	95.6%	93.6%	
2003	441	54	87.8%	115	200	0.0%	856,574	787,933	8.0%	761,990	98.4%	735,959	96.6%	95.0%	
2004	597	70	88.3%	125	245	0.4%	873,815	806,375	7.7%	776,941	97.8%	691,109 ⁶	89.0%	87.0%	
2005	526	57	89.2%	136	241	0.0%	907,199	835,890	7.9%	796,559	98.1%	769,484	96.6%	94.7%	
2006	519	45	91.3%	122	239	1.7%	772,357	703,657	8.9%	631,691	97.3%	574,361 ⁷	90.9%	88.3%	
2007	473	49	89.6%	149	216	0.9%	798,729	760,189	4.8%	713,814	98.9%	676,602	94.8%	93.7%	
2008	480	38	92.1%	151	253	2.0%	915,563	832,938	9.0%	809,862	99.0%	752,109 ⁸	97.3%	96.3%	
2009	486	57	88.3%	142	219	1.4%	850,404	848,339	0.2%	770,706	98.2%	744,170	96.6%	94.6%	
2010	483	20	95.9%	102	193	0.5%	787,953	753,464	4.4%	726,325	98.9%	702,751	96.8%	95.6%	
2011	455	28	93.8%	103	197	0.0%	798,229	765,221	4.1%	721,197	98.1%	684,481	94.9%	93.0%	
2012	363	14	96.1%	111	209	0.0%	819,775	788,605	3.8%	737,705	98.2%	712,036	96.5%	94.7%	
2013	385	15	96.1%	153	179	0.6%	683,484	658,796	3.6%	613,493	98.9%	575,156	93.8%	92.6%	
2014	384	39	89.8%	133	188	0.0%	679,374	639,989	5.8%	636,092	96.5%	599,908	94.3%	91.1%	
2015	436	116	73.4%	128	182	0.5%	654,361	615,189	6.0%	613,796	97.0%	594,736	96.9%	94.1%	
2016	394	57	85.5%	142	173	0.0%	687,218	652,110	5.1%	593,514	96.2%	588,139	99.1%	95.2%	
2017	396	27	93.2%	152	193	2.1%	707,232	671,605	5.0%	642,836	95.7%	634,390	98.7%	94.5%	
2018	305	6	98.0%	122	166	0.0%	565,221	534,753	5.4%	515,596	98.2%	498,011	96.6%	94.8%	
2019	313	25	92.0%	103	174	2.3%	541,760	504,630	6.9%	482,177	94.7%	450,377	93.4%	88.5%	
2020	423	29	93.1%	144	230	1.7%	708,208	676,954	4.4%	674,954	97.5%				
Mean	461	49	89.6%	135	209	3.9%	743,552	696,996	6.2%	669,848	97.7%	637,907	95.5%	93.3%	

1. Total collected minus total mortalities does not equal total spawned. This is because some fish are used in the spawning channel, some have been released back to the river, and some have not been used.
2. Includes jacks.
3. All documented egg loss at spawn time plus dead eggs counted at shock divided by the estimated total egg take.
4. Based on physical counts at mark time and all documented rearing mortality from ponding to release, except for BY2013 it is live eggs (est.) minus fry loss.
5. Approximately one-half of these were jacks, many of which were not used in spawning.
6. Approximately 45,000 smolts lost at Jack Creek due to frozen equipment in February, 2006.
7. EWOS feed treatment had high mortality and was discontinued in May 2007; resulted in lower survival to release.
8. Approximately 36,000 NoR (Table 33) and 12,000 HoR (Table 34) fish were culled in July 2009 to reduce pond densities; these fish were added back in to fry-smolt and live-egg-smolt survival calculations.

Table 34. Cle Elum Supplementation and Research Facility spawning and survival statistics (HoR brood only), 2002 - present.

Brood Year	Total Collected	Total Morts.	PreSpawn Survival	No. Fish Spawmed ¹		% BKD Loss	Total Egg Take ⁹	Live Eggs ¹⁰	% Egg Loss ³	Fry Ponedged ⁴	Live-Egg-Fry Survival	Smolts Released	Fry-Smolt Survival	Live-Egg-Smolt Survival
				Males ²	Females									
2002	201	22	89.1%	26	72	4.2%	258,226	100,011	7.8%	91,300	98.2%	87,837	96.2%	94.4%
2003	143	12	91.6%	30	51	0.0%	219,901	83,128	7.3%	91,204	98.8%	88,733	97.3%	96.1%
2004	126	19	84.9%	22	49	0.0%	187,406	94,659	5.9%	100,567	98.3%	94,339	93.8%	92.2%
2005	109	6	94.5%	26	45	0.0%	168,160	89,066	12.2%	92,903	98.1%	90,518	97.4%	95.6%
2006	136	21	84.6%	28	41	2.4%	112,576	80,121	8.6%	74,735	97.6%	68,434	91.6%	89.4%
2007	110	15	86.4%	26	35	0.0%	125,755	90,162	3.2%	96,912	99.2%	94,663	97.7%	96.9%
2008	194	10	94.8%	51	67	1.5%	247,503	106,122	5.1%	111,797	98.9%	97,196	97.4%	96.4%
2009	164	24	85.4%	30	38	0.0%	148,593	91,994	0.8%	91,221	98.3%	88,771	97.3%	95.6%
2010	162	9	94.4%	29	55	1.8%	215,814	94,925	8.4%	96,144	97.9%	92,030	95.7%	93.7%
2011	166	7	95.8%	28	49	0.0%	188,075	89,107	4.5%	88,852	98.4%	84,701	95.3%	93.8%
2012	140	8	94.3%	29	42	0.0%	148,932	95,438	2.0%	94,031	98.8%	90,680	96.4%	95.3%
2013	186	5	97.3%	38	43	0.0%	155,383	80,534	2.9%	75,842	98.2%	71,599	94.4%	92.7%
2014	86	11	87.2%	21	29	0.0%	104,121	74,843	1.6%	91,702	97.2%	85,322	93.0%	90.4%
2015	61	23	62.3%	15	22	13.6%	66,238	64,646	2.4%	62,625	96.9%	60,211	96.1%	93.1%
2016	114	25	78.1%	33	35	0.0%	129,355	121,466	6.1%	85,910	95.8%	81,069	94.4%	90.4%
2017	127	8	93.7%	46	55	0.0%	195,070	187,173	4.0%	88,905	97.9%	76,279	85.8%	84.0%
2018	101	6	94.1%	33	54	0.0%	179,083	172,211	3.8%	150,126 ¹¹	96.1%	144,409	96.2%	92.4%
2019	126	12	90.5%	43	46	0.0%	128,677	115,667	10.1%	120,071 ¹¹	92.6%	100,021	83.3%	77.1%
2020	131	18	86.3%	43	50	4.0%	133,970	124,494	7.1%	97,324	97.3%			
Mean	136	14	88.7%	31	46	1.5%	163,834	154,468	5.5%	94,851	97.6%	88,712	94.4%	92.2%

Continued from footnotes for Table 33 above.

9. Table 34 -- From 2002 to present this is the estimated total egg take from all HxH crosses. Due to the large surplus of eggs over the approximately 100K needed for the HxH line, many surplus fry were planted in nearby land-locked lakes and some surplus eggs were destroyed.

10. Table 34 -- For only those HxH fish which were actually ponded.

11. The number of segregated, hatchery-control line brood raceways was increased from 2 to 4 for this brood due to overall brood shortages.

Female BKD Profiles

Adults used for spawning and their progeny are tested for a variety of pathogens accepted as important in salmonid culture (USFWS Inspection Manual, 2003), on a population or "lot" basis. At the CESRF, and in the Columbia Basin it has been accepted that the most significant fish pathogen for spring Chinook is *Renibacterium salmoninarum*, the causative agent of Bacterial Kidney Disease (BKD). All adult females and 30-60 juveniles from each acclimation pond are individually tested for levels of *Renibacterium salmoninarum* using ELISA (Enzyme linked Immuno-sorbant Assay). ELISA data are reported annually to CESRF and YKFP staff for management purposes, eventual data entry and comparisons of ponds and rearing parameters. To date, no significant occurrences of other pathogens have been observed. Periodic field exams for external parasites and any signs of disease are performed on an "as needed" basis. Facility staff have been trained to recognize early signs of behavior changes or diseases and would report any abnormalities to the USFWS, Olympia Fish Health Center for further diagnostic work.

Adult females are ranked from 0 to 13 based on the relative amounts of BKD in the tissue samples of the tested fish. All BKD ranks below 5 are considered low risk for transferring significant BKD organisms through the egg to cause significant disease in progeny receiving proper care. The progeny of adults with BKD rank 6 are considered to be moderate risk and those with BKD rank 7 or greater are considered to be high risk. Given these data, the CESRF chose to rear only the progeny of females with a BKD rank of 6 or less through brood year 2001. Beginning with brood year 2002, the progeny of fish with BKD rank 6 (moderate risk) or greater (high risk) have not been used for production purposes at the CESRF. For additional information, see Appendix B.




Figure 4. Proportion of wild/natural females spawned at CESRF by BKD rank, 1997 – present.

Fecundity

Fish collected at Roza Dam are taken to the CESRF for spawning and/or research purposes. Egg loss due to spill or other reasons at spawn time is documented. When eggs are shocked, unfertilized (dead) eggs are hand-counted and remaining eggs are machine counted. Due to error associated with machine counts, average fecundity is calculated using spawn-time egg sample data (see discussion above under CESRF Spawning and Survival) and adding in documented egg loss for all females divided by the number of females (N) in the sample.

Table 35. Mean fecundity by age of adult females (BKD rank < 6) spawned at CESRF, 1997-present.

Brood Year	Wild/Natural (SN)						CESRF (HC)					
	Age-3		Age-4		Age-5		Age-3		Age-4		Age-5	
N	Fecundity	N	Fecundity	N	Fecundity	N	Fecundity	N	Fecundity	N	Fecundity	
1997			105	3,842.0	4	4,069.9						
1998	2 ¹	3,908.9	161	3,730.3	15	4,322.5						
1999	3 ¹	4,470.4	183	3,968.1	14	4,448.6						
2000			224	3,876.5	2	5,737.9						
2001			72	3,966.9	9	4,991.2			18	4,178.9		
2002	1	1,038.0	205	3,934.7	7	4,329.4			60	3,820.0	1	4,449.0
2003			163	4,160.2	31	5,092.8			30	3,584.1	19	5,459.9
2004			224	3,555.4	2	4,508.3			42	3,827.2		
2005	1	1,769.0	218	3,815.5	5	4,675.1			38	3,723.9	5	4,014.7
2006			196	3,396.4	24	4,338.9			36	3,087.3		
2007			178	3,658.3	24	4,403.3			33	3,545.2	2	4,381.9
2008			207	3,814.0	10	4,139.9			58	3,898.0		
2009	1	2,498.2	195	4,018.9	6	4,897.1			34	3,920.3		
2010			185	4,103.0					54	3,996.6		
2011	1 ¹	3,853.1	179	4,000.1	4	5,692.1			41	3,843.3	2	4,098.2
2012			186	3,901.0	5	4,982.8			41	3,537.4	1	3,900.5
2013			159	3,760.3	6	5,068.0			36	3,498.7	2	4,955.3
2014			171	3,889.4	4	4,599.5			25	3,627.1	1	5,335.8
2015			166	3,963.0	2	5,249.3			14	3,975.1	1	3,793.3
2016			159	3,969.1	7	4,959.4			34	3,675.9	1	4,375.5
2017	2	2,150.6	161	4,013.8	1	3,805.5	1	1,645.0	53	3,609.1		
2018			130	3,452.4	6	3,643.9			49	3,348.3		
2019	1	1,500.8	129	3,573.2	2	3,519.3	2	1,520.5	40	3,466.3	1	3,204.0
2020	2	1,899.4	147	3,418.8					33	3,423.3		
Mean				3,824.2		4,612.5				3,678.7		4,360.7

1. Given their length and fecundity, these fish may have been incorrectly aged.

Juvenile Salmon Evaluation

Food Conversion Efficiency

At the end of each month that fish are in the rearing ponds at the CESRF or the acclimation sites, a sample of fish are weighed and measured to estimate growth. These data, in addition to monthly mortality and pond feed data are entered into the juvenile growth and survival tracking database. Hatchery managers monitor food conversion (total pounds fed during a month divided by the total pounds gained by the fish) to track how well fish are converting feed into body mass and to evaluate the amount of feed that needs to be provided on a monthly basis. Average monthly food conversion and growth statistics for the CESRF facilities by brood year are provided in the following tables and figures.

Table 36. Mean food conversion (lbs fed/lbs gained) of CESRF juveniles by brood year and growth month, 1997 – present.

Brood Year	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar
1997	2.2		1.1	0.8	1.2	0.8	1.5	1.5		1.9		5.3
1998		1.0	0.9	1.0	0.9	0.8	2.4	1.4	2.1	-0.3	1.0	1.2
1999		1.0	1.1	1.1	1.2	1.5	1.8	1.0		-0.5	0.3	1.7
2000	0.8	0.8	1.0	1.5	1.2	1.4	2.2	2.0	1.6	2.1	2.5	2.4
2001	1.1	1.1	2.6	1.1	1.3	1.2	1.6	2.0	2.3	2.5	2.8	0.9
2002	0.9	1.0	1.4	1.2	1.4	1.1	1.5	2.2	4.0	-1.4	2.9	1.0
2003	0.6	1.0	0.9	1.4	1.2	1.2	4.6	0.7	0.9	-0.2	1.8	1.0
2004	0.9	1.0	1.2	1.6	2.4	1.2	1.7	2.0	2.8	0.9	-2.6	1.1
2005	0.8	0.7	1.3	1.0	1.3	1.2	1.5	-0.8	0.4	-0.4	2.2	
2006	0.8	0.7	0.6	0.9	0.8	1.0	1.6	-1.0		-2.6	0.6	0.6
2007	0.7	0.7	0.9	0.9	1.0	0.8	2.2	-1.6	1.9	2.0	0.7	0.9
2008	0.5	0.6	0.9	0.9	1.0		0.8	1.7	-1.1	0.9	0.9	0.6
2009	0.5	1.2	1.0	0.7	1.1	1.0	1.5	4.1	0.6	-2.8	0.8	0.9
2010	0.6	0.8	1.3	0.8	0.8	1.8	2.8	1.3		0.8	0.8	0.7
2011	0.9	0.6	0.8	0.7	1.1	0.9		0.7		0.6	0.9	1.0
2012	0.8	1.4	1.1	0.8	1.3	1.4	1.0	1.1		1.0	3.1	1.2
2013	0.6	0.9	0.7	0.9	1.0	1.1	2.7	1.4		0.4	0.8	2.5
2014	0.5	2.2	0.7	1.0	2.4	0.7	4.3	0.5		1.7	0.9	0.8
2015	0.8	0.9	0.8	1.0	1.3	0.9	-1.8	0.7	-0.8	1.0	0.5	0.9
2016	0.6	0.9	0.8	1.0	1.1	1.1	2.1	1.8	1.0	0.6	0.4	0.8
2017	0.8	0.8	0.9	0.9	1.7	0.8	2.1	2.9	3.8	0.4	0.1	0.6
2018	0.7	0.8	0.9	0.9	1.3	1.1		0.9		0.6	1.3	1.6
2019	0.8	1.7	1.1	0.8	1.3	1.5	1.1	1.6	3.3	0.6	1.5	0.9
Mean	0.8	0.9	1.0	1.0	1.3	1.1	1.9	1.2	1.6	0.4	1.1	1.1

Length and Weight Growth Profiles




Figure 5. Mean fork length (cm) of CESRF juveniles by brood year and growth month, 1997 - present.





Figure 6. Mean Weight (fish/lb) of CESRF juveniles by brood year and growth month, 1997 - present.

Juvenile Fish Health Profile

Approximately 50-100 juveniles were sacrificed for juvenile fish health samples in the spring (usually in March) of their release year. Tissue samples from these fish were processed at USFWS laboratories in Olympia, Washington for presence of bacterial kidney disease (BKD) using enzyme-linked immunosorbent assay (ELISA) tests (see Female BKD Profiles and Appendix B for additional discussion). Fish were ranked high, moderate, or low (risk) based on the relative amounts of BKD in the tissue samples of the tested fish. These relative risk levels assume a good fish culture and rearing environment (i.e., water temperature and flows, nutrition, densities, etc. all must be conducive to good fish health). As indicated in Figure 7, juvenile fish released from the CESRF are largely in the low risk category for all brood years sampled to date. Due to budget issues and the low incidence observed over twenty years of testing, the USFWS discontinued testing of juveniles beginning with brood year 2017.

Figure 7. ELISA-risk profile of CESRF juveniles by brood year, 1997 – present (data source: USFWS).



Incidence of Precocialism

For brood years 2002-2004, the YKFP tested two different feeding regimes to determine whether a slowed-growth regime reduces the incidence of precocialism without a reduction in post-release survival. The two growth regimes tested were a normal (High) growth regime resulting in fish which were about 30/pound at release and a slowed growth regime (Low) resulting in fish which were about 45/pound at release. As a critical part of this study, a team from NOAA Fisheries conducted research to characterize the physiology and development of wild and hatchery-reared spring Chinook salmon in the Yakima River Basin. While precocious male maturation is a normal life-history strategy, the hatchery environment may be potentiating this developmental pathway beyond natural levels resulting in potential loss of anadromous adults, skewing of sex ratios, and negative genetic and ecological impacts on wild populations. Previous studies have indicated that age of maturation is significantly influenced by endogenous energy stores and growth rate at specific times of the year. These studies will help direct rearing strategies at the CESRF to allow production of hatchery fish with physiological and life-history attributes that are more similar to their wild cohorts.

Relevant Publications:

- Larsen, D. A., B. R. Beckman, K. A. Cooper, D. Barrett, M. Johnston, P. Swanson, and W. W. Dickhoff. 2004. Assessment of High Rates of Precocious Male Maturation in a Spring Chinook Salmon Supplementation Hatchery Program. *Transactions of the American Fisheries Society* 133:98-120.
- Beckman, B.R. and Larsen D.A. 2005. Upstream Migration of Minijack (Age-2) Chinook Salmon in the Columbia River: Behavior, Abundance, Distribution, and Origin. *Transactions of the American Fisheries Society* 134:1520–1541.
- Larsen, D.A., B.R. Beckman, C.R. Strom, P.J. Parkins, K.A. Cooper, D.E. Fast, W.W. Dickhoff. 2006. Growth Modulation Alters the Incidence of Early Male Maturation and Physiological Development of Hatchery-reared Spring Chinook Salmon: a Comparison with Wild Fish. *Transactions of the American Fisheries Society* 135:1017-1032.
- Pearsons, T.N., C.L. Johnson, B.B. James, and G.M. Temple. 2009. Abundance and Distribution of Precociously Mature Male Spring Chinook Salmon of Hatchery and Natural Origin in the Yakima River. *North American Journal of Fisheries Management* 29:778-790.
- Larsen, D.A., B.R. Beckman, and K.A. Cooper. 2010. Examining the Conflict between Smolting and Precocious Male Maturation in Spring (Stream-Type) Chinook Salmon. *Transactions of the American Fisheries Society* 139: 564-578.
- Larsen, D.A., D.L. Harstad, C.R. Strom, M.V. Johnston, C.M. Knudsen, D.E. Fast, T.N. Pearsons, and B.R. Beckman. 2013. Early Life History Variation in Hatchery- and Natural-Origin Spring Chinook Salmon in the Yakima River, Washington. *Transactions of the American Fisheries Society* 142:2, 540-555.

CESRF Smolt Releases

The number of release groups and total number of fish released diverged from facility goals in some years. In brood year 1997, the Jack Creek acclimation facility was not yet complete and project policy and technical teams purposely decided to under-collect brood stock to allow a methodical testing of the new facility's operations with less risk to live fish, which resulted in the stocking of only 10 of the 18 raceways. In brood year 1998, the project did not meet facility release goals due to a biological specification that no more than 50% of returning wild fish be taken for brood stock. As a result only 16 raceways were stocked with progeny of the 1998 brood. In the same year, raceway 4 at the Jack Creek acclimation site suffered mechanical failures causing loss of flow and reduced oxygen levels and resulted in the loss of approximately one-half the fish in this raceway prior to release. In the drought year of 2001, a large number of returning adults presented with high enzyme-linked immunosorbent assay (ELISA) levels of *Renibacterium salmoninarum*, the causative agent of bacterial kidney disease (BKD). The progeny of these females were purposely destroyed. As a result, only nine raceways were stocked with fish. The project decided to use the fish from an odd raceway for a predator avoidance training sub-experiment (these fish were subsequently acclimated and released from the Easton acclimation site).

Table 37. CESRF total releases by brood year, treatment, and acclimation site.

Brood Year	Acclimation Site					Total
	Control ¹	Treatment ²	CFJ	ESJ	JCJ	
1997	207,437	178,611	229,290	156,758		386,048
1998 ³	284,673	305,010	221,460	230,860	137,363	589,683
1999	384,563	374,226	232,563	269,502	256,724	758,789
2000	424,554	409,731	285,954	263,061	285,270	834,285
2001 ⁴	183,963	186,273	80,782	39,106	250,348	370,236
2002	420,764	416,140	266,563	290,552	279,789	836,904
2003	414,175	410,517	273,377	267,711	283,604	824,692
2004 ⁵	378,740	406,708	280,598	273,440	231,410	785,448
2005	431,536	428,466	287,127	281,150	291,725	860,002
2006	351,063	291,732	209,575	217,932	215,288	642,795
2007	387,055	384,210	265,907	254,540	250,818	771,265
2008	421,290	428,015	280,253	287,857	281,195	849,305
2009	418,314	414,627	279,123	281,395	272,423	832,941
2010	395,455	399,326	264,420	264,362	265,999	794,781
2011	382,195	386,987	255,290	248,454	265,438	769,182
2012	401,059	401,657	256,732	276,210	269,774	802,716
2013	No Experiment		215,933	214,745	216,077	646,755
2014	337,548	347,682	232,440	226,257	226,533	685,230
2015	331,316	323,631	208,239	218,225	228,483	654,947
2016	339,816	329,392	230,490	218,676	220,042	669,208
2017	351,656	359,013	244,236	233,449	232,984	710,669
2018	322,219	320,201	213,833	206,619	221,968	642,420
2019	270,242	280,156	153,575	193,042	203,781	550,398
Mean	356,347	353,741	237,729	235,387	244,865	707,335

Table 38. CESRF average pond densities at release by brood year, treatment, and acclimation site.

Brood Year	Treatment		Acclimation Site		
	Control ¹	Treatment ²	CFJ	ESJ	JCJ
1997	41,487	35,722	38,215	39,190	
1998 ³	35,584	38,126	36,910	38,477	34,341
1999	42,729	41,581	38,761	44,917	42,787
2000	47,173	45,526	47,659	43,844	47,545
2001 ⁴	41,116	41,667	40,391	6,518	41,725
2002	46,752	46,238	44,427	48,425	46,632
2003	46,019	45,613	45,563	44,619	47,267
2004 ⁵	42,082	45,190	46,766	45,573	38,568
2005	47,948	47,607	47,855	46,858	48,621
2006	39,007	32,415	34,929	36,322	35,881
2007	43,006	42,690	44,318	42,423	41,803
2008	46,810	47,557	46,709	47,976	46,866
2009	46,479	46,070	46,521	46,899	45,404
2010	43,939	44,370	44,070	44,060	44,333
2011	42,466	42,999	42,548	41,409	44,240
2012	44,562	44,629	42,789	46,035	44,962
2013	No Experiment		35,989	35,791	36,013
2014	37,505	38,631	38,740	37,710	37,756
2015	36,813	35,959	34,707	36,371	38,081
2016	37,757	36,599	38,415	36,446	36,674
2017	39,073	39,890	40,706	38,908	38,831
2018	35,802	35,578	35,639	34,437	36,995
2019	30,027	31,128	25,596	32,174	33,964
Mean	41,552	41,172	40,792	39,799	41,331

1. Brood years 1997-2001: Optimum Conventional Treatment (OCT). Brood Years 2002-2004: Normal (High) growth. Brood Years 2005-2012: Normal feed at Cle Elum or accl. sites.
2. Brood years 1997-2001: Semi-natural Treatment (SNT). Brood Years 2002-2004: Slowed (Low) growth. Brood Year 2005, 2007-2012: saltwater transition feed at accl. Sites; BY2014-present: BioPRO vs BioVIT diet. Brood Year 2006: EWS diet at CESRF through May 3, 2007.
3. At the Jack Creek acclimation site only 4 of 6 raceways were stocked, and raceway 4 suffered mechanical failures resulting in the loss of about 20,000 OCT (control) fish.
4. High BKD incidence in adult broodstock reduced production to just 9 ponds (Clark Flat 1-2, Jack Creek, and Easton). Easton ponds were used for predator avoidance trained (PAT) fish and a single Cle Elum pond was spread between 6 ponds at Easton with crowders used to simulate pond densities for fish at other acclimation sites. These releases were excluded from mean pond density calculations by treatment.
5. At the Jack Creek acclimation site raceway 3 suffered mechanical failures resulting in the loss of about 45,000 high-growth (control) fish.

Mean length and weight at release by brood year are shown in Figures 5 and 6 under Juvenile Salmon Evaluation, length and weight growth profiles. Mark information and volitional release dates are given in Appendix A.

Smolt Outmigration Timing

The Chandler Juvenile Monitoring Facility (CJMF) located on the fish bypass facility of Chandler Canal at Prosser Dam (Rkm 75.6; Figure 1) serves as the cornerstone facility for estimating smolt production in the Yakima Basin for several species and stocks of salmonids.

Daily species counts in the livebox at the CJMF are expanded by the canal entrainment, canal survival, and sub-sampling rates in order to estimate daily passage at Prosser Dam (Pandit 2020). Expansion techniques for deriving Chandler smolt passage estimates are continually being reviewed and revised to incorporate new information. A subset of fish passing through the CJMF is sampled for presence of internal (CWT or PIT) or external (fin-clip) marks. All fish with marks are assumed to be of hatchery origin; otherwise, fish are presumed to be of natural origin.




Figure 8. Mean flow approaching Prosser Dam versus mean estimated smolt passage at Prosser of aggregate wild/natural and CESRF spring Chinook for outmigration years 1999-2020.

Smolt-to-Smolt Survival

OCT-SNT Treatment (Brood Years 1997-2001, Migration Years 1999-2003)

Results of this experiment have been published:

Fast, D. E., D. Neeley, D.T. Lind, M. V. Johnston, C.R. Strom, W. J. Bosch, C. M. Knudsen, S. L. Schroder, and B.D. Watson. 2008. Survival Comparison of Spring Chinook Salmon Reared in a Production Hatchery under Optimum Conventional and Seminatural Conditions. *Transactions of the American Fisheries Society* 137:1507–1518.

Abstract — We found insufficient evidence to conclude that seminatural treatment (SNT; i.e., rearing in camouflage-painted raceways with surface and underwater structures and underwater feeders) of juvenile Chinook salmon *Oncorhynchus tshawytscha* resulted in higher survival

indices than did optimum conventional treatment (OCT; i.e., rearing in concrete raceways with surface feeding) for the specific treatments and environmental conditions tested. We reared spring Chinook salmon from fry to smolt in paired raceways under the SNT and OCT rearing treatments for five consecutive years. For four to nine SNT and OCT raceway pairs annually, we used passive integrated transponder, coded wire, and visual implant elastomer tags to compare survival indices for juvenile fish from release at three different acclimation sites 340–400 km downstream to passage at McNary Dam on the Columbia River, and for adults from release to adult return to Roza Dam in the upper Yakima basin. The observed differences in juvenile and adult survival between the SNT and OCT fish were either statistically insignificant, conflicting in their statistical significance, or explained by significant differences in the presence of the causative agents of bacterial kidney disease in juvenile fish at release.

High-Low Growth Treatment (Brood Years 2002-04, Migration Years 2004-2006)

Two early-rearing nutritional regimes were tested using hatchery-reared Yakima Upper spring Chinook for brood years 2002 through 2004. A low nutrition-feeding rate (low treatment or low) was administered at the Cle Elum Hatchery through early rearing to determine whether that treatment would reduce the proportion of precocials produced compared to a conventional feeding rate during early rearing. The conventional feeding rate, which served as a control treatment, is referred to here as a high nutrition-feeding rate (high treatment or high). Feed was administered at a rate of 10 grams/fish for the low treatment and 15 grams/fish for the high treatment through mid-October, after which sufficient feed was administered to both sets of treated fish to meet their feeding demands. The treatments were allocated within pairs of raceways (blocks), there being a total of nine pairs. The Low nutritional feed (Low) had a significantly lower release-to-McNary survival than did the High nutritional feed (High), respective survivals being 18.1% and 21.2% ($P < 0.0001$; D. Neeley, Appendix B of [2008 annual report](#)). The Low survival to McNary was consistently lower than the High at all sites in all years. Low-treated fish were smaller fish at the time of release and had somewhat later McNary passage times than high-treated fish. See also:

Larsen, D.A., B.R. Beckman, C.R. Strom, P.J. Parkins, K.A. Cooper, D.E. Fast, W.W. Dickhoff. 2006. Growth Modulation Alters the Incidence of Early Male Maturation and Physiological Development of Hatchery-reared Spring Chinook Salmon: a Comparison with Wild Fish. *Transactions of the American Fisheries Society* 135:1017-1032.

Larsen, D. A., D. L. Harstad, C. R. Strom, M. V. Johnston, C. M. Knudsen, D. E. Fast, T. N. Pearsons, and B. R. Beckman. 2013. Early life history variation in hatchery- and natural-origin spring Chinook Salmon in the Yakima River, Washington. *Transactions of the American Fisheries Society* 142:540–555.

Feed Treatments (Brood Years 2005, 2007- 2010; Migration Years 2007, 2009- 2018)

Prior to releases in 2007, and 2009- 2018, two feed treatments were allocated to raceways within adjacent raceway pairs. The feeds tested included Bio-Oregon's BioPro, BioVita, and BioTransfer diets (see <https://www.bio-oregon.com/>). The intent of the experiments was to determine whether any of the various feeds conferred any life-stage survival advantages. Preliminary analyses indicated no significant or substantial differences between the feeds when

averaged over years. See Appendix H of our [2015 annual report](#) and Appendix F of our [2019 annual report](#) for additional detail.

Control (Bio-Oregon) versus EWOS Feed Comparison (Brood Year 2006, Migration Year 2008)

This experimental design was similar to that for other studies described above with standard Bio-Oregon pellets fed to half of the rearing ponds and an EWOS (<https://www.cargill.com/animal-nutrition/brands/ewos>) diet fed to the other ponds. The different feed treatments only lasted about 6 weeks from the time of initial ponding as we found substantially higher mortalities for fish receiving the EWOS feed. From May 7, 2007 until these fish were released in 2008 all fish in this study received the Bio-Oregon diet. For the parameters of interest, we found no significant or substantial differences between the two feeding treatments (Appendix B of [2008 annual report](#)).

Smolt-to-Adult Survival

Calculation of smolt-to-adult survival rates for Yakima River spring Chinook is complicated by the following factors:

- 1) Downstream of the confluence of the Yakima and Naches rivers the three populations of spring Chinook (Upper Yakima, Naches, and American) are aggregated. A subsample of the aggregate wild/natural populations is PIT-tagged as part of the Chandler juvenile sampling operation but their origin is not known at the time of tagging. Through 2003, the primary purpose of this subsampling effort was to derive entrainment and canal survival estimates (see 2 below). Due to issues such as tag retention and population representation, adult detections of smolts PIT-tagged at Chandler cannot be used in any valid smolt-to-adult survival analyses.
- 2) Smolt accounting at Prosser is based on statistical expansion of Chandler smolt trap sampling data using available flow data and estimated Chandler entrainment rates. Chandler smolt passage estimates are prepared primarily for the purpose of comparing relative wild versus CESRF passage estimates and not for making survival comparisons. While these Chandler smolt passage estimates represent the best available data, there may be a relatively high degree of error associated with these estimates due to inherent complexities, assumptions, and uncertainties in the statistical expansion process. Therefore, these estimates are subject to revision. We are continuing to develop methods to subdivide the wild/natural outmigration into Upper Yakima, Naches, and American components based on DNA samples of juveniles taken at Chandler since 1998.
- 3) Installation of adult PIT detection equipment at all three ladders at Prosser Dam was not completed until the fall of 2005. Therefore, detection of upstream-migrating PIT-tagged adult spring Chinook at Prosser Dam was not possible for all returning fish until the spring of 2006. Periods of high flow may preclude use of automated detection gear so 100% detection of upstream migrants is not possible in all years.
- 4) Through 2006, detection of upstream-migrating PIT-tagged adult spring Chinook at Roza Dam occurred at an approximate 100% rate only for marked CESRF fish and wild/natural

fish taken for broodstock. The majority of wild/natural fish were passed directly back to the river without PIT interrogation.

- 5) For the 1997 brood (1999 out-migration), 400 Khz PIT-tags were used. Mainstem detection facilities were not configured to detect these tags at nearly the efficiency that they can detect the newer 134.2 kHz ISO tags. Although all marked adult fish are trapped and hand-wanded for PIT detections of adults at Roza Dam, the reliability of the 400kHz detection gear and problems with hand-sampling in general likely precluded a complete accounting of all 1997 brood PIT returns.
- 6) All CESRF fish are adipose-fin clipped and subjected to higher harvest rates than unmarked wild/natural fish in marine and Columbia River mark-selective fisheries. No adjustments have yet been made in the following tables to account for differential harvest rates in these mark-selective fisheries.
- 7) PIT tag retention is a factor in estimating survival rates (Knudsen et al. 2009). No attempt has been made to correct the data in the following tables for estimates of tag retention.
- 8) The ISAB has indicated that “more attention should be given to the apparent documentation that PIT-tagged fish do not survive as well as untagged fish. This point has major implications for all uses of PIT-tagged fish as surrogates for untagged fish.” Our data appear to corroborate this point (Tables 44-45). However, these data are not corrected for tag loss. If a fish loses its PIT tag after detection upon leaving the acclimation site, but before it returns as an adult to Roza Dam, it would be included only as a release in Table 45 and only as an adult return in Table 46. Knudsen et al. (2009) found that smolt-to-adult return rates (SARS) based on observed PIT tag recoveries were significantly underestimated by an average of 25% and that after correcting for tag loss, SARS of PIT-tagged fish were still 10% lower than SARS of non-PIT-tagged fish. Thus, the data in Table 45 under-represent “true” SARS for PIT-tagged fish and SARS for PIT-tagged and non-PIT-tagged fish are likely closer than those reported in Tables 44 and 45.
- 9) Due to issues relating to water permitting, size required for tagging, and allowing sufficient time for acclimation, CESRF juveniles are not allowed to migrate until at least March 15 of their smolt year. However, juvenile sampling observations at Roza Dam indicate that a substantial number of wild/natural juveniles migrate downstream during the summer, fall, and winter months prior to their smolt outmigration year (Figure 7). Comparison of SAR data for non-contemporaneously migrating juveniles may be invalid (see Copeland et al. 2015).

Given these complicating factors, Tables 39-45 present available smolt-to-adult survival data for Yakima River CESRF and wild/natural spring Chinook. Unfortunately, true “apples-to-apples” comparisons of CESRF and wild/natural smolt-to-adult survival rates are not possible from these tables due to complexities noted above. The reader is cautioned to correct these data for, or acknowledge the factors noted above prior to any use of these data.

Table 39. Estimated smolt passage at Chandler and smolt-to-adult return indices (Chandler smolt to Yakima R. mouth adult) for Yakima Basin wild/natural and CERSF-origin spring Chinook.

Brood Year	Smolt Migr. Year	Mean Flow ¹ at Prosser Dam	Estimated Smolt Passage at Chandler		CERSF smolt-to-smolt survival ³	Yakima R. Mouth Adult Returns ⁴		Smolt-to-Adult Return Index ⁴	
			Wild/Natural ²	CERSF Total		Wild/Natural ²	CERSF Total	Wild/Natural ²	CERSF Total
1982	1984	4134	381,857			6,753		1.8%	
1983	1985	3421	146,952			5,198		3.5%	
1984	1986	3887	227,932			3,932		1.7%	
1985	1987	3050	261,819			4,776		1.8%	
1986	1988	2454	271,316			4,518		1.7%	
1987	1989	4265	76,362			2,402		3.1%	
1988	1990	4141	140,218			5,746		4.1%	
1989	1991		109,002			2,597		2.4%	
1990	1992	1960	128,457			1,178		0.9%	
1991	1993	3397	92,912			544		0.6%	
1992	1994	1926	167,477			3,790		2.3%	
1993	1995	4882	172,375			3,202		1.9%	
1994	1996	6231	218,578			1,238		0.6%	
1995	1997	12608	52,028			1,995		3.8%	
1996	1998	5466	491,584			21,151		4.3%	
1997	1999	5925	584,016	187,669	48.6%	12,855	8,670	2.2%	4.6%
1998	2000 ⁵	4946	199,416	303,688	51.5%	8,240	9,782	4.1%	3.2%
1999	2001	1321	148,460	281,256	37.1%	1,764	864	1.2%	0.3%
2000	2002	5015	467,359	366,950	44.0%	11,434	4,819	2.4%	1.3%
2001	2003	3504	308,959	154,329	41.7%	8,597	1,251	2.8%	0.8%
2002	2004	2439	169,397	290,950	34.8%	3,743	2,557	2.2%	0.9%
2003	2005	1285	134,859	236,443	28.7%	2,746	1,020	2.0%	0.4%
2004	2006	5652	133,238	300,508	38.3%	2,802	4,482	2.1%	1.5%
2005	2007	4551	99,341	351,359	40.9%	4,295	5,004	4.3%	1.4%
2006	2008	4298	120,013	265,485	41.3%	6,004	10,577	5.0%	4.0%
2007	2009	5784	237,228	415,923	53.9%	7,952	7,604	3.4%	1.8%
2008	2010	3592	220,950	382,878	45.1%	7,385	8,036	3.3%	2.1%
2009	2011	9414	304,322	442,564	53.1%	3,766	3,606	1.2%	0.8%
2010	2012	8556	258,106	391,446	49.3%	6,602	5,592	2.6%	1.4%
2011	2013	4875	365,486	372,079	48.4%	7,343	4,160	2.0%	1.1%
2012	2014	4923	263,266	408,222	50.9%	3,969	1,932	1.5%	0.5%
2013	2015	1555	125,150	332,715	51.4%	3,415	3,139	2.7%	0.9%
2014	2016	5765	185,442	403,938	58.9%	1,800	2,865	1.0%	0.7%
2015	2017	7804	208,929	273,248	41.7%	1,171	1,319	0.6%	0.5%
2016	2018 ⁶	5652	131,489	290,644	43.4%	1,724 ⁶	1,220 ⁶	1.3% ⁶	0.4% ⁶
2017	2019 ⁶	3595	175,427	319,579	45.0%				
2018	2020 ⁶	2850	151,265	371,069	57.8%				

1. Mean flow (cfs) approaching Prosser Dam March 29-July 4 of juvenile migration year. No data available for migration year 1991. In high flow years (flows at or > 5000 cfs) operation of the Chandler smolt sampling facility may be precluded during portions of the outmigration. Data courtesy of [U.S. BOR hydromet.](#)
2. Aggregate of Upper Yakima, Naches, and American wild/natural populations.
3. Estimated smolt-to-smolt (release from upper Yakima River acclimation sites to Chandler) survival for CERSF juveniles.
4. Includes combined age-3 through age-5 returns. CERSF adult returns and smolt-to-adult survival values are understated relative to wild/natural values since these figures are not adjusted for differential harvest rates in mark selective fisheries in marine and lower Columbia River fisheries.
5. Available data were not sufficient to estimate juvenile flow-entrainment and passage of wild/natural fish.
6. Data for most recent years are preliminary; return data do not include age-5 adult fish.

Table 40. Estimated wild/natural smolt-to-adult return rates (SAR) based on adult detections of PIT tagged fish. Roza tagged smolts to Bonneville Dam adult returns. Footnotes follow Table 42.

Brood Year	Number Tagged	Wild/Natural smolts tagged at Roza				SAR ¹
		Adult Returns at Age ¹			Total	
		Age 3	Age 4	Age 5		
1997	310	0	1	0	1	0.32% ²
1998	6,209	15	171	14	200	3.22%
1999	2,179	2	8	0	10	0.46%
2000	8,718	1	51	1	53	0.61%
2001	7,804	9	52	3	64	0.82%
2002	3,931	2	46	4	52	1.32%
2003	1,733	0	6	1	7	0.40%
2004	2,333	1	8	1	10	0.43%
2005	1,200	0	8	0	8	0.67%
2006	1,675	12	33	2	47	2.81%
2007	3,795 ^a	6	47	2	55	1.45%
2008	105	0	1	0	1	0.95%
2009	2,087	0	3	1	4	0.19%
2010	2,647	4	22	1	27	1.02%
2011	2,473	1	9	1	11	0.44%
2012		No Releases				
2013	524	1	5	0	6	1.15%
2014	136	0	0	0	0	0.00%
2015	181	0	0	0	0	0.00%
2016	382	0	1		1	0.26%
2017	292	2				

a. Includes 1752 fish tagged and released in late August and early Sept.

Table 41. Estimated CESRF smolt-to-adult return rates (SAR) based on adult detections of PIT tagged fish. Roza tagged smolts to Bonneville Dam adult returns.

Brood Year	Number Tagged	CESRF smolts tagged at Roza				SAR ¹
		Adult Returns at Age ¹			Total	
		Age 3	Age 4	Age 5		
1997	407	0	2	0	2	0.49% ²
1998	2,999	5	42	2	49	1.63%
1999	1,744	1	0	0	1	0.06%
2000	1,503	0	1	0	1	0.07%
2001	2,146	0	4	0	4	0.19%
2002	2,201	4	5	0	9	0.41%
2003	1,418	0	3	1	4	0.28%
2004	4,194	3	13	0	16	0.38%
2005	2,358	0	3	0	3	0.13%
2006	4,130	32	31	2	65	1.57%
2007	3,736	10	21	0	31	0.83%
2008	1,071	4	3	0	7	0.65%
2009	3,641	2	4	0	6	0.16%
2010	4,064	4	13	1	18	0.44%
2011	513	0	0	0	0	0.00%
2012	201	0	0	0	0	0.00%
2013	1,432	0	0	0	0	0.00%
2014	1,104	0	3	0	3	0.27%
2015	1,783	2	2	0	4	0.22%
2016	2,578	1	0		1	0.04%
2017	2,238	2				

1. CESRF adult returns and smolt-to-adult survival values are understated relative to wild/natural values since these figures are not adjusted for differential harvest rates in mark selective fisheries in marine and lower Columbia River fisheries.
2. The reliability of the 400kHz detection gear precluded an accurate accounting of all 1997 brood PIT returns. Therefore, this is not a true SAR. It is presented for relative within-year comparison only and should NOT be compared to SARs for other years.

Table 42. Overall McNary Dam smolt to Bonneville Dam adult return rates (SAR) based on juvenile and adult detections of wild/natural Yakima R. spring Chinook PIT-tagged and released at Roza Dam (Table B.74 in McCann et al. 2020).

Juvenile migration year	Smolts arriving MCN ^A	MCN-to-BOA without Jacks			MCN-to-BOA with Jacks		
		%SAR Estimate	Non-parametric CI		%SAR Estimate	Non-parametric CI	
			90% LL	90% UL		90% LL	90% UL
2000	2,581	6.82	6.04	7.72	7.40	6.58	9.34
2001	521	1.54	0.75	2.52	1.92	0.98	3.04
2002	2,130	2.25	1.75	2.83	2.30	1.79	2.87
2003	2,143	2.47	1.97	3.03	2.89	2.34	3.50
2004	1,297	3.70	2.90	4.57	3.78	2.94	4.64
2005	521	1.34	0.57	2.22	1.34	0.57	2.22
2006	565	1.59	0.74	2.53	1.77	0.87	2.80
2007	362	1.93	0.84	3.17	1.93	0.84	3.17
2008	509	6.87	4.97	8.80	9.23	7.05	11.40
2009	983	4.99	3.85	6.29	5.60	4.35	6.97
2010 ^B	---	---	---	---	---	---	---
2011	411	0.97	0.23	1.82	0.97	0.23	1.82
2012	826	2.79	1.89	3.88	3.27	2.28	4.43
2013	704	1.42	0.70	2.19	1.56	0.82	2.37
2014 ^B	---	---	---	---	---	---	---
2015	238	2.10	0.57	4.11	2.52	0.76	4.86
2016 ^B	---	---	---	---	---	---	---
2017 ^B	---	---	---	---	---	---	---
2018 ^C	160	0.62	0.00	2.40	0.62	0.00	2.00
Arithmetic mean (incl. zeros)		2.76			3.14		
Geometric mean (excl. zeros)		2.22			2.44		

^A Estimated population of tagged study fish alive to MCN tailrace (included fish detected at the dam and those estimated to pass undetected). CJS estimation of S1 uses PIT-tags detected on bird colonies in the Columbia River estuary and adult detections to augment the NOAA Trawl detections below BON and the Logit link.

^B Too few or no PIT-tags released to obtain reliable estimate of smolts arriving at MCN. Therefore, estimate of SAR not possible.

^C Incomplete, 2-salt returns through September 25, 2020.

Table 43. Overall McNary Dam smolt to Bonneville Dam adult return rates (SAR) based on juvenile and adult detections of CESRF PIT-tagged spring Chinook (Table B.80 in McCann et al. 2020).

Juvenile migration year	Smolts arriving MCN ^A	MCN-to-BOA without Jacks			MCN-to-BOA with Jacks		
		%SAR Estimate	Non-parametric CI		%SAR Estimate	Non-parametric CI	
			90% LL	90% UL		90% LL	90% UL
2000	14,416	3.61	3.34	3.91	3.95	3.65	4.26
2001	9,269	0.28	0.20	0.37	0.29	0.20	0.38
2002	11,753	1.36	1.18	1.54	1.72	1.52	1.91
2003	11,974	0.59	0.48	0.71	0.86	0.72	1.00
2004	7,986	1.54	1.31	1.78	1.85	1.60	2.11
2005	5,789	0.66	0.48	0.84	0.78	0.59	0.98
2006	10,285	1.23	1.06	1.43	1.59	1.39	1.81
2007	12,654	1.01	0.87	1.16	1.51	1.32	1.69
2008	11,752	3.15	2.86	3.43	5.03	4.64	5.39
2009	15,386	1.82	1.64	2.00	2.29	2.08	2.50
2010	12,479	1.51	1.33	1.71	2.53	2.27	2.78
2011	11,886	0.93	0.79	1.08	1.20	1.03	1.37
2012	15,736	1.22	1.08	1.37	1.76	1.57	1.94
2013	13,261	1.38	1.20	1.54	1.95	1.74	2.17
2014	12,856	0.58	0.48	0.70	0.84	0.72	0.98
2015	10,639	1.02	0.85	1.20	1.86	1.62	2.11
2016	13,837	0.87	0.75	1.01	1.52	1.34	1.71
2017	11,199	0.62	0.50	0.75	0.74	0.61	0.89
2018 ^B	11,809	0.53	0.42	0.65	0.83	0.68	0.98
Arithmetic mean (incl. zeros)		1.26			1.74		
Geometric mean (excl. zeros)		1.05			1.44		

^A Estimated population of tagged study fish alive to MCN tailrace (included fish detected at the dam and those estimated to pass undetected). CJS estimation of S1 uses PIT-tags detected on bird colonies in the Columbia River estuary and adult detections to augment the NOAA Trawl detections below BON and the Logit link function.

^B Incomplete, 2-salt returns through September 25, 2020.

Table 44. Estimated release-to-adult survival of PIT-tagged CESRF fish (CESRF tagged smolts to Bonneville and Roza Dam adult returns).

Brood Year	Number Tagged ¹	Adult Detections at Bonn. Dam					Adult Detections at Roza Dam				
		Age3	Age4	Age5	Total	SAR	Age3	Age4	Age5	Total	SAR
1997 ²	39,892	18	182	4	204	0.51%	65	517	16	598	1.50%
1998	37,388	49	478	48	575	1.54%	54	310	34	398	1.06%
1999	38,793	1	25	1	27	0.07%	1	22	0	23	0.06%
2000	37,582	42	159	2	203	0.54%	37	112	1	150	0.40%
2001	36,523	32	71	0	103	0.28%	22	58	0	80	0.22%
2002 ³	39,003	25	119	4	148	0.38%	15	80	2	97	0.25%
2003	38,916	7	37	1	45	0.12%	3	27	1	31	0.08%
2004	36,426	37	123	4	164	0.45%	24	98	3	125	0.34%
2005	39,119	63	126	2	191	0.49%	44	96	2	142	0.36%
2006	38,595	221	354	15	590	1.53%	187	264	11	462	1.20%
2007	38,618	73	279	3	355	0.92%	55	182	3	240	0.62%
2008	39,013	135	192	3	330	0.85%	81	132	2	215	0.55%
2009	36,239	32	110	3	145	0.40%	23	85	2	110	0.30%
2010	38,737	85	187	6	278	0.72%	62	142	3	207	0.53%
2011	38,165	77	191	2	270	0.71%	57	122	2	181	0.47%
2012	38,343	33	75	0	108	0.28%	10	59	0	69	0.18%
2013	38,278	90	110	0	200	0.52%	68	84	0	152	0.40%
2014	38,119	92	121	1	214	0.56%	64	66	1	131	0.34%
2015	38,029	15	69	0	84	0.22%	6	51	0	57	0.15%
2016	38,061	34	64		98	0.26%	20	42		62	0.16%
2017	37,709	39					26				

1. When tag detection data are available, this is the number of unique PIT tags physically detected leaving the acclimation sites. Otherwise, this is the number of fish PIT tagged less documented mortalities of PIT-tagged fish from tagging to release.
2. BY1997 used 400 kHz tags and Bonneville Dam was not fully configured for adult detection of this type of tag; therefore we saw more detections at Roza Dam where fish were manually wanded for adult PIT detections.
3. Includes HxH fish beginning with this brood year.

Table 45. Estimated release-to-adult survival of non-PIT-tagged CESRF fish (CESRF tagged smolts to Roza Dam adult returns).

Brood Year	Number Tagged ¹	Adult Returns to Roza Dam				
		Age3	Age4	Age5	Total	SAR
1997 ²	346,156	623	5,663	120	6,406	1.85%
1998	552,295	936	5,834	534	7,304	1.32%
1999	719,996	103	652	13	768	0.11%
2000	796,703	1,005	2,764	69	3,837	0.48%
2001	333,713	290	791	9	1,091	0.33%
2002 ³	797,901	332	1,771	135	2,238	0.28%
2003	785,776	115	1,568	14	1,696	0.22%
2004	749,022	683	3,688	202	4,574	0.61%
2005	820,883	1,012	5,302	22	6,336	0.77%
2006	604,200	2,383	6,427	287	9,096	1.51%
2007	732,647	1,024	5,645	87	6,756	0.92%
2008	810,292	1,552	3,680	76	5,308	0.66%
2009	796,702	389	3,106	67	3,562	0.45%
2010	756,044	721	3,618	28	4,368	0.58%
2011	731,017	780	2,318	51	3,149	0.43%
2012	764,373	172	2,274	12	2,458	0.32%
2013	608,477	718	2,386	0	3,104	0.51%
2014	647,111	644	1,511	10	2,165	0.33%
2015	616,918	237	1,242	0	1,479	0.24%
2016	631,147	158	1,211		1,369	0.22%
2017	672,960	376				

1. These fish were adipose fin-clipped, coded-wire tagged, and (beginning with 4 of 16 ponds in 1998) elastomer eye tagged. This is the number of fish physically counted at tagging.
2. BY1997 used 400 kHz tags and Bonneville Dam was not fully configured for adult detection of this type of tag; therefore we saw more detections at Roza Dam where fish were manually wanded for adult PIT detections.
3. Includes HxH fish beginning with this brood year.

Harvest Monitoring

Yakima Basin Fisheries

For spring fisheries in the Yakima River Basin, both the WDFW and the Yakama Nation employ two technicians and one biologist to monitor and evaluate in-basin harvest in the respective sport and tribal fisheries. Harvest monitoring consists of on-the-water surveys to collect catch data and to record tag information (e.g., elastomer, CWT, etc.) where possible for adipose-clipped fish. Survey data are expanded for time, area, and effort using standard methods to derive estimates of total in-basin harvest by fishery type (sport and tribal) and catch type (CESRF or wild denoted by adipose presence/absence). Results are presented in Table 46.

Columbia Basin Fisheries

Standard run reconstruction techniques are employed to derive estimates of harvest from the Columbia River mouth to the Yakima River mouth for spring Chinook. Data from databases maintained by the *United States versus Oregon* Technical Advisory Committee (TAC) are used to obtain harvest rate estimates downstream of the Yakima River for the aggregate Yakima River spring Chinook population and to estimate passage losses from Bonneville through McNary reservoirs. These data, combined with the Prosser Dam counts and estimated harvest below Prosser, are used to derive a Columbia River mouth run size estimate and Columbia River mainstem harvest estimate for Yakima spring Chinook. Results are presented in Table 47.

Table 46. Spring Chinook harvest in the Yakima River Basin, 1985-present.

Year	Tribal		Non-Tribal		River Totals		Total	Harvest Rate ¹
	CESRF	Wild	CESRF	Wild	CESRF	Wild		
1985		865		0		865	865	19.0%
1986		1,340		0		1,340	1,340	14.2%
1987		517		0		517	517	11.6%
1988		444		0		444	444	10.5%
1989		747		0		747	747	15.2%
1990		663		0		663	663	15.2%
1991		32		0		32	32	1.1%
1992		345		0		345	345	7.5%
1993		129		0		129	129	3.3%
1994		25		0		25	25	1.9%
1995		79		0		79	79	11.9%
1996		475		0		475	475	14.9%
1997		575		0		575	575	18.1%
1998		188		0		188	188	9.9%
1999		604		0		604	604	21.7%
2000	53	2,305		100	53	2,405	2,458	12.9%
2001	572	2,034	1,252	772	1,825	2,806	4,630	19.9%
2002	1,373	1,207	492	36 ²	1,865	1,243	3,108	20.6%
2003	134	306	0	0	134	306	440	6.3%
2004	289	712	569	109 ²	858	820	1,679	11.0%
2005	46	428	0	0	46	428	474	5.4%
2006	246	354	0	0	246	354	600	9.5%
2007	123	156	0	0	123	156	279	6.5%
2008	521	414	586	11 ²	1,107	426	1,532	17.8%
2009	1,089	715	541	8 ²	1,630	722	2,353	19.4%
2010	345	194	1,154	48 ²	1,499	241	1,741	13.2%
2011	1,361	1,261	1,579	179 ²	2,940	1,440	4,380	24.4%
2012	1,220	1,302	735	63 ²	1,955	1,364	3,320	27.5%
2013	846	975	786	46 ²	1,632	1,021	2,653	25.9%
2014	576	715	826	54 ²	1,402	769	2,171	19.2%
2015	121	271	385	38 ²	506	309	815	8.7%
2016	103	185	132	24 ²	235	209	444	6.4%
2017	217	201	750	104 ²	967	305	1,272	17.8%
2018	154	115	259	20 ²	413	136	548	15.2%
2019	24	16	0	0	24	16	40	1.8%
2020	26	42	0	0	26	42	68	2.0%
Mean	469	580	502	76	972	599	1,098	13.0%

1. Harvest rate is the total Yakima Basin harvest as a percentage of the Yakima River mouth run size.
2. Includes estimate of post-release mortality of unmarked fish.

Table 47. Estimated run size, harvest, and harvest rates of Yakima Basin spring Chinook in Columbia River mainstem and terminal area fisheries, 1986-present.

Year	Columbia R. Mouth Run Size	Col. R.	BON to McNary Harvest	Yakima R. Mouth Run Size	Yakima River Harvest	Columbia Basin Harvest Summary			Col. Basin Harvest Rate	
		Mouth to BON Harvest				Total	Wild	CESRF	Total	Wild
1986	13,567	280	802	9,439	1,340	2,423	2,423	0	17.9%	17.9%
1987	6,160	96	378	4,443	517	991	991	0	16.1%	16.1%
1988	5,674	363	401	4,246	444	1,208	1,208	0	21.3%	21.3%
1989	8,919	213	683	4,914	747	1,642	1,642	0	18.4%	18.4%
1990	6,954	352	480	4,372	663	1,495	1,495	0	21.5%	21.5%
1991	4,650	184	291	2,906	32	507	507	0	10.9%	10.9%
1992	6,207	103	380	4,599	345	827	827	0	13.3%	13.3%
1993	5,132	44	315	3,919	129	488	488	0	9.5%	9.5%
1994	2,251	87	113	1,302	25	225	225	0	10.0%	10.0%
1995	1,394	1	69	666	79	149	149	0	10.7%	10.7%
1996	5,898	6	309	3,179	475	790	790	0	13.4%	13.4%
1997	5,192	3	348	3,173	575	926	926	0	17.8%	17.8%
1998	2,868	3	144	1,903	188	335	335	0	11.7%	11.7%
1999	4,154	4	192	2,781	604	800	800	0	19.3%	19.3%
2000	28,753	58	1,752	19,101	2,458	4,267	4,144	123	14.8%	14.8%
2001	32,307	971	4,281	24,149	4,630	9,882	5,685	4,197	30.6%	28.7%
2002	25,256	1,275	2,877	15,814	3,108	7,259	2,736	4,524	28.7%	23.9%
2003	10,278	286	903	7,227	440	1,628	987	641	15.8%	14.7%
2004	24,212	1,023	2,330	16,820	1,679	5,031	2,877	2,154	20.8%	16.2%
2005	13,317	354	906	9,589	474	1,735	1,375	360	13.0%	12.1%
2006	12,197	311	944	6,594	600	1,855	1,068	787	15.2%	13.5%
2007	5,223	174	457	4,457	279	910	449	461	17.4%	15.1%
2008	12,554	1,204	1,870	9,273	1,532	4,607	1,360	3,247	36.7%	25.2%
2009	13,693	1,210	1,089	11,395	2,353	4,651	1,318	3,333	34.0%	23.9%
2010	18,565	1,631	2,778	13,745	1,741	6,149	1,516	4,633	33.1%	21.8%
2011	23,316	1,098	1,794	18,520	4,380	7,272	2,590	4,682	31.2%	22.4%
2012	17,315	856	1,633	12,616	3,320	5,809	2,370	3,438	33.5%	26.6%
2013	14,933	880	974	10,602	2,653	4,507	1,817	2,690	30.2%	23.3%
2014	17,303	716	2,222	11,868	2,171	5,110	2,097	3,012	29.5%	22.5%
2015	11,991	476	1,440	9,848	815	2,731	1,457	1,274	22.8%	17.8%
2016	10,107	454	996	7,281	444	1,894	971	923	18.7%	15.5%
2017	12,196	493	920	7,544	1,272	2,685	853	1,831	22.0%	13.5%
2018	6,237	248	636	3,737	548	1,433	459	975	23.0%	16.4%
2019	3,784	68	260	2,251	40	368	131	238	9.7%	8.6%
2020 ¹	5,764	62	347	3,413	68	476	276	200	8.3%	7.7%
Mean	11,381	445	1,038	7,934	1,176	2,659	1,410	1,249	20.0%	17.0%

1. Preliminary.

Marine Fisheries

Based on available CWT information, harvest managers have long assumed that Columbia River spring Chinook are not harvested in any abundance in marine fisheries as the timing of their ocean migration does not generally overlap either spatially or temporally with the occurrence of marine fisheries (TAC 1997). The Regional Mark Information System (RMIS) will be queried regularly for any CWT recoveries of CESRF releases in ocean or Columbia River mainstem fisheries. Table 48 gives the results of a query of the RMIS database run on Jan. 5, 2021 for CESRF spring Chinook CWTs released in brood years 1997-2015 and Figure 8 shows recovery locations for CWTs recovered in marine fisheries 2008-2012. Based on the information reported to RMIS to date, it is believed that marine harvest accounts for about 0-4% of the total harvest of Yakima Basin spring Chinook. The apparent increase for brood year 2013 may be attributable to a number of factors including: preliminary data or changes in fish distribution, ecological conditions, or sampling rates. CWT recovery data for brood year 2016 were considered too incomplete to report at this time.

Table 48. Marine and freshwater recoveries of CWTs from brood year 1997-2015 releases of spring Chinook from the CESRF as reported to the Regional Mark Information System (RMIS) Jan. 5, 2021.

Brood Year	Observed CWT Recoveries			Expanded CWT Recoveries		
	Marine	Fresh	Marine %	Marine	Fresh	Marine %
1997	5	56	8.2%	8	321	2.4%
1998	2	53	3.6%	2	228	0.9%
1999		2	0.0%		9	0.0%
2000		14	0.0%		34	0.0%
2001		1	0.0%		1	0.0%
2002		7	0.0%		36	0.0%
2003		4	0.0%		10	0.0%
2004	2	154	1.3%	15	526	2.8%
2005	2	96	2.0%	2	304	0.7%
2006	14	328	4.1%	16	1160	1.4%
2007	8	145	5.2%	13	1139	1.1%
2008	5	245	2.0%	7	1634	0.4%
2009	4	91	4.2%	7	588	1.2%
2010	4	164	2.4%	9	948	0.9%
2011	5	186	2.6%	5	1030	0.5%
2012	4	73	5.2%	2	273	0.7%
2013	9	65	12.2%	20	534	3.6%
2014	4	71	5.3%	8	533	1.5%
2015 ¹	2	23	8.0%	2	49	3.9%

1. Reporting of CWT recoveries to the RMIS database typically lags actual fisheries by one to two years. Therefore, CWT recovery data for brood year 2015 are considered preliminary or incomplete.




Figure 9. Marine recovery locations of coded-wire-tagged CESRF spring Chinook, recovery years 2008-2012.

Literature Cited

- BPA (Bonneville Power Administration). 1990. Yakima-Klickitat Production Project Preliminary Design Report and Appendices. Bonneville Power Administration, Portland, OR.
- Copeland, T., D.A. Venditti, and B.R. Barnett. 2014. The Importance of Juvenile Migration Tactics to Adult Recruitment in Stream-Type Chinook Salmon Populations. *Transactions of the American Fisheries Society* 143:1460-1475.
- Knudsen C.M., S.L. Schroder, T.N. Pearsons, J.A. Rau, A.L. Fritts, and C.R. Strom. 2003. Monitoring Phenotypic and Demographic Traits of upper Yakima River Hatchery and Wild Spring Chinook: Gametic and juvenile Traits. YKFP Annual Report 2002.
- Knudsen, C.M. (editor). 2004. Reproductive Ecology of Yakima River hatchery and wild spring Chinook. Annual Report 2003, Project Number 1995-063-25. BPA Report DOE/BP-00013756-3.
- Knudsen, C. M., M. V. Johnston, S. L. Schroder, W. J. Bosch, D. E. Fast, and C. R. Strom. 2009. Effects of Passive Integrated Transponder Tags on Smolt-to-Adult Recruit Survival, Growth, and Behavior of Hatchery Spring Chinook Salmon. *North American Journal of Fisheries Management* 29:658-669.
- Maynard, D. J., T. A. Flagg, and C. V. W. Mahnken. 1995. A review of seminatural culture strategies for enhancing the post-release survival of anadromous salmonids. *Am. Fish. Soc. Symp.* 15:307-314.
- McCann, J., B. Chockley, E. Cooper, G. Scheer, S. Haeseker, B. Lessard, T. Copeland, J. Ebel, A. Storch, and D. Rawding. 2020. [Comparative Survival Study](#) of PIT-tagged Spring/Summer/Fall Chinook, Summer Steelhead, and Sockeye 2020 Annual Report (BPA Project #19960200). Fish Passage Center, Portland, Oregon.
- NPPC (Northwest Power Planning Council). 1982. Columbia River Basin Fish and Wildlife Program. Adopted November 15, 1982. Northwest Power Planning Council, Portland, OR.
- Pandit, S. 2020. 2019 Annual Chandler Certification for Yearling Out-migrating Spring Chinook Smolt. Appendix C in Blodgett et al., editors, Yakima/Klickitat Fisheries Project Monitoring and Evaluation, Yakima Subbasin, Project Number 1995-063-25, 378 electronic pages (<https://www.cbfish.org/Document.mvc/Viewer/P177850>).
- TAC (United States versus Oregon Technical Advisory Committee). 1997. 1996 All Species Review, Columbia River Fish Management Plan. August 4, 1997. Columbia River Inter-Tribal Fish Commission, Portland, Oregon.

Appendix A. Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.

<i>Brood Year</i>	<i>C.E. Pond</i>	<i>Accl. Pond</i>	<i>Treatment¹</i>			<i>Tag Information</i>				<i>First Release</i>	<i>Last Release</i>	<i>CWT Code</i>	<i>No. PIT</i>	<i>No. CWT</i>	<i>Est. Tot. Release²</i>
2006	CLE01	CFJ04	BIO	WW	3.5	Right	Red	Snout		3/15/2008	5/14/2008	190101	2,000	36,945	38,607
2006	CLE02	CFJ03	EWS	WW	3.5	Left	Red	Snout		3/15/2008	5/14/2008	190102	2,000	31,027	32,790
2006	CLE03	ESJ02	BIO	WW	3.2	Right	Green	Snout		3/15/2008	5/14/2008	190103	2,000	36,931	38,762
2006	CLE04	ESJ01	EWS	WW	3.2	Left	Green	Snout		3/15/2008	5/14/2008	190104	2,000	29,635	31,400
2006	CLE05	JCJ02	BIO	WW	3.3	Right	Orange	Snout		3/15/2008	5/14/2008	190105	2,000	36,735	38,383
2006	CLE06	JCJ01	EWS	WW	3.3	Left	Orange	Snout		3/15/2008	5/14/2008	190106	2,000	28,984	30,680
2006	CLE07	ESJ04	BIO	WW	3.4	Right	Green	Snout		3/15/2008	5/14/2008	190107	2,000	38,212	40,006
2006	CLE08	ESJ03	EWS	WW	3.4	Left	Green	Snout		3/15/2008	5/14/2008	190108	2,000	32,726	34,519
2006	CLE09	CFJ02	BIO	WW	3.4	Right	Red	Snout		3/15/2008	5/14/2008	190109	2,000	36,485	38,097
2006	CLE10	CFJ01	EWS	WW	3.4	Left	Red	Snout		3/15/2008	5/14/2008	190110	2,000	29,907	31,647
2006	CLE11	JCJ04	BIO	WW	3.3	Right	Orange	Snout		3/15/2008	5/14/2008	190111	2,000	39,491	40,703
2006	CLE12	JCJ03	EWS	WW	3.3	Left	Orange	Snout		3/15/2008	5/14/2008	190112	2,000	33,418	35,273
2006	CLE13	ESJ06	BIO	WW	3.4	Right	Green	Snout		3/15/2008	5/14/2008	190113	2,000	38,609	39,841
2006	CLE14	ESJ05	EWS	WW	3.4	Left	Green	Snout		3/15/2008	5/14/2008	190114	2,000	31,573	33,404
2006	CLE15	JCJ06	BIO	WW	3.4	Right	Orange	Snout		3/15/2008	5/14/2008	190115	2,000	36,844	38,619
2006	CLE16	JCJ05	EWS	WW	3.4	Left	Orange	Snout		3/15/2008	5/14/2008	190116	2,000	29,857	31,630
2006	CLE17	CFJ06	BIO	HH	3.2	Right	Red	Posterior Dorsal		3/15/2008	5/14/2008	190117	4,000	34,299	38,045
2006	CLE18	CFJ05	EWS	HH	3.2	Left	Red	Posterior Dorsal		3/15/2008	5/14/2008	190118	4,000	26,643	30,389

¹ BIO = BioVita (BioOregon Protein Inc.) or control diet; EWS = EWOS (EWOS Canada Ltd.). All fish were switched to BioVita diet beginning May 3, 2007. All fish are progeny of wild/natural parents unless denoted as HH which designates the hatchery control line. "Avg BKD" denotes the average BKD ELISA ranking of the female parents whose progeny were in these ponds.

² The number of fish released is estimated as the total number of fish counted at marking less mortalities documented from mark to release.

Appendix A. Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.

Brood Year	C.E. Pond	Accl. Pond	Treatment¹			Tag Information			First Release	Last Release	CWT Code	No. PIT	No. CWT	Est. Tot. Release²
2007	CLE01	JCJ06	BIO	WW	2.8	Right	Orange	Snout	3/15/2009	5/15/2009	190151	2,000	38,044	39,840
2007	CLE02	JCJ05	STF	WW	2.8	Left	Orange	Snout	3/15/2009	5/15/2009	190152	2,000	40,066	41,843
2007	CLE03	JCJ04	BIO	WW	2.7	Right	Orange	Snout	3/15/2009	5/15/2009	190153	2,000	40,843	42,647
2007	CLE04	JCJ03	STF	WW	2.7	Left	Orange	Snout	3/15/2009	5/15/2009	190154	2,000	40,196	41,979
2007	CLE05	CFJ06	BIO	WW	2.8	Right	Red	Snout	3/15/2009	5/15/2009	190155	2,000	40,855	42,717
2007	CLE06	CFJ05	STF	WW	2.8	Left	Red	Snout	3/15/2009	5/15/2009	190156	2,000	40,475	42,345
2007	CLE07	ESJ06	BIO	WW	2.6	Right	Green	Snout	3/15/2009	5/15/2009	190157	2,000	42,549	44,387
2007	CLE08	ESJ05	STF	WW	2.6	Left	Green	Snout	3/15/2009	5/15/2009	190158	2,000	43,243	45,080
2007	CLE09	CFJ02	BIO	HH	2.7	Right	Red	Posterior Dorsal	3/15/2009	5/15/2009	190159	4,000	43,803	47,625
2007	CLE10	CFJ01	STF	HH	2.7	Left	Red	Posterior Dorsal	3/15/2009	5/15/2009	190160	4,000	43,256	47,038
2007	CLE11	ESJ02	BIO	WW	2.8	Right	Green	Snout	3/15/2009	5/15/2009	190161	2,000	41,098	42,945
2007	CLE12	ESJ01	STF	WW	2.8	Left	Green	Snout	3/15/2009	5/15/2009	190162	2,001	40,535	42,405
2007	CLE13	ESJ04	BIO	WW	2.7	Right	Green	Snout	3/15/2009	5/15/2009	190163	2,009	39,308	41,190
2007	CLE14	ESJ03	STF	WW	2.7	Left	Green	Snout	3/15/2009	5/15/2009	190164	2,000	36,663	38,533
2007	CLE15	JCJ02	BIO	WW	2.9	Right	Orange	Snout	3/15/2009	5/15/2009	190165	2,000	40,312	42,083
2007	CLE16	JCJ01	STF	WW	2.9	Left	Orange	Snout	3/15/2009	5/15/2009	190166	2,000	40,594	42,426
2007	CLE17	CFJ03	STF	WW	2.8	Right	Red	Snout	3/15/2009	5/15/2009	190167	2,000	40,687	42,561
2007	CLE18	CFJ04	BIO	WW	2.8	Left	Red	Snout	3/15/2009	5/15/2009	190168	2,000	41,704	43,621

¹ BIO = BioVita (BioOregon Protein Inc.) or control diet; STF = salt-water transition diet at acclimation sites. All fish are progeny of wild/natural parents unless denoted as HH which designates the hatchery control line beginning with brood year 2002. "Avg BKD" denotes the average BKD ELISA ranking of the female parents whose progeny were in these ponds.

² The number of fish released is estimated as the total number of fish counted at marking less mortalities documented from mark to release.

Appendix A. Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.

Brood Year	C.E. Pond	Accl. Pond	Treatment¹			Tag Information			First Release	Last Release	CWT Code	No. PIT	No. CWT	Est. Tot. Release²
2008	CLE01	ESJ01	STF	WW	3.3	Right	Orange	Snout	3/15/2010	5/11/2010	190191	2,000	44,917	46,704
2008	CLE02	ESJ02	BIO	WW	3.3	Left	Orange	Snout	3/15/2010	5/11/2010	190192	2,000	45,576	47,414
2008	CLE03	CFJ03	STF	WW	3.2	Right	Red	Snout	3/15/2010	5/11/2010	190193	2,000	44,099	45,931
2008	CLE04	CFJ04	BIO	WW	3.2	Left	Red	Snout	3/15/2010	5/11/2010	190194	2,000	42,464	44,271
2008	CLE05	JCJ05	STF	WW	3.0	Right	Green	Snout	3/15/2010	5/11/2010	190195	2,000	46,118	47,936
2008	CLE06	JCJ06	BIO	WW	3.0	Left	Green	Snout	3/15/2010	5/11/2010	190196	2,000	43,708	45,466
2008	CLE07	ESJ05	STF	WW	3.2	Right	Orange	Snout	3/15/2010	5/11/2010	190197	2,000	48,468	50,299
2008	CLE08	ESJ06	BIO	WW	3.2	Left	Orange	Snout	3/15/2010	5/11/2010	190198	2,000	47,611	49,419
2008	CLE09	CFJ05	STF	HH	2.9	Right	Red	Posterior Dorsal	3/15/2010	5/11/2010	190199	4,000	45,169	48,942
2008	CLE10	CFJ06	BIO	HH	2.9	Left	Red	Posterior Dorsal	3/15/2010	5/11/2010	190201	4,000	44,493	48,254
2008	CLE11	JCJ01	STF	WW	3.3	Right	Green	Snout	3/15/2010	5/11/2010	190202	2,000	44,583	46,413
2008	CLE12	JCJ02	BIO	WW	3.3	Left	Green	Snout	3/15/2010	5/11/2010	190203	2,000	45,086	46,856
2008	CLE13	ESJ03	STF	WW	3.1	Right	Orange	Snout	3/15/2010	5/11/2010	190204	2,000	45,518	47,317
2008	CLE14	ESJ04	BIO	WW	3.1	Left	Orange	Snout	3/15/2010	5/11/2010	190205	2,000	44,879	46,704
2008	CLE15	CFJ01	STF	WW	3.2	Right	Red	Snout	3/15/2010	5/11/2010	190206	2,000	45,169	46,893
2008	CLE16	CFJ02	BIO	WW	3.2	Left	Red	Snout	3/15/2010	5/11/2010	190207	2,000	44,149	45,962
2008	CLE17	JCJ03	STF	WW	3.2	Right	Green	Snout	3/15/2010	5/11/2010	190208	2,000	45,807	47,580
2008	CLE18	JCJ04	BIO	WW	3.2	Left	Green	Snout	3/15/2010	5/11/2010	190209	2,000	45,157	46,944

¹ BIO = BioVita (BioOregon Protein Inc.) or control diet; STF = salt-water transition diet at acclimation sites. All fish are progeny of wild/natural parents unless denoted as HH which designates the hatchery control line beginning with brood year 2002. "Avg BKD" denotes the average BKD ELISA ranking of the female parents whose progeny were in these ponds.

² The number of fish released is estimated as the total number of fish counted at marking less mortalities documented from mark to release.

Appendix A. Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.

Brood Year	C.E. Pond	Accl. Pond	Treatment¹			Tag Information				First Release	Last Release	CWT Code	No. PIT	No. CWT	Est. Tot. Release²
2009	CLE01	CFJ05	STF	HH	3.0	Right	Red	Posterior	Dorsal	3/15/2011	5/16/2011	190215	4,000	40,109	43,965
2009	CLE02	CFJ06	BIO	HH	3.0	Left	Red	Posterior	Dorsal	3/15/2011	5/16/2011	190216	4,000	41,012	44,806
2009	CLE03	JCJ01	STF	WW	3.0	Right	Orange	Snout		3/15/2011	3/31/2011	190217	2,000	37,245	39,048
2009	CLE04	JCJ02	BIO	WW	3.0	Left	Orange	Snout		3/15/2011	3/31/2011	190218	2,000	42,212	44,053
2009	CLE05	CFJ01	STF	WW	3.2	Right	Red	Snout		3/15/2011	5/16/2011	190219	2,000	47,016	48,761
2009	CLE06	CFJ02	BIO	WW	3.2	Left	Red	Snout		3/15/2011	5/16/2011	190220	2,000	46,733	48,569
2009	CLE07	ESJ05	STF	WW	3.1	Right	Green	Snout		3/15/2011	5/16/2011	190221	2,000	46,302	48,089
2009	CLE08	ESJ06	BIO	WW	3.1	Left	Green	Snout		3/15/2011	5/16/2011	190222	2,000	46,969	48,721
2009	CLE09	ESJ01	STF	WW	3.0	Right	Green	Snout		3/15/2011	5/16/2011	190223	2,000	43,612	45,379
2009	CLE10	ESJ02	BIO	WW	3.0	Left	Green	Snout		3/15/2011	5/16/2011	190224	2,000	43,173	44,962
2009	CLE11	JCJ05	STF	WW	3.1	Right	Orange	Snout		3/15/2011	3/31/2011	190225	2,000	47,585	49,306
2009	CLE12	JCJ06	BIO	WW	3.1	Left	Orange	Snout		3/15/2011	3/31/2011	190226	2,000	47,644	49,434
2009	CLE13	ESJ03	STF	WW	3.2	Right	Green	Snout		3/15/2011	5/16/2011	190227	2,000	45,277	47,036
2009	CLE14	ESJ04	BIO	WW	3.2	Left	Green	Snout		3/15/2011	5/16/2011	190228	2,000	45,529	47,208
2009	CLE15	JCJ03	STF	WW	3.1	Right	Orange	Snout		3/15/2011	3/31/2011	190229	2,000	43,825	45,592
2009	CLE16	JCJ04	BIO	WW	3.1	Left	Orange	Snout		3/15/2011	3/31/2011	190230	2,000	43,209	44,990
2009	CLE17	CFJ03	STF	WW	3.2	Right	Red	Snout		3/15/2011	5/16/2011	190231	2,000	45,587	47,451
2009	CLE18	CFJ04	BIO	WW	3.2	Left	Red	Snout		3/15/2011	5/16/2011	190232	2,000	43,952	45,571

¹ BIO = BioVita (BioOregon Protein Inc.) or control diet; STF = salt-water transition diet at acclimation sites. All fish are progeny of wild/natural parents unless denoted as HH which designates the hatchery control line beginning with brood year 2002. "Avg BKD" denotes the average BKD ELISA ranking of the female parents whose progeny were in these ponds.

² The number of fish released is estimated as the total number of fish counted at marking less mortalities documented from mark to release.

Appendix A. Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.

Brood Year	C.E. Pond	Accl. Pond	Treatment¹ /Avg BKD	Tag Information					First Release	Last Release	CWT Code	No. PIT	No. CWT	Est. Tot. Release²
2010	CLE01	CFJ05	STF WW 4.2	Right	Red	Snout		3/15/2012	5/14/2012	190256	2,000	40,221	41,972	
2010	CLE02	CFJ06	BIO WW 4.2	Left	Red	Snout		3/15/2012	5/14/2012	190257	2,000	40,845	42,664	
2010	CLE03	CFJ03	STF HH 4.0	Right	Red	Posterior Dorsal		3/15/2012	5/14/2012	190258	4,000	43,725	47,415	
2010	CLE04	CFJ04	BIO HH 4.0	Left	Red	Posterior Dorsal		3/15/2012	5/14/2012	190259	4,000	40,976	44,615	
2010	CLE05	ESJ01	STF WW 4.2	Right	Green	Snout		3/15/2012	5/14/2012	190260	2,000	40,710	42,374	
2010	CLE06	ESJ02	BIO WW 4.2	Left	Green	Snout		3/15/2012	5/14/2012	190261	2,000	40,419	42,157	
2010	CLE07	JCJ01	STF WW 4.0	Right	Orange	Snout		3/15/2012	5/14/2012	190262	2,000	43,833	45,471	
2010	CLE08	JCJ02	BIO WW 4.0	Left	Orange	Snout		3/15/2012	5/14/2012	190263	2,000	43,815	45,573	
2010	CLE09	ESJ03	STF WW 4.1	Right	Green	Snout		3/15/2012	5/14/2012	190264	2,000	42,528	44,257	
2010	CLE10	ESJ04	BIO WW 4.1	Left	Green	Snout		3/15/2012	5/14/2012	190265	2,000	42,649	44,443	
2010	CLE11	ESJ05	STF WW 4.2	Right	Green	Snout		3/15/2012	5/14/2012	190266	2,000	43,878	45,633	
2010	CLE12	ESJ06	BIO WW 4.2	Left	Green	Snout		3/15/2012	5/14/2012	190267	2,000	43,750	45,498	
2010	CLE13	JCJ03	STF WW 4.2	Right	Orange	Snout		3/15/2012	5/14/2012	190268	2,000	41,816	43,473	
2010	CLE14	JCJ04	BIO WW 4.2	Left	Orange	Snout		3/15/2012	5/14/2012	190269	2,000	41,052	42,772	
2010	CLE15	JCJ05	STF WW 4.1	Right	Orange	Snout		3/15/2012	5/14/2012	190270	2,000	42,894	44,603	
2010	CLE16	JCJ06	BIO WW 4.1	Left	Orange	Snout		3/15/2012	5/14/2012	190271	2,000	42,371	44,107	
2010	CLE17	CFJ01	STF WW 4.2	Right	Red	Snout		3/15/2012	5/14/2012	190272	2,000	42,329	44,128	
2010	CLE18	CFJ02	BIO WW 4.2	Left	Red	Snout		3/15/2012	5/14/2012	190273	2,000	41,829	43,626	

¹ BIO = BioVita (BioOregon Protein Inc.) or control diet; STF = salt-water transition diet at acclimation sites. All fish are progeny of wild/natural parents unless denoted as HH which designates the hatchery control line beginning with brood year 2002. "Avg BKD" denotes the average BKD ELISA ranking of the female parents whose progeny were in these ponds.

² The number of fish released is estimated as the total number of fish counted at marking less mortalities documented from mark to release.

Appendix A. Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.

<i>Brood Year</i>	<i>C.E. Pond</i>	<i>Accl. Pond</i>	<i>Treatment¹</i>			<i>Tag Information</i>			<i>First Release</i>	<i>Last Release</i>	<i>CWT Code</i>	<i>No. PIT</i>	<i>No. CWT</i>	<i>Est. Tot. Release²</i>
2011	CLE01	JCJ05	STF	WN	4.1	Right	Orange	Snout	3/15/2013	5/15/2013	190320	2,000	42,452	44,225
2011	CLE02	JCJ06	BIO	WN	4.1	Left	Orange	Snout	3/15/2013	5/15/2013	190321	2,000	42,217	44,056
2011	CLE03	CFJ05	STF	HC	4.0	Right	Red	Posterior Dorsal	3/15/2013	5/15/2013	190322	4,000	38,432	42,092
2011	CLE04	CFJ06	BIO	HC	4.0	Left	Red	Posterior Dorsal	3/15/2013	5/15/2013	190323	4,000	38,743	42,609
2011	CLE05	ESJ01	STF	WN	4.1	Right	Green	Snout	3/15/2013	5/15/2013	190324	2,000	38,404	40,250
2011	CLE06	ESJ02	BIO	WN	4.1	Left	Green	Snout	3/15/2013	5/15/2013	190325	2,000	37,931	39,731
2011	CLE07	CFJ01	STF	WN	4.1	Right	Red	Snout	3/15/2013	5/15/2013	190326	2,000	40,449	42,308
2011	CLE08	CFJ02	BIO	WN	4.1	Left	Red	Snout	3/15/2013	5/15/2013	190327	2,000	39,281	41,088
2011	CLE09	JCJ03	STF	WN	4.0	Right	Orange	Snout	3/15/2013	5/15/2013	190328	2,000	43,588	45,243
2011	CLE10	JCJ04	BIO	WN	4.0	Left	Orange	Snout	3/15/2013	5/15/2013	190329	2,000	41,715	43,288
2011	CLE11	ESJ05	STF	WN	4.0	Right	Green	Snout	3/15/2013	5/15/2013	190330	2,000	40,964	42,610
2011	CLE12	ESJ06	BIO	WN	4.0	Left	Green	Snout	3/15/2013	5/15/2013	190331	2,000	40,905	42,759
2011	CLE13	CFJ03	STF	WN	4.0	Right	Red	Snout	3/15/2013	5/15/2013	190332	2,000	42,298	44,190
2011	CLE14	CFJ04	BIO	WN	4.0	Left	Red	Snout	3/15/2013	5/15/2013	190333	2,000	41,111	43,003
2011	CLE15	JCJ01	STF	WN	3.9	Right	Orange	Snout	3/15/2013	5/15/2013	190334	2,000	42,769	44,590
2011	CLE16	JCJ02	BIO	WN	3.9	Left	Orange	Snout	3/15/2013	5/15/2013	190335	2,000	42,230	44,036
2011	CLE17	ESJ03	STF	WN	4.0	Right	Green	Snout	3/15/2013	5/15/2013	190336	2,000	39,770	41,479
2011	CLE18	ESJ04	BIO	WN	4.0	Left	Green	Snout	3/15/2013	5/15/2013	190337	2,000	39,823	41,625

¹ BIO = BioVita (BioOregon Protein Inc.) or control diet; STF = salt-water transition diet at acclimation sites. All fish are progeny of wild/natural parents unless denoted as HC which designates the hatchery control line beginning with brood year 2002. "Avg BKD" denotes the average BKD ELISA ranking of the female parents whose progeny were in these ponds.

² The number of fish released is estimated as the total number of fish counted at marking less mortalities documented from mark to release.

Appendix A. Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.

<i>Brood Year</i>	<i>C.E. Pond</i>	<i>Accl. Pond</i>	<i>Treatment¹</i>			<i>Tag Information</i>				<i>First Release</i>	<i>Last Release</i>	<i>CWT Code</i>	<i>No. PIT</i>	<i>No. CWT</i>	<i>Est. Tot. Release²</i>
2012	CLE01	ESJ03	STF	WN	3.7	Right	Green	Snout		3/15/2014	5/15/2014	190367	2,000	44,358	45,902
2012	CLE02	ESJ04	BIO	WN	3.7	Left	Green	Snout		3/15/2014	5/15/2014	190368	2,000	44,999	46,758
2012	CLE03	CFJ03	STF	HC	3.8	Right	Red	Posterior Dorsal		3/15/2014	5/15/2014	190369	4,000	42,147	45,670
2012	CLE04	CFJ04	BIO	HC	3.8	Left	Red	Posterior Dorsal		3/15/2014	5/15/2014	190370	4,000	41,497	45,010
2012	CLE05	ESJ05	STF	WN	3.8	Right	Green	Snout		3/15/2014	5/15/2014	190371	2,000	43,627	45,512
2012	CLE06	ESJ06	BIO	WN	3.8	Left	Green	Snout		3/15/2014	5/15/2014	190372	2,000	44,507	46,420
2012	CLE07	CFJ05	STF	WN	3.7	Right	Red	Snout		3/15/2014	5/15/2014	190373	2,000	41,067	42,932
2012	CLE08	CFJ06	BIO	WN	3.7	Left	Red	Snout		3/15/2014	5/15/2014	190374	2,000	37,499	39,367
2012	CLE09	CFJ01	STF	WN	3.7	Right	Red	Snout		3/15/2014	5/15/2014	190375	2,000	42,001	43,629
2012	CLE10	CFJ02	BIO	WN	3.7	Left	Red	Snout		3/15/2014	5/15/2014	190376	2,000	38,364	40,124
2012	CLE11	JCJ01	STF	WN	3.8	Right	Orange	Snout		3/15/2014	5/15/2014	190377	2,000	41,425	43,279
2012	CLE12	JCJ02	BIO	WN	3.8	Left	Orange	Snout		3/15/2014	5/15/2014	190378	2,000	44,713	46,491
2012	CLE13	ESJ01	STF	WN	3.7	Right	Green	Snout		3/15/2014	5/15/2014	190379	2,000	42,619	44,499
2012	CLE14	ESJ02	BIO	WN	3.7	Left	Green	Snout		3/15/2014	5/15/2014	190380	2,000	45,217	47,119
2012	CLE15	JCJ03	STF	WN	3.7	Right	Orange	Snout		3/15/2014	5/15/2014	190381	2,000	43,330	45,200
2012	CLE16	JCJ04	BIO	WN	3.7	Left	Orange	Snout		3/15/2014	5/15/2014	190382	2,000	42,900	44,729
2012	CLE17	JCJ05	STF	WN	3.7	Right	Orange	Snout		3/15/2014	5/15/2014	190383	2,000	43,240	45,034
2012	CLE18	JCJ06	BIO	WN	3.7	Left	Orange	Snout		3/15/2014	5/15/2014	190384	2,000	43,257	45,041

¹ BIO = BioVita (BioOregon Protein Inc.) or control diet; STF = salt-water transition diet at acclimation sites. All fish are progeny of wild/natural parents unless denoted as HC which designates the hatchery control line beginning with brood year 2002. "Avg BKD" denotes the average BKD ELISA ranking of the female parents whose progeny were in these ponds.

² The number of fish released is estimated as the total number of fish counted at marking less mortalities documented from mark to release.

Appendix A. Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.

<i>Brood Year</i>	<i>C.E. Pond</i>	<i>Accl. Pond</i>	<i>Treatment¹ /Avg BKD</i>					<i>Tag Information</i>			<i>First Release</i>	<i>Last Release</i>	<i>CWT Code</i>	<i>No. PIT</i>	<i>No. CWT</i>	<i>Est. Tot. Release²</i>
2013	CLE01	CFJ05	WN	3.8	Right	Red	Snout				3/15/2015	5/6/2015	190401	2,000	36,097	37,928
2013	CLE02	CFJ06	WN	3.8	Left	Red	Snout				3/15/2015	5/6/2015	190402	2,000	34,541	36,343
2013	CLE03	ESJ05	WN	3.7	Right	Green	Snout				3/15/2015	5/6/2015	190403	2,000	33,761	35,473
2013	CLE04	ESJ06	WN	3.7	Left	Green	Snout				3/15/2015	5/6/2015	190404	2,000	34,682	36,295
2013	CLE05	CFJ03	WN	3.9	Right	Red	Snout				3/15/2015	5/6/2015	190405	2,000	34,495	36,240
2013	CLE06	CFJ04	WN	3.9	Left	Red	Snout				3/15/2015	5/6/2015	190406	2,000	32,054	33,823
2013	CLE07	ESJ03	WN	3.8	Right	Green	Snout				3/15/2015	5/6/2015	190407	2,000	32,866	34,672
2013	CLE08	ESJ04	WN	3.8	Left	Green	Snout				3/15/2015	5/6/2015	190408	2,000	34,418	36,130
2013	CLE09	CFJ01	HC	3.8	Right	Red	Posterior Dorsal				3/15/2015	5/6/2015	190409	4,000	32,264	36,029
2013	CLE10	CFJ02	HC	3.7	Left	Red	Posterior Dorsal				3/15/2015	5/6/2015	190410	4,000	31,648	35,570
2013	CLE11	JCJ03	WN	3.7	Right	Orange	Snout				3/15/2015	5/6/2015	190411	2,000	34,948	36,725
2013	CLE12	JCJ04	WN	3.7	Left	Orange	Snout				3/15/2015	5/6/2015	190412	2,000	35,508	37,236
2013	CLE13	ESJ01	WN	3.6	Right	Green	Snout				3/15/2015	5/6/2015	190413	2,000	34,013	35,805
2013	CLE14	ESJ02	WN	3.6	Left	Green	Snout				3/15/2015	5/6/2015	190414	2,000	34,580	36,370
2013	CLE15	JCJ01	WN	3.7	Right	Orange	Snout				3/15/2015	5/6/2015	190415	2,000	32,151	33,810
2013	CLE16	JCJ02	WN	3.7	Left	Orange	Snout				3/15/2015	5/6/2015	190416	2,000	33,703	35,249
2013	CLE17	JCJ05	WN	3.8	Right	Orange	Snout				3/15/2015	5/6/2015	190417	2,000	35,987	37,604
2013	CLE18	JCJ06	WN	3.8	Left	Orange	Snout				3/15/2015	5/6/2015	190418	2,000	33,807	35,453

¹ All fish are progeny of wild/natural parents unless denoted as HC which designates the hatchery control line beginning with brood year 2002. "Avg BKD" denotes the average BKD ELISA ranking of the female parents whose progeny were in these ponds.

² The number of fish released is estimated as the total number of fish counted at marking less mortalities documented from mark to release.

Appendix A. Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.

Brood Year	C.E. Pond	Accl. Pond	Treatment¹			Tag Information			First Release	Last Release	CWT Code	No. PIT	No. CWT	Est. Tot. Release²
2014	CLE01	JCJ01	VIT	WN	1.7	Right	Orange	Snout	3/15/2016	5/12/2016	190427	2,000	35,198	37,071
2014	CLE02	JCJ02	PRO	WN	1.7	Left	Orange	Snout	3/15/2016	5/12/2016	190428	2,000	33,966	35,853
2014	CLE03	ESJ05	VIT	WN	1.6	Right	Green	Snout	3/15/2016	5/12/2016	190429	2,000	33,202	35,121
2014	CLE04	ESJ06	PRO	WN	1.6	Left	Green	Snout	3/15/2016	5/12/2016	190430	2,000	32,271	34,191
2014	CLE05	CFJ01	VIT	WN	1.5	Right	Red	Snout	3/15/2016	5/12/2016	190431	2,000	34,849	36,728
2014	CLE06	CFJ02	PRO	WN	1.4	Left	Red	Snout	3/15/2016	5/12/2016	190432	2,000	33,272	35,097
2014	CLE07	JCJ05	VIT	WN	1.5	Right	Orange	Snout	3/15/2016	5/12/2016	190433	2,000	37,322	38,943
2014	CLE08	JCJ06	PRO	WN	1.5	Left	Orange	Snout	3/15/2016	5/12/2016	190434	2,000	36,493	38,274
2014	CLE09	CFJ03	VIT	WN	1.9	Right	Red	Snout	3/15/2016	5/12/2016	190435	2,000	36,883	38,786
2014	CLE10	CFJ04	PRO	WN	1.9	Left	Red	Snout	3/15/2016	5/12/2016	190436	2,000	34,619	36,507
2014	CLE11	JCJ03	VIT	WN	1.5	Right	Orange	Snout	3/15/2016	5/12/2016	190437	2,000	37,505	39,376
2014	CLE12	JCJ04	PRO	WN	1.5	Left	Orange	Snout	3/15/2016	5/12/2016	190438	2,000	35,212	37,016
2014	CLE13	ESJ01	VIT	WN	1.4	Right	Green	Snout	3/15/2016	5/12/2016	190439	2,000	37,387	39,279
2014	CLE14	ESJ02	PRO	WN	1.4	Left	Green	Snout	3/15/2016	5/12/2016	190440	2,000	38,002	39,894
2014	CLE15	ESJ03	VIT	WN	1.4	Right	Green	Snout	3/15/2016	5/12/2016	190441	2,000	37,749	39,146
2014	CLE16	ESJ04	PRO	WN	1.4	Left	Green	Snout	3/15/2016	5/12/2016	190442	2,000	36,736	38,626
2014	CLE17	CFJ05	VIT	HC	1.2	Right	Red	Posterior Dorsal	3/15/2016	5/12/2016	190443	4,000	40,014	43,232
2014	CLE18	CFJ06	PRO	HC	1.3	Left	Red	Posterior Dorsal	3/15/2016	5/12/2016	190444	4,000	38,272	42,090

¹ All fish are progeny of wild/natural parents unless denoted as HC which designates the hatchery control line beginning with brood year 2002. "Avg BKD" denotes the average BKD ELISA ranking of the female parents whose progeny were in these ponds. PRO=BioPro diet, VIT=BioVita diet, Bio-Oregon products.

² The number of fish released is estimated as the total number of fish counted at marking less mortalities documented from mark to release.

Appendix A. Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.

<i>Brood Year</i>	<i>C.E. Pond</i>	<i>Accl. Pond</i>	<i>Treatment¹</i>			<i>Tag Information</i>			<i>First Release</i>	<i>Last Release</i>	<i>CWT Code</i>	<i>No. PIT</i>	<i>No. CWT</i>	<i>Est. Tot. Release²</i>
2015	CLE01	ESJ01	PRO	WN	2.9	Right	Green	Snout	3/15/2017	5/15/2017	190457	2,000	32,798	34,620
2015	CLE02	ESJ02	VIT	WN	2.9	Left	Green	Snout	3/15/2017	5/15/2017	190458	2,000	32,700	34,552
2015	CLE03	JCJ03	PRO	WN	2.9	Right	Orange	Snout	3/15/2017	5/15/2017	190459	2,000	38,469	40,305
2015	CLE04	JCJ04	VIT	WN	2.9	Left	Orange	Snout	3/15/2017	5/15/2017	190460	2,000	34,615	36,415
2015	CLE05	CFJ05	PRO	WN	2.9	Right	Red	Snout	3/15/2017	5/15/2017	190461	2,000	33,149	35,007
2015	CLE06	CFJ06	VIT	WN	2.9	Left	Red	Snout	3/15/2017	5/15/2017	190462	2,000	32,516	34,357
2015	CLE07	CFJ01	PRO	HC	2.6	Right	Red	Posterior Dorsal	3/15/2017	5/15/2017	190463	4,000	28,055	31,894
2015	CLE08	CFJ02	VIT	HC	2.6	Left	Red	Posterior Dorsal	3/15/2017	5/15/2017	190464	4,000	24,464	28,317
2015	CLE09	JCJ01	PRO	WN	3.0	Right	Orange	Snout	3/15/2017	5/15/2017	190465	2,000	38,098	39,927
2015	CLE10	JCJ02	VIT	WN	3.0	Left	Orange	Snout	3/15/2017	5/15/2017	190466	2,000	35,807	37,611
2015	CLE11	ESJ03	PRO	WN	2.8	Right	Green	Snout	3/15/2017	5/15/2017	190467	2,000	33,136	34,968
2015	CLE12	ESJ04	VIT	WN	2.8	Left	Green	Snout	3/15/2017	5/15/2017	190468	2,000	34,248	36,014
2015	CLE13	ESJ05	PRO	WN	2.8	Right	Green	Snout	3/15/2017	5/15/2017	190469	2,000	37,837	39,669
2015	CLE14	ESJ06	VIT	WN	2.8	Left	Green	Snout	3/15/2017	5/15/2017	190470	2,000	36,564	38,402
2015	CLE15	JCJ05	PRO	WN	2.9	Right	Orange	Snout	3/15/2017	5/15/2017	190471	2,000	34,354	36,206
2015	CLE16	JCJ06	VIT	WN	2.9	Left	Orange	Snout	3/15/2017	5/15/2017	190472	2,000	36,156	38,019
2015	CLE17	CFJ03	PRO	WN	2.8	Right	Red	Snout	3/15/2017	5/15/2017	190473	2,000	36,915	38,720
2015	CLE18	CFJ04	VIT	WN	2.8	Left	Red	Snout	3/15/2017	5/15/2017	190474	2,000	38,105	39,944

¹ All fish are progeny of wild/natural parents unless denoted as HC which designates the hatchery control line beginning with brood year 2002. "Avg BKD" denotes the average BKD ELISA ranking of the female parents whose progeny were in these ponds. PRO=BioPro diet, VIT=BioVita diet, Bio-Oregon products.

² The number of fish released is estimated as the total number of fish counted at marking less mortalities documented from mark to release.

Appendix A. Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.

<i>Brood Year</i>	<i>C.E. Pond</i>	<i>Accl. Pond</i>	<i>Treatment¹</i>	<i>Tag Information</i>					<i>First Release</i>	<i>Last Release</i>	<i>CWT Code</i>	<i>No. PIT</i>	<i>No. CWT</i>	<i>Est. Tot. Release²</i>
2016	CLE01	CFJ05	PRO WN	2.4	Right	Red	Snout	3/15/2018	5/15/2018	190490	2,000	35,447	37,354	
2016	CLE02	CFJ06	VIT WN	2.4	Left	Red	Snout	3/15/2018	5/15/2018	190491	2,000	35,568	37,468	
2016	CLE03	ESJ05	PRO WN	2.4	Right	Green	Snout	3/15/2018	5/15/2018	190492	2,000	36,330	38,195	
2016	CLE04	ESJ06	VIT WN	2.4	Left	Green	Snout	3/15/2018	5/15/2018	190493	2,000	35,002	36,943	
2016	CLE05	CFJ01	PRO HC	2.7	Right	Red	Posterior Dorsal	3/15/2018	5/15/2018	190494	4,000	36,189	40,043	
2016	CLE06	CFJ02	VIT HC	2.7	Left	Red	Posterior Dorsal	3/15/2018	5/15/2018	190495	4,000	37,147	41,026	
2016	CLE07	CJ03	PRO WN	2.4	Right	Orange	Snout	3/15/2018	5/15/2018	190496	2,000	36,599	38,400	
2016	CLE08	JCJ04 ³	VIT WN	2.4	Left	Orange	Snout	3/15/2018	5/15/2018	190497	2,000	34,080	54,569	
2016	CLE09	JCJ01	PRO WN	2.5	Right	Orange	Snout	3/15/2018	5/15/2018	190498	2,000	34,189	36,048	
2016	CLE10	JCJ02 ³	VIT WN	2.5	Left	Orange	Snout	3/15/2018	5/15/2018	190499	2,000	32,004	52,475	
2016	CLE11	CFJ03	PRO WN	2.6	Right	Red	Snout	3/15/2018	5/15/2018	190501	2,000	36,470	38,334	
2016	CLE12	CFJ04	VIT WN	2.6	Left	Red	Snout	3/15/2018	5/15/2018	190502	2,000	34,372	36,265	
2016	CLE13	ESJ03	PRO WN	2.5	Right	Green	Snout	3/15/2018	5/15/2018	190503	2,000	31,448	33,380	
2016	CLE14	ESJ04	VIT WN	2.5	Left	Green	Snout	3/15/2018	5/15/2018	190504	2,000	31,093	33,025	
2016	CLE15	JCJ05	PRO WN	2.5	Right	Orange	Snout	3/15/2018	5/15/2018	190505	2,000	36,688	38,550	
2016	CLE16	JCJ06 ³	VIT WN	2.5	Left	Orange	Snout	3/15/2018	5/15/2018	190506	2,000	35,244	0	
2016	CLE17	ESJ01	PRO WN	2.5	Right	Green	Snout	3/15/2018	5/15/2018	190507	2,000	37,553	39,512	
2016	CLE18	ESJ02	VIT WN	2.5	Left	Green	Snout	3/15/2018	5/15/2018	190508	2,000	35,689	37,621	

¹ All fish are progeny of wild/natural parents unless denoted as HC which designates the hatchery control line beginning with brood year 2002. "Avg BKD" denotes the average BKD ELISA ranking of the female parents whose progeny were in these ponds. PRO=BioPro diet, VIT=BioVita diet, Bio-Oregon products.

² The number of fish released is estimated as the total number of fish counted at marking less mortalities documented from mark to release.

³ Due to problems at the acclimation site, Jack Creek raceway 6 was closed and all fish transferred and split between raceways 2 and 4 in February 2018.

Appendix A. Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.

<i>Brood Year</i>	<i>C.E. Pond</i>	<i>Accl. Pond</i>	<i>Treatment¹</i>			<i>Tag Information</i>			<i>First Release</i>	<i>Last Release</i>	<i>CWT Code</i>	<i>No. PIT</i>	<i>No. CWT</i>	<i>Est. Tot. Release²</i>
2017	CLE01	CFJ01	PRO	WN	3.4	Right	Red	Snout	3/15/2019	5/9/2019	190535	2,000	38,689	40,527
2017	CLE02	CFJ02	VIT	WN	3.4	Left	Red	Snout	3/15/2019	5/9/2019	190536	2,000	39,792	41,650
2017	CLE03	ESJ05	PRO	WN	3.5	Right	Green	Snout	3/15/2019	5/9/2019	190537	2,000	34,646	36,556
2017	CLE04	ESJ06	VIT	WN	3.5	Left	Green	Snout	3/15/2019	5/9/2019	190538	2,000	35,655	37,493
2017	CLE05	JCJ05 ³	PRO	WN	3.1	Right	Orange	Snout			190539	2,000	35,118	0
2017	CLE06	JCJ06 ³	VIT	WN	3.1	Left	Orange	Snout			190540	2,000	36,475	0
2017	CLE07	ESJ03	PRO	WN	3.3	Right	Green	Snout	3/15/2019	5/9/2019	190541	2,000	37,843	39,737
2017	CLE08	ESJ04	VIT	WN	3.3	Left	Green	Snout	3/15/2019	5/9/2019	190542	2,000	38,689	40,579
2017	CLE09	CFJ03	PRO	WN	3.4	Right	Red	Snout	3/15/2019	5/9/2019	190543	2,000	40,551	42,423
2017	CLE10	CFJ04	VIT	WN	3.4	Left	Red	Snout	3/15/2019	5/9/2019	190544	2,000	41,529	43,357
2017	CLE11	JCJ03 ³	PRO	WN	3.3	Right	Orange	Snout	3/15/2019	5/7/2019	190545	2,000	38,702	58,941
2017	CLE12	JCJ04 ³	VIT	WN	3.3	Left	Orange	Snout	3/15/2019	5/7/2019	190546	2,000	39,368	60,266
2017	CLE13	ESJ01	PRO	WN	3.3	Right	Green	Snout	3/15/2019	5/9/2019	190547	2,000	37,502	39,385
2017	CLE14	ESJ02	VIT	WN	3.3	Left	Green	Snout	3/15/2019	5/9/2019	190548	2,000	37,829	39,699
2017	CLE15	CFJ05	PRO	HC	3.2	Right	Red	Posterior Dorsal	3/15/2019	5/9/2019	190549	4,000	33,390	37,153
2017	CLE16	CFJ06	VIT	HC	3.2	Left	Red	Posterior Dorsal	3/15/2019	5/9/2019	190550	4,000	35,413	39,126
2017	CLE17	JCJ01 ³	PRO	WN	3.3	Right	Orange	Snout	3/15/2019	5/7/2019	190551	2,000	36,661	56,934
2017	CLE18	JCJ02 ³	VIT	WN	3.3	Left	Orange	Snout	3/15/2019	5/7/2019	190552	2,000	35,946	56,843

¹ All fish are progeny of wild/natural parents unless denoted as HC which designates the hatchery control line beginning with brood year 2002. "Avg BKD" denotes the average BKD ELISA ranking of the female parents whose progeny were in these ponds. PRO=BioPro diet, VIT=BioVita diet, Bio-Oregon products.

² The number of fish released is estimated as the total number of fish counted at marking less mortalities documented from mark to release.

³ Due to problems at the acclimation site, Jack Creek raceways 5&6 were closed and all fish transferred and split between raceways 1-4 in February 2019.

Appendix A. Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.

<i>Brood</i>	<i>C.E.</i>	<i>Accl.</i>	<i>Treatment¹</i>				<i>Tag Information</i>				<i>First</i>	<i>Last</i>	<i>CWT</i>	<i>No.</i>	<i>No.</i>	<i>Est. Tot.</i>
<i>Pond</i>					<i>/Avg BKD</i>					<i>Release</i>	<i>Release</i>	<i>Code</i>	<i>PIT</i>	<i>CWT</i>	<i>Release²</i>	
2018	CLE01	ESJ01	Pro	WN	4.2	Left	Green	Snout		3/15/2020	5/15/2020	190573		2,773	31,833	34,524
2018	CLE02	ESJ02	Vit	WN	4.2	Right	Green	Snout		3/15/2020	5/15/2020	190574		2,000	31,213	33,105
2018	CLE03	CFJ01	Pro	HC	3.2	Left	Red	Posterior Dorsal		3/15/2020	5/15/2020	190575		2,000	35,285	37,228
2018	CLE04	CFJ02	Vit	HC	3.2	Right	Red	Posterior Dorsal		3/15/2020	5/15/2020	190576		2,000	34,672	36,594
2018	CLE05	ESJ03	Pro	WN	4.0	Left	Green	Snout		3/15/2020	5/15/2020	190577		2,000	33,397	35,301
2018	CLE06	ESJ04	Vit	WN	4.0	Right	Green	Snout		3/15/2020	5/15/2020	190578		2,000	33,772	35,692
2018	CLE07	CFJ05	Pro	HC	3.1	Left	Red	Posterior Dorsal		3/15/2020	5/15/2020	190579		2,000	32,461	34,384
2018	CLE08	CFJ06	Vit	HC	3.1	Right	Red	Posterior Dorsal		3/15/2020	5/15/2020	190580		2,000	34,276	36,203
2018	CLE09	JCJ03	Pro	WN	3.9	Left	Orange	Snout		3/15/2020	5/15/2020	190581		2,000	39,166	41,015
2018	CLE10	JCJ04	Vit	WN	3.9	Right	Orange	Snout		3/15/2020	5/15/2020	190582		2,000	38,910	40,780
2018	CLE11	JCJ05	Pro	WN	4.2	Left	Orange	Snout		3/15/2020	5/15/2020	190583		2,000	32,561	34,449
2018	CLE12	JCJ06	Vit	WN	4.2	Right	Orange	Snout		3/15/2020	5/15/2020	190584		2,000	32,726	34,621
2018	CLE13	JCJ01	Pro	WN	3.2	Left	Orange	Snout		3/15/2020	5/15/2020	190585		2,000	34,595	36,473
2018	CLE14	JCJ02	Vit	WN	3.2	Right	Orange	Snout		3/15/2020	5/15/2020	190586		2,000	32,739	34,630
2018	CLE15	CFJ04	Pro	WN	4.1	Left	Red	Snout		3/15/2020	5/15/2020	190587		4,000	30,681	34,579
2018	CLE16	CFJ03	Vit	WN	4.1	Right	Red	Snout		3/15/2020	5/15/2020	190588		4,000	30,934	34,845
2018	CLE17	ESJ05	Pro	WN	4.0	Left	Green	Snout		3/15/2020	5/15/2020	190589		2,000	32,347	34,266
2018	CLE18	ESJ06	Vit	WN	4.0	Right	Green	Snout		3/15/2020	5/15/2020	190590		2,000	31,802	33,731

¹ All fish are progeny of wild/natural parents unless denoted as HC which designates the hatchery control line beginning with brood year 2002. "Avg BKD" denotes the average BKD ELISA ranking of the female parents whose progeny were in these ponds. PRO=BioPro diet, VIT=BioVita diet, Bio-Oregon products.

² The number of fish released is estimated as the total number of fish counted at marking less mortalities documented from mark to release.

Appendix A. Tag and Release Information by Cle Elum Pond Id, Brood Years 2006-2019.

<i>Brood</i>	<i>C.E.</i>	<i>Accl.</i>	<i>Treatment¹</i>			<i>Tag Information</i>			<i>First</i>	<i>Last</i>	<i>CWT</i>	<i>No.</i>	<i>No.</i>	<i>Est. Tot.</i>
<i>Pond</i>					<i>/Avg BKD</i>				<i>Release</i>	<i>Release</i>	<i>Code</i>	<i>PIT</i>	<i>CWT</i>	<i>Release²</i>
2019	CLE01	ESJ05	VIT	WN	3.8	Left	Green	Snout	3/15/2021	5/13/2021	190632	2,000	33,560	35,472
2019	CLE02	ESJ06	PRO	WN	3.8	Right	Green	Snout	3/15/2021	5/13/2021	190631	2,000	30,989	32,896
2019	CLE03	CFJ01	VIT	HC	3.6	Left	Red	Posterior Dorsal	3/15/2021	5/13/2021	190630	2,000	28,346	30,283
2019	CLE04	CFJ02	PRO	HC	3.6	Right	Red	Posterior Dorsal	3/15/2021	5/13/2021	190629	2,000	26,327	28,236
2019	CLE05	JCJ05	VIT	WN	3.4	Left	Orange	Snout	3/15/2021	5/13/2021	190628	2,000	30,806	32,703
2019	CLE06	JCJ06	PRO	WN	3.4	Right	Orange	Snout	3/15/2021	5/13/2021	190627	2,000	32,103	33,984
2019	CLE07	ESJ03	VIT	WN	3.6	Left	Green	Snout	3/15/2021	5/13/2021	190626	2,000	33,106	34,985
2019	CLE08	ESJ04	PRO	WN	3.6	Right	Green	Snout	3/15/2021	5/13/2021	190625	2,000	31,724	33,590
2019	CLE09	JCJ03	VIT	WN	3.7	Left	Orange	Snout	3/15/2021	5/13/2021	190624	2,000	33,462	35,333
2019	CLE10	JCJ04	PRO	WN	3.7	Right	Orange	Snout	3/15/2021	5/13/2021	190623	2,000	34,274	36,137
2019	CLE11	CFJ03	VIT	WN	3.9	Left	Red	Snout	3/15/2021	5/13/2021	190622	4,000	22,653	26,457
2019	CLE12	CFJ04	PRO	WN	3.9	Right	Red	Snout	3/15/2021	5/13/2021	190621	4,000	23,275	27,097
2019	CLE13	JCJ01	VIT	WN	3.5	Left	Orange	Snout	3/15/2021	5/13/2021	190620	2,000	33,085	34,904
2019	CLE14	JCJ02	PRO	WN	3.5	Right	Orange	Snout	3/15/2021	5/13/2021	190619	2,000	28,839	30,720
2019	CLE15	CFJ05	VIT	HC	3.9	Left	Red	Posterior Dorsal	3/15/2021	5/13/2021	190618	2,000	19,755	21,678
2019	CLE16	CFJ06	PRO	HC	3.9	Right	Red	Posterior Dorsal	3/15/2021	5/13/2021	190617	2,000	17,875	19,824
2019	CLE17	ESJ01	VIT	WN	3.7	Left	Green	Snout	3/15/2021	5/13/2021	190616	2,000	26,511	28,341
2019	CLE18	ESJ02	PRO	WN	3.7	Right	Green	Snout	3/15/2021	5/13/2021	190615	2,000	26,240	27,758

¹ All fish are progeny of wild/natural parents unless denoted as HC which designates the hatchery control line beginning with brood year 2002. "Avg BKD" denotes the average BKD ELISA ranking of the female parents whose progeny were in these ponds. PRO=BioPro diet, VIT=BioVita diet, Bio-Oregon products.

² The number of fish released is estimated as the total number of fish counted at marking less mortalities documented from mark to release.

Appendix C

2020 Annual Chandler Certification for Outmigrating Spring (Yearling) Chinook Smolts



Photo courtesy of Dr. Dave Fast, Yakima

Prepared by:

Shubha Pandit, Bill Bosch & Mark Johnston
Yakima-Klickitat Fisheries Project
Yakama Nation Fisheries
P.O. Box 151, Toppenish,
WA 98948, USA

July 14, 2021

<u>EXECUTIVE SUMMARY</u>	<u>3</u>
<u>1. INTRODUCTION</u>	<u>6</u>
<u>2.0 METHODOLOGY.....</u>	<u>9</u>
2.1. ESTIMATING SAMPLE RATE AND CALIBRATION	10
2.2. MISSING DATA IMPUTATION	13
2.3. PIT-TAG DATA	13
2.4. GENETIC INFORMATION	13
2.5. ESTIMATING PROSSER BYPASS DETECTION RATE.....	14
2.6. WILD AND HATCHERY PASSAGE ESTIMATE	15
2.7. MODEL VALIDATION (ESTIMATES COMPARISONS)	16
2.8. ESTIMATED DAILY SMOLT OUTMIGRATION FROM PROSSER	16
<u>3.0 RESULTS AND DISCUSSION.....</u>	<u>18</u>
3.1. SPECIES COMPOSITION AND DAILY COUNTS IN THE COUNTING FACILITY.....	19
3.2. COUNTS OF WILD AND HATCHERY SPRING CHINOOK.....	22
3.3. DETECTION RATE OF THE SAMPLING FACILITY AND DOWNSTREAM DAMS (MCJ, JOHN DAY, BON).....	24
3.4. PREDICTED NUMBER OF OUTMIGRATING WILD AND HATCHERY SPRING CHINOOK SMOLTS	26
3.5. ANNUAL TREND OF JUVENILE PROSSER-PASSAGE ESTIMATES (HATCHERY AND WILD) BY STOCK	27
3.6. GENETIC VARIATION AMONG STOCKS (UPPER YAKIMA, NACHES, AMERICAN).....	32
3.7. CONTRIBUTION OF EACH STOCK TO OUTMIGRATION	34
3.9. RELATIONSHIP BETWEEN WILD JUVENILE PASSAGE ESTIMATES AND ESTIMATED ADULT RETURNS.....	38
3.9. RELATIONSHIP BETWEEN ESTIMATED JUVENILE PASSAGE AND RIVER FLOW.....	40
3.9.1. ANNUAL.....	40
3.9.2. DAILY.....	41
<u>4. REFERENCES</u>	<u>44</u>
<u>5. SUPPLEMENTARY INFORMATION: DETAILED PASSAGE-ESTIMATES</u>	<u>45</u>

Executive Summary

Outmigrating smolts have been monitored since 1983 at the Chandler Diversion Canal in the Yakima River at Prosser, Washington. Chandler Juvenile Monitoring Facility (CJMF) improvements over the years have made it possible to count all species entering the juvenile bypass system each year from January into July, encompassing the entire juvenile outmigration period. Winter operations are made possible by the dual purpose of the canal, which supplies a hydroelectric plant as well as an irrigation district. The diversion is located downstream from all Spring Chinook, Summer Chinook, Coho and Steelhead spawning and juvenile rearing areas in the Yakima River Basin.

Numerous projects to restore and protect channel and riparian habitat, along with fish reintroduction programs, have been implemented in the Yakima Basin since the 1990s. The population status and trends for the different species in their freshwater life stages are important measures of management success, and the data collected at the facility have allowed us to answer several management questions that can help to improve these programs. This report provides estimates of 2020 outmigrating Spring (yearling) Chinook smolt populations (hatchery and wild) past Prosser Dam; its temporal (annual) trend from 1999 through 2020; and evaluation of whether the production and releases of hatchery smolts into the upper Yakima had an effect on the production of wild smolts and on the relative abundance of the three stock sources of wild smolts (Naches, American, and Upper Yakima rivers). This evaluation is part of an ongoing study that was initiated in 1999 with the first release of hatchery Spring Chinook smolts.

The entire bypass flow leaving the juvenile screens enters the counting facility but only a portion is manually counted. A timer gate on an hourly cycle directs bypass flow to a holding tank for a portion of each hour that can be adjusted as often as once per day to compensate for fluctuations in fish abundance so as to not overwhelm the capacity of the staff to tally those smolts by species and stock.

In 2020, the CJMF was in operation from January 15th to July 8th, a total of 178 days, with occasional closures (4 days) due to high stream flows or adverse river conditions. There were three gate timing settings (TR) for fish sampling. Over the 178 operating/sampling days, the timer gate was set at a 33% sample rate (20 minutes per hour) for 144 days, 50% for 9 days, and 100% for 20 days.

Several statistical methods/approaches were applied for expanding the subsample data and analyzing them. Most of the methods described in this report were based on the methods described in previous years' annual reports. To address the objectives of the study, we answered the following research questions:

1. Which species and runs were captured during the 2020 sampling period and what were the relative abundances of each group?

During the 2020 sampling period, a total of 97,447 individuals (raw, unadjusted value) of 18 species and runs were captured in the sampling room (Table 2, Figure 5). Among salmonids, hatchery and wild Spring Chinook (56,669) comprised more than 50% of the total count; and the

second highest count was Steelhead (4946). Total counts varied over time, with 5% of the total pre-March, 2% in March, 47% in April, 46% in May and 0.5% post-May.

2. What was the PIT-tag detection efficiency of the monitoring facility and did the efficiencies vary among the sampling periods (pre-March, March, April, May, Post-May)?

The overall PIT-tag detection rate at the CJMF in 2020 was 43.96 ± 1.0 %, which was higher than in 2019 (27.85 ± 0.07 %, mean \pm SE), but the rate varied through the season. The highest detection rates occurred in April and May as diversion rate increased with decreasing river flow.

3. How many wild and hatchery Spring Chinook smolts were estimated to pass Prosser Dam during 2020 and was there any temporal trend from the 1999 through 2020 juvenile migration years?

Wild (natural-origin) spring chinook can be separated genetically into three stocks: Upper Yakima, from the Yakima River and tributaries above the Naches River confluence; American River, a tributary of the Naches River; and Naches River, from the Naches River and tributaries exclusive of the American River. Only the Upper Yakima stock receives hatchery supplementation.

The estimated number of wild Spring Chinook smolts passing Prosser Dam during the 2020 migration period ranged from 115,300 to 201,313; whereas hatchery smolts ranged from 371,069 to 500,195. The estimated total number of hatchery Spring Chinook smolts passing Prosser Dam during the 2020 sampling period was almost double that of wild Spring Chinook smolts. On average over outmigration years 2000-2020, $226,910 \pm 25,682$ wild and $322,679 \pm 15,931$ hatchery Spring Chinook smolts passed Prosser Dam each year. The total number of wild outmigrating smolts as well as the upper Yakima component stock seemed to be decreasing over time, whereas the population of Upper Yakima hatchery smolts seemed to be increasing but none of these trends were statistically significant.

4. What proportions of wild Spring Chinook populations that outmigrated from Prosser were contributed by different stocks (Naches, American, Upper Yakima) in the Yakima Basin? Did the proportions of these stocks in the outmigrating smolt population vary by migration year?

About 61% of the total count of wild outmigrating Spring Chinook smolts was contributed by the Upper Yakima stock; while 28% and 11% of the total wild outmigration were contributed by the Naches and American river stocks, respectively. The rate of decline in the wild Upper Yakima stock averaged $-2498/\text{year}$, which was the highest of the three wild populations (Naches, American, Upper Yakima), but the estimated decline was not significant (Upper Yakima; $R^2=0.02$, $p=0.47$). The rate of decrease for Naches stock was $-394/\text{year}$, it was also not significant; however, only the American stock average reduction was significant (Slope= $-1087/\text{year}$, $R^2=0.228$, $p=0.04$).

5. Did the production and release of hatchery smolts into the upper Yakima affect the production of wild smolts?

To evaluate whether the hatchery program affected wild production, we tested a hypothesis that the rate of decline of outmigration should be higher in Upper Yakima wild Spring Chinook, because only the Upper Yakima stock receives hatchery supplementation. We found that there was no significant negative linear trend in the relative proportions of the three stocks of outmigrating smolt populations with the outmigration year, indicating that there was no influence of hatchery supplementation on wild abundance as measured by outmigrating smolt abundance at Prosser in the lower Yakima River. If the proportion of wild Upper Yakima smolts would have decreased significantly over time, this could represent a hatchery effect, environmental effects, or a combination of the two.

6. What was the effect of river flow (daily as well as annual flow) on the number of outmigrating Spring Chinook smolts?

The annual juvenile passage estimate of wild and hatchery Spring Chinook at Prosser Dam tends to increase with average annual river flow for the outmigration period, but this relationship was significant only for Upper Yakima hatchery smolts. The results further showed that daily estimated outmigrating smolts increased when flow pulses occurred, whether due to natural runoff or reservoir releases made to facilitate outmigration.

1. Introduction

Conservation and management of culturally and economically important species rely on monitoring programs to provide accurate and robust estimates of population size. Numerous projects to restore and protect channel and riparian habitat have been implemented on the Yakima River in coordination with reintroduction/supplementation programs. Quantifying and understanding whether juvenile outmigration or Smolt-to-Adult-Return (SAR) are increased/decreased over time, or which stocks perform better, are fundamental questions in determining whether species management and production goals are being reached.

Outmigrating smolts have been monitored since 1983 at the Chandler Diversion Canal in the Yakima River at Prosser, Washington (Figures 2 -4). The diversion canal is located downstream from all Spring Chinook, Summer Chinook, Coho and Steelhead spawning and juvenile rearing areas in the Yakima River Basin. Improvements at the Chandler Juvenile Monitoring Facility (CJMF) over the years have made it possible to count all species entering the juvenile bypass system each year from January into July, encompassing the entire juvenile (smolt) outmigration period. Winter operations are made possible by the dual purpose of the canal, which supplies a hydroelectric plant as well as an irrigation district. Chandler Diversion canal typically conveys 1000 cfs with a maximum of 1500 cfs over the course of a year. Water not used for irrigation is returned to the Yakima River eleven miles downstream at the Chandler Powerhouse. The Yakima River at Prosser is characterized by a high spring runoff peaking in March, and low summer flows reaching a minimum in August, but there is wide variation in this flow pattern and the timing of high and low flows from year to year.

At the CJMF, fish are counted from the portion of river flow that is diverted into the irrigation canal and then into the juvenile fish bypass system. The monitoring data collected at the facility over the 6-month outmigration period can be useful to determine the status and trends of different species and runs at the outmigrating smolt stage, identify potential life-cycle bottlenecks, and evaluate the effectiveness of ongoing reintroduction and habitat improvement actions on population dynamics. The number of smolts of different species that outmigrate from the river basin are influenced by the numbers and fecundity of spawners and by the conditions their progeny encounter before and during outmigration, including river water temperature and river flows. Yakima River flow is modified by storage and releases from five large reservoirs in

YKFP Project Year 2020 M&E Annual Report, Appendix C, Chandler Certification

the upper Yakima Basin, and by irrigation and hydropower withdrawals and return flow. Under various agreements, minimum flows below storage and diversion dams are maintained to sustain ecological processes during periods of low natural runoff. Snowmelt exacerbated by occasional rain-on-snow events causes considerable variation in the flow of unregulated tributaries and in the Yakima River itself from November through June. When irrigation demand exceeds this runoff during the fish outmigration period, unnatural delays and poor outmigration survival can result. Studies of the relationship of river flow and outmigration have shown that river flow pulses from natural events and reservoir releases can accelerate smolt movement downstream and enhance survival to the ocean. Relying entirely on annual outmigration totals may obscure the role of in-season flow fluctuations and the importance of flow pulses during this critical period.

The main objectives of the study were to estimate prior-year (2020) outmigrating smolt populations (hatchery and wild) of spring Chinook; assess its temporal trend from 1999 through 2020; determine whether the production and releases of hatchery smolts into the upper Yakima had an effect on the production of wild smolts and on the relative abundances of the three stock sources of wild smolts (Naches, American, and Upper Yakima rivers); evaluate whether outmigration is higher in years of high river flow; and within years, on days with greater flow. To address the objectives, we answered the following research questions:

- Which species and runs were captured during the 2020 sampling period and what were the relative abundances of each group?
- What was the PIT-tag detection efficiency of the monitoring facility, and did the efficiencies vary among the sampling periods (pre-March, March, April, May, Post-May) in 2020?
- How many wild and hatchery Spring Chinook smolts emigrated from Prosser during 2020 and was there any temporal trend from 1999 through the 2020 juvenile migration year?
- What proportions of wild Spring Chinook populations that outmigrated from Prosser were contributed by different stocks (Naches, American, Upper Yakima) in the Yakima Basin? Did the proportions of these stocks in the outmigrating smolt population vary by migration year?

- Did the production and release of hatchery smolts into the upper Yakima affect the production of wild smolts?
- What was the effect of river flow (daily as well as annual flow) on the number of outmigrating Spring Chinook smolts?

2.0 Methodology

The CJMF is located on the fish bypass outlet of Chandler Canal at Prosser Dam (Figure 1), which is about 76 river km (47 river miles) upstream from the mouth of the Yakima River. The canal supplies water for irrigation and to generate power. The Chandler Canal typically conveys 1000 cfs with a maximum of 1500 cfs over the course of a year (Pyper and Smith, 2005). The proportion of river flow diverted, and thus the proportion of smolts entrained, varies widely during the outmigration season, due mostly to fluctuations in river flow. Juvenile fish screens (Figure 2) allow fish to exit the canal. The bypass flow enters a juvenile counting facility before returning to the river, where a portion of the fish are manually counted. A timer gate on an hourly cycle directs bypass flow to a holding tank for a portion of each hour that can be adjusted as often as once per day to compensate for fluctuations in fish abundance and avoid overwhelming the capacity of the staff to tally those smolts by species and stock. For this study, several methods were used to enumerate smolts and are outlined in Figure 3.




Figure 1. Yakima basin and the location of the Chandler Juvenile Monitoring Facility at Prosser and different sub-basins or genetic stocks (Naches, Upper Yakima River and American River).



Figure 2. Composite photo depicting the Chandler canal location and the key sampling components at the Chandler Juvenile Monitoring Facility.

2.1. Estimating Sample Rate and Calibration

Figure 4 is a schematic of the CJMF layout and the details of the sampling area. The sampling period was continuous from January 15th to July 8th in 2020 except for a few days in which the facility was shut down due to adverse river conditions. In 2019, the sampling period was from January 9th to July 6th.




Figure 3. Outline of the methodology used for data analysis in this report

In 2020, three timer-gate settings (TR) were used to control the proportion of bypassed smolts that were manually counted: 33% (20 minutes per hour), 50% (30 minutes per hour), and 100%. There are two PIT-tag detectors in the bypass system (Figure 4): one upstream of the timer gate and one in the exit from the counting facility downstream of the timer gate where the daily subsamples of smolts are tallied. Along with detectors in the Prosser adult ladders, these detectors comprise site PRO in the PIT Tag Information System (PTAGIS) maintained by the Pacific States Marine Fisheries Commission.




Figure 4. Site Overview of Chandler Juvenile Monitoring Facility at Prosser. The layout was adapted from the site configuration at <https://www.ptagis.org/>.

The timer gate, when opened, directs the Prosser bypass flow from Chandler Canal into the sample tank where smolts are tallied. Data regarding species, life stage, and abundance were tallied and counted daily during the sampling period. The timer gate setting has to be corrected because some bypassed fish swim against the bypass flow and may not enter the counting facility in strict proportion to the gate setting. For a given daily TR setting, the observed sample rate was computed as:

$$SR_{ti} = \frac{\text{the number of PIT-tagged Spring Chinook smolts detected leaving the counting facility}}{\text{the total number detected by the bypass detector located upstream of the timer gate } (TG_i)}; \text{ or}$$

$$SR_{ti} = \frac{n[\text{counting facility}]}{n[\text{bypass (TR)}]}; \text{ Where } ti \text{ is the timer setting.}$$

Once we estimated the daily sample rate, the calibration value was computed as:

$$\text{Calibration value (CV)} = w(33\%) \times [SR(TR=33\%)/33\%] + w(50\%) \times [SR(TR=50\%)/50\%]$$

Where $w(33\%)$ and $w(50\%)$ are the weight, which are the proportion of bypass detections within the TR setting 0.33 and 0.50, respectively. The weights being the proportions of bypass detections within the TR setting and estimated as (see, Neeley 2012):

$$w(33\%) = \frac{n[\text{bypass}(\text{TR}=33\%)]}{\{n[\text{bypass}(\text{TR}=33\%)] + n[\text{bypass}(\text{TR}=50\%)]\}}$$

$$w(50\%) = \frac{n[\text{bypass}(\text{TR}=50\%)]}{\{n[\text{bypass}(\text{TR}=33\%)] + n[\text{bypass}(\text{TR}=50\%)]\}}$$

2.2. Missing data imputation

Spring Chinook smolts were tallied each day as to source (hatchery-spawned or wild) on the basis of external marks. However, the sampling facility was shut down for a few days due to flow conditions or other technical problems. Data were missing for those days in which the sampling facility was closed. Linear interpolation was used to impute counts for days with missing information.

2.3. PIT-tag data

We queried the PTAGIS database (<https://www.ptagis.org/>) in April 2021 to retrieve available PIT-tag detection information for all tagged hatchery Spring Chinook smolts released upstream from Prosser Dam. Altogether 43,130 hatchery Spring Chinook were tagged, (about 6% of the total release) but not all the tagged fish were detected at the acclimation site exits, either because of mortality and tag shedding over the 3-to-5-month period between tagging and volitional release, or detection failure on exit. We used only those fish which were detected on exit from acclimation sites or captured, tagged and released in the Roza Dam bypass in the upper Yakima River. A total of 35,522 PIT-tagged smolts were used for this analysis. An encounter history for each fish with detection events (date and detection site) was constructed for further analysis.

2.4. Genetic information

During the sampling period each year, tissue samples were taken from subsamples of wild smolts passing through the counting facility. In order to minimize bias, samples of smolts were distributed proportionally among five time strata (January-February, March, April, May and

June). These tissue samples were processed in the Molecular Genetics Laboratory of the Washington Department of Fish and Wildlife (WDFW). Results of 2019 and 2020 molecular samples are available (Seamons and Bowman, 2020) and these information was used to estimate 2019 and 2020 outmigrating smolts.

2.5. Estimating Prosser bypass detection rate

The proportions of all PIT- tagged smolts released above Prosser and detected at mid-Columbia dams that were previously detected in the Chandler Canal bypass serve as estimates of bypass-detection rate. Detections at the three downstream sites with juvenile PIT tag detection (McNary, John Day, and Bonneville dams) were pooled to estimate the Prosser bypass detection rate.

Daily estimates of Prosser detection rate from downstream dams are not possible because smolts migrate at different rates between Prosser and downstream dams, and one day's detections in the Prosser bypass are detected at a given downstream dam over several subsequent days. For this study, the detection rate was estimated for five strata over the outmigration period (pre-March, March, April, May and post-May) based on McNary Dam alone, or pooled over the three Columbia River dams. The detection efficiency (DE) was estimated as:

$$DE = n(\text{daily joint site detections})/n(\text{total site detections})$$

These detection rates based on upper Yakima hatchery Spring Chinook were also applied to the three stocks of wild Spring Chinook smolts, few of which were tagged. The wild Spring Chinook were made up of Naches, American, and Upper-Yakima stock (See fig. 1). All hatchery Spring Chinook smolts were coded-wire tagged and most were elastomer tagged in addition to about 6% being PIT-tagged. Elastomer tags allowed visual separation of hatchery smolts and adults by acclimation site, with fish released from the Clark Flat, Easton, and Jack Creek sites, receiving red, green, and orange elastomer tags, respectively. Elastomer-tagged smolts were also tallied by elastomer color. PIT-tagged hatchery smolts were not elastomer-tagged.

The wild and elastomer-tagged hatchery tallies were expanded by four different estimates of Prosser detection rates as mentioned above.

1. McNary-based un-stratified detection rate estimate

2. McNary-based stratified detection rate estimate
3. Pooled-lower-dam-based un-stratified detection rate estimate
4. Pooled-lower-dam-based stratified detection rate estimate

Detailed methodology is given in Neeley (2019).

Of these four estimators, the one chosen for further analysis was a pooling of stratified estimates from the detection efficiencies from McNary, John Day, and Bonneville Dams on the Columbia Rivers; the strata being established for each of these dams by combining daily estimates that were deemed similar using Logistic stepwise regression of the daily detection efficiencies on Julian-date indicators that take the value 1 if the estimate was from a given date or a later date or 0 if the estimate was from an earlier date (see, Neeley (2019) for further details).

2.6. Wild and hatchery passage estimate

On a daily basis the sampled Spring Chinook smolts were tallied as to source (hatchery-spawned or wild). On those days when the facility was shut down, linear interpolation was used to impute values to the missing information as mentioned above. The daily actual and imputed tallies were divided by the sample rates in use on those days (SR). The sample-rate-adjusted tallies for each source were added over days within each of five time periods and were then divided by the respective period's detection rate. The wild and hatchery smolts were tallied separately. Wild smolts were identified by the lack of a coded-wire tag or external mark. Hatchery smolts could be identified by the presence of an elastomer tag, a coded wire tag, an adipose fin clip and a PIT tag if there was no elastomer tag. Expanded elastomer-tagged tallies were then divided by the proportion of hatchery smolts to obtain estimates of the passage of all hatchery smolts.

Within each of the five time periods (pre-March, March, April, May, post-May), the tallied sample of wild smolts was subsampled and genetically classified as to brood origin (American, Naches, or Upper Yakima rivers) by the Washington Department of Fish and Wildlife Molecular Genetics Laboratory so that brood-origin proportions could be estimated for each stratum. The wild passage estimates within each period were multiplied by each of the period's brood-source proportions. Each wild brood's time-period passage estimates were then added over the time

periods to estimate the brood's total passage, as were the hatchery passage estimates. The detailed methodology can be found in Neeley (2019).

2.7. Model validation (estimates comparisons)

The estimates of the number of smolts passing Prosser Dam can vary slightly with different entrainment-based estimation methods. To ascertain which of these passage estimates is the best to report and use for further analysis, we compared flow/entrainment-based estimates of hatchery Spring Chinook smolts at Prosser to another estimate that was derived using a PIT-tag-based survival rate from release site to Prosser Dam. Since we know the total number of hatchery Spring Chinook smolts released in the upper Yakima, we multiplied the survival rate by the total release, which provided the total hatchery smolt population passing Prosser. This estimate can be viewed as an independent estimate but it can also be biased because we assumed there was no variation in the survival rate among the sampling days time strata. If detection rate is not homogeneous, survival rate cannot be homogeneous. However, this survival-based estimator has value because it is independent of the flow/entrainment-based method.

In addition to the survival-based method, each of the flow/entrainment methods' estimates of hatchery juvenile passage (see section 2.5 above) was also compared with hatchery adult returns at Prosser (Bosch, 2020). If the estimate is a reasonable value, it should be highly correlated with the hatchery adult returns from that outmigration.

2.8. Estimated Daily smolt outmigration from Prosser

One of our objectives was to determine whether river flows influence the size of the population of outmigrating smolts. If larger number of smolts outmigrated during high river flow, the rate of outmigration would be a function of river flow. To estimate daily passage at Prosser Dam, daily counts of each species in the live box at the (CJMF) were expanded by the canal entrainment, canal survival (from prior paired releases), and sub-sampling rates using the following formula (Neeley, 2012).

$$\text{Entrainment rate (ER)} = 1/1 + \exp(-5.60081 + 13.5861 * \text{diversion rate}) \text{..eq. 1}$$

$$\text{Survival Probability} = 1/1 + \exp(-2.84815 + 0.0154 * \text{Julian date} - 0.00017 * (\text{canal flow} + 132)) \text{..eq.2}$$

Estimated daily count: $\text{Count}/(\text{Survival Probability} * \text{sample.rate}(\text{SR}) * \text{ER})$.. eq.3

The model for the Entrainment Rate (ER) was based on the logistic regression using the daily proportion of Yakima River flow diverted into the canal. The Entrainment Rate (ER) is the predicted daily proportion of fish passing Prosser that are entrained into Chandler Canal, the Canal-Survival Rate (Survival probability) is the daily predicted proportion of those entrained fish that survive the canal from below the head-gate down the canal and into the bypass to a point just above the sampling station, and Sampling Rate (SR) is the estimated proportion of fish that are sampled from the bypass and enumerated.

2.8.1. Relationship between river flow and estimated daily count

To determine whether high river flow helped to increase the rate of smolt outmigration from Prosser, we built univariate relationships using two datasets (annual and daily).

- A. Annual total estimates: A univariate linear relationship between the estimated total annual number of hatchery Spring Chinook smolts passing Prosser (2000-2020 outmigration years) and the average March-June river flows (corresponding to the March-June volitional exit of hatchery Spring Chinook from acclimation sites) for each year from 2000 through 2019.
- B. Daily estimates: A univariate linear relationship between the estimated daily count of wild Spring Chinook and daily river flow above Prosser Dam, which is the sum of the daily flows measured at the Bureau of Reclamation gaging stations CHCW (Chandler Canal) and YRPW (Yakima River below Chandler Canal). River flow data were accessed in May, 2021 from

<https://www.usbr.gov/pn/hydromet/yakima/yakwebarcread.html>.

3.0 Results and discussion

In 2020 the CJMF was operated from January 15th to July 8th (178 days total), with occasional closures (4 days) due to excessively high stream flows or other technical problems. There were three timer gate settings (TR) for sampling, representing the percentage of time in each hourly cycle that bypassed fish were directed into the sample tank. Over the sampling period, the timer gate setting (TR) was 33% for 144 days, 50% for 9 days, and 100% for 20 days. As noted earlier, adjustments are applied to timer gate settings because some bypassed fish swim against the bypass flow upstream from the gate and may not enter the counting facility in strict proportion to the gate setting, unless there is no alternative, i.e. the gate is set to sample 100% of bypass flow. This occurs at the end of the season when lethal lower river conditions require transportation of entrained smolts to the Columbia River instead of discharge past the sample room detector to the Yakima River.

The SR is usually less than the TR, indicating not all fish passing through the bypass when the timer gate is open are actually entering and being detected in the counting facility. In 2020, when TR was 33%, sample rate (SR) was 26.1%, and at the 50% TR setting the SR was 39.7% (Table 1). The 2020 sample rate was lower than the rate in 2019 and 2018 but similar to the rates in 2017, 2009, 2007, 2000 and 1992 (Table 1). Why there was variation in sampling rate among years is a topic for further investigation, with potential causes including water temperature, fish size or other factors.

Table 1. Sample-room sample rates for given timer-gate settings. Timer Gate Rate (TR) is the proportion of time that the bypass gate is opened to Sample Room.

Out-Migration Year	Calibration Value	Estimated Sample Rates (SR) for different Timer-Gate Rates								
		Timer-Gate Rate (TR)								
		0.05	0.1	0.2	0.25	0.33	0.4	0.45	0.5	0.75
1998	0.778	0.039	0.078	0.156	0.194	0.257	0.311	0.350	0.389	0.583
1999	0.833	0.042	0.083	0.167	0.208	0.275	0.333	0.375	0.417	0.625
2000	0.794	0.040	0.079	0.159	0.198	0.262	0.318	0.357	0.397	0.595
2001	0.278	0.014	0.028	0.056	0.070	0.092	0.111	0.125	0.139	0.209
2002	0.838	0.042	0.084	0.168	0.209	0.277	0.335	0.377	0.419	0.628
2003	0.669	0.033	0.067	0.134	0.167	0.221	0.267	0.301	0.334	0.501
2004	0.693	0.035	0.069	0.139	0.173	0.229	0.277	0.312	0.346	0.520

2005	0.776	0.039	0.078	0.155	0.194	0.256	0.310	0.349	0.388	0.582
2006	1.000	0.050	0.100	0.200	0.250	0.330	0.400	0.450	0.500	0.750
2007	0.800	0.040	0.080	0.160	0.200	0.264	0.320	0.360	0.400	0.600
2008	0.651	0.033	0.065	0.130	0.163	0.215	0.260	0.293	0.326	0.488
2009	0.770	0.038	0.077	0.154	0.192	0.254	0.308	0.346	0.385	0.577
2010	0.584	0.029	0.058	0.117	0.146	0.193	0.234	0.263	0.292	0.438
2011	1.000	0.050	0.100	0.200	0.250	0.330	0.400	0.450	0.500	0.750
2012	0.979	0.049	0.098	0.196	0.245	0.323	0.391	0.440	0.489	0.734
2013	0.973	0.049	0.097	0.195	0.243	0.321	0.389	0.438	0.486	0.729
2014	0.903	0.045	0.090	0.181	0.226	0.298	0.361	0.407	0.452	0.678
2015	0.830	0.041	0.083	0.166	0.207	0.274	0.332	0.373	0.415	0.622
2016	0.873	0.044	0.087	0.175	0.218	0.288	0.349	0.393	0.437	0.655
2017	0.819	0.041	0.082	0.164	0.205	0.270	0.327	0.368	0.409	0.614
2018	0.910	0.046	0.091	0.182	0.228	0.300	0.364	0.410	0.455	0.683
2019	0.906	0.045	0.091	0.181	0.226	0.299	0.362	0.408	0.453	0.679
2020	0.794	0.040	0.079	0.158	0.199	0.261	0.318	0.357	0.397	0.596

Note: Estimates for the year 1998-2018 were adopted from Neeley (2019)

3.1. Species composition and daily counts in the counting facility

During the 2020 sampling period, a total of 97,447 individuals (raw, unadjusted value) of 18 species and runs were captured in the sampling room (Table 2, Figure 5). Among salmonids, hatchery and wild Spring Chinook (56,669) comprised more than 50% of the total count; and the second highest count was Steelhead (4,946). Total counts varied over time, with 5% of the total pre-March, 2% in March, 47% in April, 46% in May and 0.5% post-May.

Table 2. Total counts by species in the Sample-room sample rates for 2019 and 2020.

Species	2019	2020
Bass	84	87
Bigmouth Minnow	187	131
Bluegill	68	113
Carp	22	176
Catfish	809	757
Chiselmouth	2393	280
Crappie	19	47
Dace	3	0
Lamprey	3654	138
Hatchery Spring Chinook	29532	39047
Perch	17	24
Pumpkinseed	1	0
Shiner	33	11
Sockeye	32	5593
Sucker	1079	590
Whitefish	357	215
Summer/Fall Chinook	13411	26497
Wild Spring Chinook	13507	14925
Coho	8075	1850
Wild Steelhead	5440	4946




Figure 5: Daily catch (raw count) of different species from January through July 2019 (sampling period). Number in green color is the total counts in the sampled during the sampling period.

3.2. Counts of wild and hatchery Spring Chinook

Daily raw counts of the of the hatchery and wild Spring Chinook were divided by the daily sampling rate to arrive at the total number bypassed each sampling day. Missing counts were estimated by linear interpolation for those days in which no sampling was done as mentioned in methodology. After the adjustments, total counts of bypassed hatchery and wild spring Chinook during the sampling period in the sampling facility were estimated to be 148,967 and 53,095, respectively (Table 3). Wild Spring Chinook passed Prosser Dam earlier than their hatchery counterparts, starting with the January initiation of sampling, while hatchery Spring Chinook were not observed until after their volitional release from acclimation sites began in mid-March. The outmigration of both groups was nearly complete by the end of May and ending in late June. The wild counts peaked in April and the hatchery counts peaked in May when fish remaining in acclimation sites were forced out (Table 3, Figure 6).

Table 3 and Figure 6 also contain 2019 counts to illustrate some of the annual variability in Spring Chinook outmigration timing. In 2019 the pre-March and May periods had higher bypass counts of wild fish than April, while hatchery counts were somewhat later in 2019 than in 2020. Annual fluctuations in outmigration timing are a subject for further investigation.

Table 3. Adjusted total count (raw count x sample rate (SR)) of bypassed hatchery and wild Spring Chinook smolts in the Chandler Juvenile Monitoring Facility over 5 temporal strata in 2019 and 2020.

Migration year	Origin	Counts (adjusted)					Total
		Pre-March	March	April	May	Post-May	
2019	Wild	15,489	3,937	10,596	23,290	63	53,374
	Hatchery	0	904	24,775	76,824	198	102,701
2020	Wild	8,843	2,602	30,737	10,851	58	53,092
	Hatchery	8	1,419	64,446	82,305	789	148,967




Figure 6. Percent of outmigration of bypassed wild and hatchery Spring Chinook by temporal stratum for 2019 and 2020




Figure 7. Cumulative catch proportion of bypassed wild and hatchery Spring Chinook by date for 2019 and 2020

3.3. Detection rate of the sampling facility and downstream Dams (MCJ, John Day, BON)

A total of 9519 PIT-tagged hatchery Spring Chinook were detected at the CJMF in 2020. The number of tagged fish detected at Prosser that were also detected at downstream dams depended on downstream detection probability in addition to downstream mortality. Joint detections of hatchery Spring Chinook between Prosser (PRO) and McNary (MCJ), PRO and John Day (JDJ); and PRO and Bonneville Dam (B2J/BCC) in 2020 were found to be 240, 402, and 644, respectively (1286 total). In 2019, joint detections were lower for all dams except at PRO-MCJ: 369, 320, and 465, (1154 total), for PRO-MCJ, PRO-JDJ and PRO-BON, respectively (Table 4).

Overall detection rate at Prosser in 2020 (pooled 43.96 ± 1.0 %) was higher than in 2019 (27.85 ± 0.07 %). The higher detection rate in 2020 was true regardless of which downstream dam was used for a reference to evaluate the Prosser detection rate (Table 4). The most obvious reason for fluctuations in detection rates is fluctuations in diversion rates, but the mean diversion rate for the hatchery spring chinook outmigration period in both years was similar. The timing differences within the outmigration period that were discussed earlier, and their interaction with irrigation demand and river flow could result in an effect on detection rate not evident from a simple mean of diversion rate over the entire outmigration period, or other factors could be responsible. The joint detection rate also varied by stratum or time window (before March, March, April, May and after May) for the tagged hatchery Spring Chinook. Because volitional exit began March 15 in 2020, Prosser detections of hatchery smolts are not expected until late March. Therefore, there were no detections pre-March, and the most detections were in April and May (Table 5). The highest detection rate was in May; whereas the lowest was pre-March the smolts in both years (2019 and 2020) regardless whether it was based on the MCJ reference or all downstream detection sites pooled together (MCJ, JDJ and B2J/BCC, Table 4). This may be related at least in part to normal within-season variations in river flow and diversion quantity.

Table 4. Detection efficiencies of Prosser (PRO) and joint detection of hatchery Spring Chinook smolts between PRO and McNary (MCJ), PRO and John Day (JDJ), PRO and Bonneville (B2J/BCC); and PRO and the pooled detections at MCJ and downstream sites. Detection at Bonneville included the juvenile (smolt) population of hatchery Spring Chinook detected by B2J at BCC antennas, which cover different passage routes. The pooled (ALL) estimate represents the detection probability at Prosser based on detection at one or more downstream dams (MCJ, JDJ and BON).

Migration year	Joint Detection at PRO with	Months					Total	Rate of Detection at Prosser (%)
		Pre-March	March	April	May	Post-May		
2019	McNary (MCJ)	0	6	143	220	0	369	27.09 ±1.2
	John Day (JDJ)	0	2	94	224	0	320	29.46 ±1.4
	Bonneville (BON)	0	4	174	287	0	465	27.04 ±1.8
	Pooled (All)	0	12	411	731	0	1154	27.85 ±0.7
2020	McNary (MCJ)	0	7	117	116	0	240	33.44 ±1.8
	John Day (JDJ)	0	4	146	252	1	402	46.20 ±1.8
	Bonneville (BON)	0	5	295	342	2	644	46.66 ±1.4
	Pooled (All)	0	16	558	710	3	1286	43.96 ±1.0

Table 5. Detection rate of hatchery Spring Chinook smolts at Prosser Dam based on strata (Un-stratified and Stratified) with the reference of McNary Dam and pooled over the three Columbia River dams (MCJ, JDJ and B2J/BCC) for 2019 and 2020. The redistributed detection rate was estimated by pooling the five time periods into two groups: pre-March through April, and May through Post-May.

Migration Year	Reference	Stratified / unstratified	Pre-March	March	April	May	Post-May
2019	Based on McNary Dam	Un-stratified	27.61%	27.61%	27.61%	27.61%	27.61%
		Stratified	0.00%	19.38%	17.72%	39.63%	0.00%
		Stratified (Redistributed)	18.47%	18.47%	18.47%	39.63%	39.63%
	Based on pooled Dams (McJ, JDJ and BON)	Un-stratified	27.93%	27.93%	27.93%	27.93%	27.93%
		Stratified	0.00%	24.64%	20.27%	36.13%	0.00%
		Stratified (Redistributed)	20.07%	20.07%	20.06%	35.88%	35.88%
2020	Based on McNary Dam	Un-stratified	33.44%	33.44%	33.44%	33.44%	33.44%
		Stratified	0.00%	11.85%	22.72%	57.96%	0.00%
		Stratified (Redistributed)	23.68%	23.68%	23.68%	57.96%	57.96%
	Based on pooled Dams (MCJ, JDJ and BON)	Un-stratified	43.96%	43.96%	43.96%	43.96%	43.96%
		Stratified	0.00%	14.87%	33.21%	59.30%	24.56%
		Stratified (Redistributed)	32.26%	32.26%	33.30%	59.23%	59.23%

3.4. Predicted number of outmigrating wild and hatchery Spring Chinook smolts

The total number of hatchery Spring Chinook smolts passing Prosser Dam during the 2019 and 2020 migration periods was almost double that of the wild (natural-origin) run (Table 6).

Furthermore, outmigrating smolts of hatchery origin were more numerous in 2020 than in the 2019 outmigration period by all of the four estimation methods employed. Applying the detection rates derived from hatchery Spring Chinook to their wild counterparts (Table 5), the estimates of wild Spring Chinook smolts passing Prosser Dam also varied between years.

Depending on the estimation method, wild outmigration ranged from 154,530 to 175,427 in 2019; and from 115,300 to 201,313 in 2020. The hatchery smolt estimates ranged from 310,836 to 353,803 in 2019 and from 371,069 to 500,195 in 2020.

The details of the juvenile Spring Chinook passage estimates at Prosser Dam based on different estimators from 1999-2020 are given in Appendix A of this report. The estimates based on the method with temporal strata Pre-May, May, June, Post-June was found to be slightly higher than the estimates based on non-stratified detection rates.

Table 6. The estimated number of wild and hatchery Spring Chinook smolts migrating past Prosser Dam in 2019 and 2020 using four estimation methods.

Migration Year	Origin	Estimates of outmigration population based on different methods			
		McN_UnStr (Method 1)	McN_Str (Method 2)	Pooled_UnStr (Method 3)	Pooled_Str (Method 4)
2019	Natural Origin	168,119	154,848	175,427	154,530
	Hatchery	310,836	353,803	319,579	343,212
2020	Natural Origin	201,313	168,133	151,265	115,300
	Hatchery	456,852	500,195	371,069	380,494

Choosing the best estimate was challenging. We compared these estimates with another independent estimate derived from survival rate (Table 7). In migration year 2020, the average survival rate from the three acclimation sites to Prosser Dam was $71.22 \pm 3.91\%$ (based on the CJS model) and the total number of released hatchery Spring Chinook smolts during 2020 was 624,200. Multiplying the survival rate by the released population, the total outmigration of hatchery Spring Chinook from Prosser was estimated to be $444,555 \pm 47,958$ (mean \pm 95 % confidence Intervals). This estimate was closest to the estimate derived from the pooled stratified (redistributed) method (Tables 6 and 7).

Table 7. Number of Spring Chinook (hatchery) smolts release at Acclimation sites and its survival rate from the acclimation sites to Below Prosser based on CJS model and the estimated outmigration smolts from Prosser Dam for the migration year 2019 and 2020.

Migration Year	No. of smolts at Acclimation sites	Survival rate from the acclimation site to below Prosser		Estimated outmigration smolt from Prosser	
		Average	SE	Average	95% CI
2019	673,218	50.82	2.2	342,129	29,103
2020	624,200	61.22	3.91	382,135	47,958

However, the estimates based on survival rate may still have some bias because the survival rate may not be homogeneous among the sampling months, especially due to variation in river flow at Prosser within the sampling period. However, previous years’ analyses also showed that the estimate based on pooled-lower-dam-based stratified detection rate (method 4) was highly correlated with hatchery returns.

3.5. Annual trend of juvenile Prosser-passage estimates (hatchery and wild) by stock

Annual juvenile Prosser-passage estimates from outmigration years 1999 through 2020 are given in Table 8 by stock of wild/Natural origin (Naches, American, and Upper Yakima rivers) plus hatchery Upper Yakima River origin. It showed that Prosser juvenile estimates for both wild (natural) and hatchery vary among the outmigration year. Total Spring Chinook outmigration per year was $551,589 \pm 29,107$. However, in an average year, $226,910 \pm 25,682$ wild and $324,679 \pm 15,931$ hatchery Spring Chinook smolt passed downstream at Prosser (Table 8 and Figure 8). Wild (natural) Spring Chinook from the American River had the lowest average, Naches had the second, and the supplemented Upper Yakima stock had the highest average among the wild stocks (Figure 8). Hatchery juvenile Spring Chinook passage at Prosser Dam in 2020 was the highest since 2014 (Table 8).

Table 8. Annual estimated wild and hatchery-origin smolt passage at Prosser Dam from 1999 through 2020. Estimates for the outmigration years from 1998 through 2018 were adopted from Neeley (2019).

Brood Year (BY)	Outmigration Year	Wild Stock Estimates				Hatchery (Upper Yakima)	Total Wild & Hatchery
		Total Wild	Naches	American	Upper Yakima		
1997	1999	584,016	93,427	63,000	427,588	187,669	771,685
1998	2000	199,416	55,737	50,944	92,795	303,688	503,104
1999	2001	148,460	Genetic samples not taken			281,256	429,716
2000	2002	467,359	92,323	17,835	357,201	366,950	834,309
2001	2003	308,959	74,498	42,867	191,594	154,329	463,288
2002	2004	169,397	59,978	35,800	73,619	290,950	460,347
2003	2005	134,859	45,321	35,564	5,374	236,443	371,302
2004	2006	133,238	49,947	7,882	75,409	300,508	433,746
2005	2007	99,341	26,684	11,103	61,554	351,359	450,700
2006	2008	120,013	32,589	6,811	80,613	265,485	385,498
2007	2009	237,228	80,756	26,498	128,974	415,923	653,151
2008	2010	220,950	77,397	30,354	113,198	382,878	603,828
2009	2011	304,322	58,904	17,882	227,536	442,564	746,886
2010	2012	258,106	81,483	23,609	153,014	391,446	649,552
2011	2013	365,386	85,577	25,681	254,228	372,079	737,465
2012	2014	263,266	79,450	28,622	155,194	408,222	671,488
2013	2015	125,150	29,885	13,769	81,496	332,715	457,865
2014	2016	185,442	57,657	15,378	112,407	403,938	589,380
2015	2017	208,929	62,190	24,455	122,285	273,248	482,177
2016	2018	131,489	37,500	9,824	76,150	290,644	422,133
2017	2019	175,427	41,690	22,379	127,176	319,579	495,006
2018	2020	151,265	34,770	5,007	115,288	371,069	522,333
Average/year		226,910	59,894	24,536	146,728	324,679	551,589
Standard Error (SE)		25,682	5,348	3,266	21,313	15,931	29,107




Figure 8. Average annual Prosser-passage estimates for wild and hatchery Spring Chinook by stock, outmigration years 2000-2020. The dot within each box is the mean and the horizontal line is the median. Outmigration year 1999 was not included for this box plot because in that year only a few raceways were used for hatchery production, and 1999 was also the last outmigration year in which “ISO” PIT tags with poor read range were used.

Because the smolt passage estimates for the three largest stock groupings (Total wild, Upper Yakima wild, and Upper Yakima hatchery) varied by outmigration year, we further estimated whether the outmigration smolt decreased over years (temporal trends) and whether there were differences among stocks. In 1999, only 14 of 18 raceways were used for hatchery production. As a result, the Prosser passage estimates for hatchery smolts in 1999 were low, which might not compare well with other years’ hatchery estimates. Two relationships were developed using the data with and without 1999’s passage estimates for all three groups (total wild, Upper Yakima wild, and Upper Yakima hatchery). In both datasets, the total number of outmigrating wild smolts and the number of wild upper Yakima smolts seemed to be decreasing over time, whereas the population of hatchery in Upper Yakima sub-basin seemed to be increasing; but neither trend was statistically significant (Figure 9).




Figure 9. Estimated passage at Prosser Dam for total wild, Upper Yakima wild, and Upper Yakima hatchery Spring Chinook smolts with predicted trends by outmigration year. A1, B1 and C1 are for outmigration years 2000-2020 (omitting 1999 data); whereas the A2, B2 and C2 relationships were developed using all outmigration years from 1999 through 2020.

Although outmigration of hatchery smolts is increasing but not significantly, there is a possibility that a true positive increase in hatchery smolts coming from the Upper Yakima would have an associated true negative decrease in Upper Yakima wild passage. Therefore, the assessment of wild and hatchery trends was further examined using the percentage changes in outmigrating smolt populations of wild and hatchery over years (hatchery plus wild).

Table 9. Percentage of wild and hatchery spring Chinook stocks in juvenile Prosser passage estimates, comparing the hatchery stock to all wild stocks and to the Upper Yakima wild stock by itself.

Brood Year (BY)	Out-migration Year	Total Yakima Basin		Only Upper Yakima River	
		% Hatchery of Total	% Wild of Total	% Hatchery of Upper Yakima Stock	% Wild of Upper Yakima stock
1997	1999	24.32%	75.68%	30.50%	69.50%
1998	2000	60.36%	39.64%	76.60%	23.40%
1999	2001	65.45%	34.55%	Genetic samples not taken	
2000	2002	43.98%	56.02%	50.67%	49.33%
2001	2003	33.31%	66.69%	44.61%	55.39%
2002	2004	63.20%	36.80%	79.81%	20.19%
2003	2005	63.68%	36.32%	97.78%	2.22%
2004	2006	69.28%	30.72%	79.94%	20.06%
2005	2007	77.96%	22.04%	85.09%	14.91%
2006	2008	68.87%	31.13%	76.71%	23.29%
2007	2009	63.68%	36.32%	76.33%	23.67%
2008	2010	63.41%	36.59%	77.18%	22.82%
2009	2011	59.25%	40.75%	66.04%	33.96%
2010	2012	60.26%	39.74%	71.90%	28.10%
2011	2013	50.45%	49.55%	59.41%	40.59%
2012	2014	60.79%	39.21%	72.45%	27.55%
2013	2015	72.67%	27.33%	80.33%	19.67%
2014	2016	68.54%	31.46%	78.23%	21.77%
2015	2017	56.67%	43.33%	69.08%	30.92%
2016	2018	68.85%	31.15%	79.24%	20.76%
2017	2019	64.56%	35.44%	71.53%	28.47%
2018	2020	71.04%	28.96%	76.30%	23.70%

Note: Estimates for the outmigration years from 1998 through 2018 were adopted from Neeley (2019)

We found that while the rate of change in the outmigrating hatchery smolt population over years seemed to be positive (Figure 10) and the trend for wild stocks were negative, the relationship of hatchery passage to wild passage (all wild stocks or only the Upper Yakima wild stock) was not statistically significant. This indicates that the production and releases of spring Chinook hatchery smolts into the upper Yakima do not have an effect on the production of wild smolts. The reduction of the production of wild smolts could be influenced by many factors including habitat loss that limits the carrying capacity and it eventually reduces the survival rate and the total outmigration.




Figure 10. Linear trend on the percentage of hatchery and wild components of the total outmigrating spring Chinook populations for all Yakima Basin stocks and only the Upper Yakima River stock

3.6. Genetic variation among stocks (Upper Yakima, Naches, American)

As discussed above, wild Yakima Basin Spring Chinook are comprised of multiple stocks, of which Upper Yakima River, Naches River, and American River stocks have been identified by demographic characteristics and supported by genetic analysis. Reproductively isolated

populations usually differ in productivity. We, therefore, further evaluated whether the rate of outmigration of these genetic stocks has changed over time. Because no hatchery program has been implemented in the American and Naches rivers, we hypothesized that the rate of decline should be higher in the Upper Yakima's wild Spring Chinook, if the hatchery program affected wild productivity.

The annual outmigration estimates showed that the wild Spring Chinook smolt population declined over the 2000-2020 outmigration years (Figure 9) for all three stocks. The rate of decline of the smolt in the Wild Upper Yakima stock was -1529 smolts/year, but the trend was not significant ($R^2=0.02$, $p=0.469$), nor was the rate of decline for the Naches River stock (Slope=881/year, $R^2=0.67$, $p=0.273$). Only the American stock declined significantly (Slope= -1072/year, $R^2=0.228$, $p=0.04$, Figure 9); there has been no introduction of hatchery smolts into the American River.

In fact, the American River seems to have suffered a relatively low anthropogenic effect compared to the other rivers. It is also the coldest and has entirely natural flow that persists through the summer. If hatchery or other local anthropogenic factors had a negative influence, the American River stock should have declined the least, but the opposite was true in terms of outmigrant abundance.




Figure 11. The relationship between estimated smolt passage of Wild Spring Chinook of Naches, American, and Upper Yakima stock by outmigration year.

3.7. Contribution of each stock to outmigration

For outmigration years 1999-2020, about 61% of the total wild outmigration was contributed by the Upper Yakima wild stock; while 28% and 11% were contributed by Naches and American River stocks, respectively (Table 10).

Table 10. American, Naches and Upper Yakima Percentages of Prosser passage of wild Spring Chinook smolts at Prosser Dam. Data for outmigration years 1998 through 2017 were adopted from Neeley (2018).

Brood Year	Outmigration Year	Naches	American	Upper Yakima
1997	1999	16.00%	10.79%	73.22%
1998	2000	27.95%	25.55%	46.53%
1999	2001			
2000	2002	19.75%	3.82%	76.43%
2001	2003	24.11%	13.87%	62.01%

2002	2004	35.41%	21.13%	43.46%
2003	2005	33.61%	26.37%	40.02%
2004	2006	37.49%	5.92%	56.60%
2005	2007	26.86%	11.18%	61.96%
2006	2008	27.15%	5.68%	67.17%
2007	2009	34.04%	11.17%	54.37%
2008	2010	35.03%	13.74%	51.23%
2009	2011	19.36%	5.88%	74.77%
2010	2012	31.57%	9.15%	59.28%
2011	2013	23.42%	7.03%	69.58%
2012	2014	30.18%	10.87%	58.95%
2013	2015	23.88%	11.00%	65.12%
2014	2016	31.09%	8.29%	60.62%
2015	2017	29.77%	11.70%	58.53%
2016	2018	28.52%	7.47%	57.91%
2017	2018	23.76%	12.76%	72.50%
2018	2020	22.99%	3.31%	76.22%
Mean		27.71%	11.27%	61.26%
SE		1.25%	1.32%	2.20%

Table 11. Estimated Wild Spring Chinook stock distributions (American, Naches and Upper Yakima River) within the genetic sampling periods (Pre-March through Post-May). The data were provided by WDFW.

migrat ion year	American					Naches					U. Yakima				
	Pre- March	March	April	May	Post- May	Pre- March	March	April	May	Post- May	Pre- March	March	April	May	Post-May
1999	8.08%	8.08%	8.08%	12.00%	28.00%	6.06%	6.06%	6.06%	29.00%	33.00%	85.86%	85.86%	85.86%	59.00%	39.00%
2000	16.18%	16.18%	22.14%	46.94%	46.94%	22.06%	22.06%	30.99%	36.73%	36.73%	61.76%	61.76%	46.88%	16.33%	16.33%
2002	3.81%	3.81%	3.81%	3.86%	3.86%	19.68%	19.68%	19.68%	20.29%	20.29%	76.51%	76.51%	76.51%	75.85%	75.85%
2003	13.43%	13.43%	13.43%	16.03%	16.03%	21.64%	21.64%	21.64%	34.24%	34.24%	64.93%	64.93%	64.93%	49.73%	49.73%
2004	6.46%	4.27%	21.50%	34.72%	31.25%	33.84%	29.27%	36.47%	34.03%	18.75%	59.70%	66.46%	42.03%	31.25%	50.00%
2005	21.39%	18.87%	29.57%	32.14%	0.00%	35.32%	7.55%	35.36%	23.21%	17.86%	43.28%	73.58%	35.07%	44.64%	82.14%
2006	7.36%	0.00%	5.52%	5.45%	2.27%	39.88%	25.96%	35.95%	39.11%	15.91%	52.76%	74.04%	58.53%	55.45%	81.82%
2007	9.10%	14.50%	6.81%	16.75%	11.54%	18.20%	32.30%	24.72%	29.78%	26.07%	72.70%	53.20%	68.47%	53.47%	62.39%
2008	8.33%	0.00%	5.22%	5.00%	14.81%	8.33%	14.29%	25.22%	31.11%	51.85%	83.33%	85.71%	69.57%	63.89%	33.33%
2009	9.80%	10.93%	12.06%	10.95%	36.29%	35.60%	32.43%	29.25%	40.78%	28.23%	54.60%	56.64%	58.69%	48.27%	35.48%
2010	30.31%	0.00%	14.16%	11.88%	0.00%	7.35%	19.50%	37.13%	33.63%	75.49%	62.34%	80.50%	48.71%	54.49%	24.51%
2011	8.64%	0.00%	3.49%	5.92%	16.65%	18.19%	19.75%	23.96%	13.10%	0.00%	73.17%	80.25%	72.55%	80.98%	83.35%
2012	10.99%	5.31%	6.17%	13.65%	23.46%	31.62%	29.60%	29.32%	38.48%	29.45%	57.39%	65.09%	64.51%	47.87%	47.09%
2013	8.23%	2.30%	5.72%	16.96%	6.39%	17.43%	20.59%	27.50%	29.53%	7.85%	74.34%	77.11%	66.78%	53.51%	85.76%
2014	11.65%	12.03%	9.09%	11.95%	13.86%	41.19%	21.74%	30.16%	38.12%	0.00%	47.16%	66.23%	60.74%	49.93%	86.14%
2015	13.86%	11.62%	8.92%	14.74%	14.74%	16.80%	26.32%	23.13%	24.09%	24.09%	69.34%	62.06%	67.96%	61.17%	61.17%
2016	5.69%	7.42%	9.44%	13.00%	3.71%	26.41%	23.18%	38.42%	34.52%	0.00%	67.90%	69.40%	52.13%	52.49%	96.29%
2017	10.20%	11.21%	15.80%	10.78%	37.16%	31.70%	27.73%	27.10%	29.57%	11.47%	58.10%	61.06%	57.10%	59.65%	51.37%
2018	8.80%	3.30%	5.82%	10.40%	25.00%	23.20%	33.00%	35.11%	41.94%	25.00%	68.00%	63.70%	59.08%	47.66%	50.00%
2019	9.90%	12.44%	14.70%	14.71%	0.00%	17.82%	21.89%	23.32%	35.29%	0.00%	72.28%	65.67%	61.98%	50.00%	100.0%

2020	3.78%	6.50%	2.84%	3.60%	0.00%	3.78%	6.50%	2.84%	3.60%	0.00%	76.22%	73.17%	74.47%	66.19%	100.0%
------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	--------	--------	--------	--------	--------

3.8. Relationship between Wild Juvenile passage estimates and estimated Adult Returns

Since the number of smolts outmigrating from Prosser (Prosser-passage estimates) varied among years, we further evaluated whether this variation corresponded to adult returns. Or in other words, does the fluctuation of annual wild juvenile passage at Prosser synchronize with the fluctuation of the adult returns at Prosser? To answer the question, we built a univariate relationship between the total Juvenile Prosser estimates of wild Spring Chinook and the predicted adult return to Prosser. Table 12 presents the brood year Prosser escapement (the escapement measures are taken as a surrogate of spawner number) of the parental generation in addition to total juvenile Prosser passage and Prosser return. The relationship between juvenile-to-adult correlation of total wild juvenile passage to adult return from each outmigration was significantly high, with an R^2 of 79% and p value < 0.01 (Figure 12), indicating that estimated number of outmigration smolts are reasonably accurate.

Table 12. Total estimated escapement (Estimated Spawners (wild/natural) at Yakima river mouth), juvenile passage and return to Prosser of each wild Spring Chinook brood for brood years 1997-2018. Estimated value for the Prosser escapement and Prosser return were adopted from Table 10 and Table 3 of Bosch (2020), respectively. The shaded yellow color indicates that adult returns from these brood years are incomplete.

Brood Year	Out-migration Year	Estimated Spawners (wild/natural) at Yakima river mouth	Total Juvenile Prosser Passage	Prosser return
1997	1999	2,337	584,016	12,808
1998	2000	1,307	199,476	7,283
1999	2001	1,439	148,460	4,090
2000	2002	15,976	467,359	11,128
2001	2003	17,916	308,959	7,731
2002	2004	11,113	169,397	3,850
2003	2005	5,933	134,859	2,195
2004	2006	12,893	133,218	3,687
2005	2007	7,617	99,265	4,089
2006	2008	5,050	123,735	5,118

2007	2009	3,308	250,846	7,610
2008	2010	5,922	221,228	6,739
2009	2011	8,172	303,711	4,167
2010	2012	9,875	252,029	6,148
2011	2013	11,644	365,468	7,002
2012	2014	7,383	267,433	3,941
2013	2015	6,352	123,289	3,736
2014	2016	7,882	53,478	1,928
2015	2017	7,569	57,051	870
2016	2018	5,613	131,489	1876
2017	2019	5,015	175,427	79
2018	2020	2,451	151,265	




Figure 12. The relationship between total smolts outmigration and Prosser returns of progeny (adult returns) of wild Spring Chinook. Since the Spring Chinook can spend as many as 4 years in the ocean, the relationship was made for the populations that outmigrated from 1997 through 2018. Each point is the outmigration year from 1999 to 2018.




Figure 13. Year-to-year trends in juvenile outmigration from Prosser and Adult returns of Spring Chinook (wild) for outmigration years 1999-2018.

3.9. Relationship between estimated juvenile passage and river flow

3.9.1. Annual

The annual juvenile passage estimate of wild and hatchery Spring Chinook at Prosser Dam tends to increase with average river flow for the outmigration period, but this relationship was significant only for Upper Yakima hatchery smolts (Figure 14).




Figure 14. The relationship between the annual passage estimate for Total Wild, Upper Yakima Wild and Upper Yakima hatchery stocks of juvenile Yakima Spring Chinook and the March-June average river flow approaching Prosser Dam. The flow is the sum of the average flow measured at the gaging stations CHCW and YRPW.

3.9.2. Daily

The annual juvenile passage estimate of wild and hatchery Spring Chinook at Prosser Dam tends to increase with average river flow for the outmigration period, but this relationship was significant only for Upper Yakima hatchery smolts (Figure 14).

We further evaluated whether the daily estimated number of wild out-migrating smolts in 2020 was affected by daily river flow (the river flow approaching the Prosser Dam, which is the sum of the flow measured at the gaging stations CHCW and YRPW). Figure 15 shows day-to-day trends of the estimated daily counts and the daily river flow and water temperature, and shows that daily estimated outmigrating smolts increased when flow pulses occurred, whether due to natural runoff or reservoir releases made to facilitate outmigration.




Figure 15. Daily estimated counts of hatchery and wild/natural Spring Chinook smolts, water temperature and river flow approaching Prosser Dam for the period in which the CJMF was operated during 2020. The river flow (blue line) is the flow approaching the dam (sum of the flow measured at the gaging stations CHCW and YRPW). Red line is the daily average temperature measured at Prosser.

We analyzed scatter plots and linear relationships between the daily estimated smolt counts of both runs (natural origin and hatchery origin) and daily river flow; and it showed that in general, daily estimated out-migrating smolts was high if the river flow was high (see Figure 16).

Furthermore, the smolts were found to be out-migrated more when flow pulse events occurred whether due to natural runoff or reservoir releases made to facilitate outmigration, indicating that when smolt migration appears to be stalled, releases of pulses in flow from reservoirs can improve the rate of smolt out-migration (Figure 15).




Figure 16. The relationship between daily estimated wild/natural and hatchery Spring Chinook smolt passage and river flow approaching Prosser Dam for the period in which the Chandler Canal monitoring facility was operated during 2020.

4. References

- Bosch, B. 2020. Run Size Forecast for Yakima River Adult Spring Chinook, 2021 (preliminary). Yakima Klickitat Fisheries Project, Yakama Nation Fisheries Resource Management, 760 Pence Road, Yakima, Washington, 98908
- Fiander, W., D. E. Fast, and W. J. Bosch (editors). 2019. Yakima-Klickitat Fisheries Project Monitoring and Evaluation- Yakima Subbasin, Final Report for the performance period May/2018-April/2019. Project Number 1995-063-25, 275 electronic pages.
- Neeley, D. 2019. Methods of Estimating Smolt Survival and Passage. Appendix C *In* Final report for the performance period May 1, 2018 through April 30, 2019, edited by W. Fiander, D. E. Fast, and W. J. Bosch. Yakima/Klickitat Fisheries Project.
- Neeley, D. 2012. 2012 Annual Report: Chandler Certification for Yearling Outmigrating Spring Chinook Smolt.
- Pyper, B. J. and Smith, D. L. 2005. Evaluation Of Salmonid Survival Resulting From Flow Alterations To The Lower Yakima River, S.P. Cramer and Associates, submitted to Kennewick Irrigation District and US Bureau of Reclamation, Yakima
- Seamons, T. R. and Bowman, C. M. 2019 and 2020. DNA-Based Population-of-Origin Assignments of Chinook Smolts Outmigrating Past Chandler Trap at Prosser Dam (Yakima River) in 2018. Washington Department of Fish and Wildlife, Molecular Genetics Laboratory, 600 Capital Way N., Olympia, WA.
- Zabel, R. W. & Achord, S. (2004) Relating size of juveniles to survival within and among populations of Chinook salmon. *Ecology*, 85, 795–806.
- Zabel, R. W., T. Wagner, J. L. Congleton, S. G. Smith, and J. G. Williams. 2005. Survival and selection of migrating salmon from capture-recapture models with individual traits. *Ecological Applications* 15:1427–1439.

5. Supplementary information: Detailed Passage-Estimates

Detailed Passage-Estimates for each year from 1998 through 2020

Supplementary information: Detailed Passage-Estimates for each year from 1998 through 2020

5.1. Year 1998

1998	Brood-Year 1996	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer				
Wild	Prosser Wild Tally	0	10618	106253	6174	292	123337	123337				
	American	WDFW Percent	0	0.00	0.02	0.02	0.12					
		Estimated Prosser Tally	0	0.00	2125.06	124.72	35.06	2284.84	2284.84			
		WDFW Percent	0.21	0.21	0.24	0.24	0.51					
	Naches	Estimated Prosser Tally	0	2230	25501	1497	149	29376	29376			
		WDFW Percent	0.79	0.79	0.74	0.74	0.37					
	Upper Yakima	Estimated Prosser Tally	0	8388	78627	4552	108	91676	91676			
		Yakima Passage Wild Tally	0	10618	106253	6174	292	123337	Expanded Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
	Estimate a.	Detection Efficiency										
		Total Passage										
		American Passage										
		Naches Passage										
		American & Naches Passage										
		Upper Yakima Passage										
	Estimate b.	Detection Efficiency										
		Total Passage										
		American Passage										
		Naches Passage										
		American & Naches Passage										
		Upper Yakima Passage										
	Estimate c.	Detection Efficiency										
		Total Passage										
		American Passage										
		Naches Passage										
		American & Naches Passage										
		Upper Yakima Passage										
	Estimate e.	Detection Efficiency										
		Total Passage										

*

American Passage
 Naches Passage
 American & Naches Passage
 Upper Yakima Passage

			Expanded Elastomer	Expanded PIT	PIT- Tag/Total	Calibration Index
Hatchery		Prosser Hatchery Tally				
McN-Str Hatch	Estimate a.	Total Passage				
McN-UnStr Hatch	Estimate b.	Total Passage				
Pooled Str Hatch	Estimate c.	Total Passage				
Pooled UnStr Hatch	Estimate e.	Total Passage				

5.2. Year 1999

1999	Brood-Year 1997	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer				
Wild	Prosser Wild Tally	41232.89541	407	29431	51920	1577	124569	124569				
	American	WDFW Percent	0.08	0.08	0.08	0.12	0.28					
		Estimated Prosser Tally	3332	33	2378	6230	442	12415	12415			
		WDFW Percent	0.06	0.06	0.06	0.29	0.33					
	Naches	Estimated Prosser Tally	2499	25	1784	15057	520	19885	19885			
		WDFW Percent	0.86	0.86	0.86	0.59	0.39					
	Upper Yakima	Estimated Prosser Tally	35401.98091	350	25269	30633	615	92269	92269			
		Yakima Passage Wild Tally	41233	407	29431	51920	1577	124569	Expanded Elastomer	Calibrated Total	PIT- Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	18.5%	18.5%	18.5%	25.5%	5.0%					
		Total Passage	222873	2201	159082	203681	31262	619099	619099	571397		0.9229
		American Passage	18010	178	12855	24442	8753	64238	64238	59288		
		Naches Passage	13507	133	9641	59067	10316	92666	92666	85526		
		American & Naches Passage	31517	311	22496	83509	19070	156904	156904	144815		
		Upper Yakima Passage	191355	1890	136586	120172	12192	462195	462195	426583		
McN UnStr Wild	Estimate b.	Detection Efficiency	23.0%	23.0%	23.0%	23.0%	23.0%					
		Total Passage	179338	1771	128008	225822	6860	541799	541799	502917		0.9282
		American Passage	14492	143	10344	27099	1921	53998	53998	50123		

		Naches Passage	10869	107	7758	65488	2264	86486	86486	80280		
		American & Naches Passage	25361	251	18102	92587	4184	140485	140485	130403		
		Upper Yakima Passage	153977	1521	109906	133235	2675	401314	401314	372514		
Pooled Str Wild	Estimate c.	Detection Efficiency	19.4%	19.4%	19.4%	23.0%	3.8%					
		Total Passage	212650	2101	151786	225518	41751	633805	633805	584016	0.9214	
		American Passage	17184	170	12266	27062	11690	68371	68371	63000		
		Naches Passage	12888	127	9199	65400	13778	101392	101392	93427		
		American & Naches Passage	30072	297	21465	92462	25468	169764	169764	156428		
		Upper Yakima Passage	182579	1803	130321	133056	16283	464042	464042	427588		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	20.3%	20.3%	20.3%	20.3%	20.3%					
		Total Passage	203022	2005	144913	255644	7766	613350	613350	569333	0.9282	
		American Passage	16406	162	11710	30677	2174	61130	61130	56743		
		Naches Passage	12304	122	8783	74137	2563	97908	97908	90882		
		American & Naches Passage	28710	284	20493	104814	4737	159038	159038	147624		
		Upper Yakima Passage	174312	1722	124420	150830	3029	454312	454312	421709		
Hatchery		Prosser Hatchery Tally	0	7	1812	31529	1371	34719	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estim a.	Total Passage	0	39	9796	123685	27175	160696	179215	165406	0.1033	0.9229
McN-UnStr Hatch	Estimate b.	Total Passage	0	32	7883	137130	5963	151007	168410	156324		0.9282
Pooled Str Hatch	Estimate c.	Total Passage	0	38	9347	136946	36292	182622	203668	187669		0.9214
Pooled UnStr Hatch	Estimate e.	Total Passage	0	36	8924	155240	6750	170950	190650	176968		0.9282

5.3. Year 2000

2000	Brood-Year 1998	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer
		12636.7108						
Wild	Prosser Wild Tally	9	252	11172	19815	814	44690	44690
	American	WDFW Percent	0.16	0.16	0.22	0.47	0.47	
		Estimated Prosser Tally	2044	41	2473	9301	382	14241
		WDFW Percent	0.22	0.22	0.31	0.37	0.37	
	Naches	Estimated Prosser Tally	2788	56	3462	7279	299	13883
		WDFW Percent	0.62	0.62	0.47	0.16	0.16	
	Upper Yakima	Estimated Prosser Tally	7805	156	5237	3235	133	16566

		Yakima Passage Wild Tally	12637	252	11172	19815	814	44690	Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	12.5%	12.5%	31.6%	52.6%	31.0%					
		Total Passage	100754	2008	35311	37686	2627	178387	178387	222645		1.2481
		American Passage	16298	325	7816	17689	1233	43362	43362	54120		
		Naches Passage	22225	443	10943	13844	965	48420	48420	60433		
		American & Naches Passage	38524	768	18759	31533	2199	91782	91782	114553		
		Upper Yakima Passage	62231	1240	16552	6153	429	86605	86605	108091		
McN UnStr Wild	Estimate b.	Detection Efficiency	41.7%	41.7%	41.7%	41.7%	41.7%					
		Total Passage	30333	605	26818	47564	1955	107274	107274	132166		1.2320
		American Passage	4907	98	5936	22326	918	34184	34184	42116		
		Naches Passage	6691	133	8311	17472	718	33326	33326	41059		
		American & Naches Passage	11598	231	14247	39798	1636	67510	67510	83175		
		Upper Yakima Passage	18735	373	12571	7765	319	39764	39764	48991		
Pooled Str Wild	Estimate c.	Detection Efficiency	15.9%	15.9%	30.0%	51.1%	30.0%					
		Total Passage	79697	1589	37229	38770	2713	159998	159998	199476		1.2467
		American Passage	12892	257	8241	18198	1273	40862	40862	50944		
		Naches Passage	17580	350	11537	14242	997	44707	44707	55737		
		American & Naches Passage	30472	607	19778	32440	2270	85568	85568	106681		
		Upper Yakima Passage	49224	981	17451	6330	443	74430	74430	92795		
Pooled UnStr Wild	Estimate e.	Total Passage	41.2%	41.2%	41.2%	41.2%	41.2%					
		Total Passage	30699	612	27141	48137	1979	108568	108568	133760		1.2320
		American Passage	4966	99	6008	22595	929	34596	34596	42624		
		Naches Passage	6772	135	8411	17683	727	33728	33728	41554		
		American & Naches Passage	11738	234	14419	40278	1656	68324	68324	84178		
		Upper Yakima Passage	18961	378	12722	7859	323	40244	40244	49582		
Hatchery	Estimate a.	Prosser Hatchery Tally	0	11	12187	59659	21234	93091	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
		Total Passage	0	91	38517	11346	6	68501	220575	235507	293937	0.0634

McN-UnStr Hatch	Estimate b.	Total Passage	0	27	29253	14320 6	50971	223458	238585	293946	1.2320
Pooled Str Hatch	Estimate c.	Total Passage	0	72	40610	11673 1	70728	228141	243585	303688	1.2467
Pooled UnStr Hatch	Estimate e.	Total Passage	0	28	29606	14493 3	51586	226152	241461	297490	1.2320

5.4. Year 2001

2001	Brood-Year 1999	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer			
Wild	Prosser Wild Tally	4678.6417 82	3236	101993	27763	1307	138977	138977			
	American	WDFW Percent									
		Estimated Prosser Tally						0	Genetic Sample Analysis not Performed		
		WDFW Percent									
	Naches	Estimated Prosser Tally	genetic assignment to Upper Yakima Stock not possible						0		
		WDFW Percent									
	Upper Yakima	Estimated Prosser Tally						0			
		Yakima Passage Wild Tally					138977	Elastomer			
McN Str Wild	Estimate a.	Detection Efficiency	76.1%	76.1%	76.1%	86.8%	91.9%				
		Total Passage	6150	4253	134076	31992	1421	177893	177893	149124	0.8383
		American Passage									
		Naches Passage									
		American & Naches Passage									
		Upper Yakima Passage									
McN UnStr Wild	Estimate b.	Detection Efficiency	83.9%	83.9%	83.9%	83.9%	83.9%				
		Total Passage	5577	3857	121571	33092	1558	165654	165654	143613	0.8669
		American Passage									
		Naches Passage									
		American & Naches Passage									
		Upper Yakima Passage									

Pooled Str Wild	Estimate c.	Detection Efficiency	77.3%	77.3%	77.3%	85.9%	90.9%					
		Total Passage	6052	4185	131931	32310	1438	175917	175917	148460	0.8439	
		American Passage										
		Naches Passage										
		American & Naches Passage										
Pooled UnStr Wild	Estimate e.	Detection Efficiency	83.7%	83.7%	83.7%	83.7%	83.7%					
		Total Passage	5589	3865	121828	33162	1561	166004	166004	143917	0.8669	
		American Passage										
		Naches Passage										
		American & Naches Passage										
		Upper Yakima Passage										
Hatchery		Prosser Hatchery Tally	0	4	96207	148783	16931	261925	Expanded Elastomer	Expand ed PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	0	5	126468	171448	18415	316337	333380	279467	0.0511	0.8383
McN-UnStr Hatch	Estimate b.	Total Passage	0	5	114674	177343	20181	312202	329022	285245		0.8669
Pooled Str Hatch	Estimate c.	Total Passage	0	5	124446	173151	18633	316235	333273	281256		0.8439
Pooled UnStr Hatch	Estimate e.	Total Passage	0	5	114916	177717	20223	312862	329717	285847		0.8669

5.5. Year 2002

2002	Brood-Year 2000	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer			
Wild	Prosser Wild Tally	66506.36024	26080	101052	40512	62	234213	234213			
American	WDFW Percent	0.04	0.04	0.04	0.04	0.04					
	Estimated Prosser Tally	2534	994	3850	1566	2	8945	8945			
Naches	WDFW Percent	0.20	0.20	0.20	0.20	0.20					
	Estimated Prosser Tally	13090	5133	19890	8220	13	46345	46345			
Upper Yakima	WDFW Percent	0.77	0.77	0.77	0.76	0.76					
	Estimated Prosser Tally	50882.64387	19954	77313	30726	47	178922	178922			
	Yakima Passage Wild Tally	66506	26080	101052	40512	62	234213	Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	31.7%	31.7%	56.3%	65.9%	25.2%				
		Total Passage	209858	82295	179367	61477	247	533244	533244	466904	0.8756

		American Passage	7995	3135	6833	2376	10	20348	20348	17817		
		Naches Passage	41305	16198	35304	12474	50	105331	105331	92227		
		American & Naches Passage	49300	19333	42137	14850	60	125679	125679	110044		
		Upper Yakima Passage	160558	62963	137230	46628	187	407565	407565	356861		
McN UnStr Wild	Estimate b.	Detection Efficiency	59.5%	59.5%	59.5%	59.5%	59.5%					
		Total Passage	111740	43819	169781	68066	104	393510	393510	349322	0.8877	
		American Passage	4257	1669	6468	2631	4	15028	15028	13341		
		Naches Passage	21993	8625	33417	13810	21	77867	77867	69123		
		American & Naches Passage	26250	10294	39885	16441	25	92895	92895	82464		
		Upper Yakima Passage	85490	33525	129896	51625	79	300615	300615	266858		
Pooled Str Wild	Estimate c.	Detection Efficiency	32.8%	32.8%	53.9%	65.2%	7.9%					
		Total Passage	202911	79571	187367	62093	784	532726	532726	467359	0.8773	
		American Passage	7730	3031	7138	2400	30	20329	20329	17835		
		Naches Passage	39938	15662	36879	12599	159	105236	105236	92323		
		American & Naches Passage	47668	18693	44016	14998	189	125565	125565	110158		
		Upper Yakima Passage	155243	60878	143350	47095	595	407161	407161	357201		
Pooled UnStr Wild	Estimate e.	Total Passage	57.6%	57.6%	57.6%	57.6%	57.6%					
		Total Passage	115447	45272	175414	70324	108	406565	406565	360912	0.8877	
		American Passage	4398	1725	6682	2718	4	15527	15527	13784		
		Naches Passage	22723	8911	34526	14269	22	80450	80450	71416		
		American & Naches Passage	27121	10635	41208	16986	26	95977	95977	85200		
		Upper Yakima Passage	88326	34637	134206	53337	82	310588	310588	275712		
Hatchery		Prosser Hatchery Tally	5	2254	126919	101160	171	230509	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	16	7111	225281	153510	680	386599	404834	354470	0.0450	0.8756
McN-UnStr Hatch	Estimate b.	Total Passage	9	3786	213241	169962	288	387287	405555	360015		0.8877
Pooled Str Hatch	Estimate c.	Total Passage	16	6876	235328	155049	2164	399432	418273	366950		0.8773
Pooled UnStr Hatch	Estimate e.	Total Passage	9	3912	220316	175601	298	400136	419010	371959		0.8877

5.6. Year 2003

2003	Brood-Year 2001	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer
Wild	Prosser Wild Tally	30359.49166	16582	98537	33294	272	179045	179045

	American	WDFW Percent	0.13	0.13	0.13	0.16	0.16					
		Estimated Prosser Tally	4078	2227	13236	5338	44	24923	24923			
		WDFW Percent	0.22	0.22	0.22	0.34	0.34					
	Naches	Estimated Prosser Tally	6570	3589	21325	11400	93	42977	42977			
		WDFW Percent	0.65	0.65	0.65	0.50	0.50					
	Upper Yakima	Estimated Prosser Tally	19711.01324	10766	63975	16557	135	111144	111144			
		Yakima Passage Wild Tally	30359	16582	98537	33294	272	179045	Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	45.1%	45.1%	61.9%	54.7%	13.4%					
		Total Passage	67353	36787	159149	60921	2035	326245	326245	308309		0.9450
		American Passage	9047	4941	21378	9767	326	45461	45461	42961		
		Naches Passage	14576	7961	34443	20859	697	78536	78536	74218		
		American & Naches Passage	23624	12903	55821	30626	1023	123997	123997	117180		
		Upper Yakima Passage	43729	23884	103328	30295	1012	202248	202248	191129		
McN UnStr Wild	Estimate b.	Detection Efficiency	58.5%	58.5%	58.5%	58.5%	58.5%					
		Total Passage	51891	28342	168422	56908	466	306029	306029	289106		0.9447
		American Passage	6970	3807	22624	9124	75	42600	42600	40244		
		Naches Passage	11230	6134	36450	19485	159	73458	73458	69395		
		American & Naches Passage	18201	9941	59073	28609	234	116058	116058	109640		
		Upper Yakima Passage	33691	18401	109349	28299	232	189971	189971	179466		
Pooled Str Wild	Estimate c.	Detection Efficiency	47.3%	47.3%	61.3%	51.8%	11.4%					
		Total Passage	64119	35020	160800	64329	2398	326666	326666	308959		0.9458
		American Passage	8613	4704	21600	10314	93	45324	45324	42867		
		Naches Passage	13877	7579	34800	22026	487	78768	78768	74498		
		American & Naches Passage	22490	12283	56400	32339	579	124091	124091	117365		
		Upper Yakima Passage	41630	22737	104400	31990	1819	202575	202575	191594		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	57.1%	57.1%	57.1%	57.1%	57.1%					
		Total Passage	53199	29056	172667	58342	477	313743	313743	296392		0.9447
		American Passage	7146	3903	23194	9354	77	43674	43674	41259		
		Naches Passage	11513	6288	37368	19976	163	75309	75309	71145		
		American & Naches Passage	18659	10191	60562	29330	240	118983	118983	112403		
		Upper Yakima Passage	34540	18865	112105	29013	237	194760	194760	183989		

Hatchery			Prosser Hatchery Tally	0	2058	67386	15896	233	85573	Expanded Elastomer	Expanded PIT	PIT- Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	0	4565	108836	29087	1743	144230	160014	151217	0.0986	0.9450	
McN-UnStr Hatch	Estimate b.	Total Passage	0	3517	115178	27170	399	146264	162271	153297		0.9447	
Pooled Str Hatch	Estimate c.	Total Passage	0	4346	109965	30714	2054	147078	163174	154329		0.9458	
Pooled UnStr Hatch	Estimate e.	Total Passage	0	3605	118081	27855	409	149950	166361	157161		0.9447	

5.7.Year 2004

2004	Brood-Year 2002	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer				
		5652.215										
Wild	Prosser Wild Tally	163	7240	70520	19028	346	102786	102786				
	American	WDFW Percent	0.06	0.04	0.21	0.35	0.31					
		Estimated Prosser Tally	365	309	15160	6607	108	22549	22549			
	Naches	WDFW Percent	0.34	0.29	0.36	0.34	0.19					
		Estimated Prosser Tally	1913	2119	25721	6475	65	36292	36292			
	Upper Yakima	WDFW Percent	0.60	0.66	0.42	0.31	0.50					
		Estimated Prosser Tally	3374.136	048	4812	29639	5946	173	43944	43944		
		Yakima Passage Wild Tally	5652	7240	70520	19028	346	102786	Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	58.4%	58.4%	58.4%	87.2%	87.2%					
		Total Passage	9680	12400	120771	21832	397	165079	165079	171641		1.0398
		American Passage	626	529	25963	7580	124	34822	34822	36206		
		Naches Passage	3276	3629	44049	7429	74	58457	58457	60781		
		American & Naches Passage	3901	4158	70012	15009	198	93280	93280	96987		
		Upper Yakima Passage	5778	8241	50759	6822	198	71799	71799	74653		
McN Str Wild	Estimate b.	Detection Efficiency	64.5%	64.5%	64.5%	64.5%	64.5%					
		Total Passage	8760	11221	109291	29489	536	159296	159296	170539		1.0706
		American Passage	566	479	23495	10239	167	34947	34947	37413		
		Naches Passage	2964	3284	39862	10034	100	56245	56245	60215		
		American & Naches Passage	3531	3763	63357	20274	268	91192	91192	97628		
		Upper Yakima Passage	5229	7458	45934	9215	268	68104	68104	72910		
McN UnStr Wild	Estimate c.	Detection Efficiency	59.4%	59.4%	59.4%	86.8%	86.8%					
		Total Passage	9511	12183	118664	21916	398	162673	162673	169397		1.0413
		American Passage	615	520	25510	7610	124	34379	34379	35800		
		Naches Passage	3219	3566	43281	7458	75	57597	57597	59978		
		American & Naches Passage	3833	4086	68791	15068	199	91976	91976	95778		
		Upper Yakima Passage	5678	8097	49873	6849	199	70696	70696	73619		

Pooled Str Wild	Estimate e.	Detection Efficiency	66.8%	66.8%	66.8%	66.8%	66.8%					
		Total Passage	8465	10843	105611	28496	518	153933	153933	164797	1.0706	
		American Passage	547	463	22704	9894	162	33770	33770	36153		
		Naches Passage	2865	3174	38520	9697	97	54352	54352	58188		
		American & Naches Passage	3412	3636	61224	19591	259	88122	88122	94341		
		Upper Yakima Passage	5053	7207	44387	8905	259	65811	65811	70456		
Pooled UnStr Wild		Prosser Hatchery Tally	0	1662	99011	83912	283	184868	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	0	2847	169565	96276	324	269013	282162	293378	0.0466	1.0398
McN-UnStr Hatch	Estimate b.	Total Passage	0	2576	153446	130045	438	286505	300510	321719		1.0706
Pooled Str Hatch	Estimate c.	Total Passage	0	2797	166606	96651	326	266380	279400	290950		1.0413
Pooled UnStr Hatch	Estimate e.	Total Passage	0	2490	148280	125667	423	276860	290392	310888		1.0706

5.8. Year 2005

2005	Brood-Year 2003	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer				
		37617.03										
Wild	Prosser Wild Tally	993	3569	66596	6246	63	114092	114092				
	American	WDFW Percent	0.21	0.19	0.30	0.32	0.00					
		Estimated Prosser Tally	8047	673	19689	2008	30418	30418				
	Naches	WDFW Percent	0.35	0.08	0.35	0.23	0.18					
		Estimated Prosser Tally	13288	269	23550	1450	38568	38568				
	Upper Yakima	WDFW Percent	0.43	0.74	0.35	0.45	0.82					
		Estimated Prosser Tally	16282.00	236	23357	2789	45106	45106				
		Yakima Passage Wild Tally	37617	3569	66596	6246	63	114092	Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	60.7%	60.7%	71.4%	69.2%	69.2%					
		Total Passage	61931	5876	93219	9028	92	170146	170146	131650		0.7737
		American Passage	13249	1109	27561	2902	0	44820	44820	34679		
		Naches Passage	21876	443	32965	2096	16	57396	57396	44410		
		American & Naches Passage	35125	1552	60525	4998	16	102216	102216	79090		
		Upper Yakima Passage	26806	4324	32694	4030	75	67930	67930	52560		
McN UnStr Wild	Estimate b.	Detection Efficiency	70.0%	70.0%	70.0%	70.0%	70.0%					

		Total Passage	53727	5097	95116	8921	91	162952	162952	125864	0.7724		
		American Passage	11494	962	28121	2868	0	43444	43444	33556			
		Naches Passage	18978	385	33635	2071	16	55085	55085	42548			
		American & Naches Passage	30472	1346	61757	4939	16	98530	98530	76104			
		Upper Yakima Passage	23255	3751	33360	3983	74	64422	64422	49760			
Pooled Str	Wild	Estimate c.	Detection Efficiency	60.1%	60.1%	71.9%	57.1%	57.1%					
			Total Passage	62602	5939	92669	10945	111	172267	172267	134859	0.7828	
			American Passage	13392	1121	27398	3518	0	45429	45429	35564		
			Naches Passage	22113	448	32770	2541	20	57892	57892	45321		
			American & Naches Passage	35506	1569	60168	6059	20	103321	103321	80885		
			Upper Yakima Passage	27096	4370	32501	4886	91	68946	68946	53974		
Pooled UnStr	Wild	Estimate e.	Detection Efficiency	68.4%	68.4%	68.4%	68.4%	68.4%					
			Total Passage	54999	5218	97370	9133	93	166813	166813	128846	0.7724	
			American Passage	11766	985	28788	2936	0	44474	44474	34351		
			Naches Passage	19428	394	34432	2120	17	56390	56390	43556		
			American & Naches Passage	31194	1378	63220	5056	17	100864	100864	77907		
			Upper Yakima Passage	23806	3840	34150	4077	76	65949	65949	50939		
Hatchery			Prosser Hatchery Tally	21	8	159590	37455	16	197090	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.		Total Passage	35	13	223388	54132	24	277593	291340	225424	0.0472	0.7737
McN-UnStr Hatch	Estimate b.		Total Passage	31	11	227934	53495	23	281494	295434	228194		0.7724
Pooled Str Hatch	Estimate c.		Total Passage	36	13	222070	65629	29	287777	302028	236443		0.7828
Pooled UnStr Hatch	Estimate e.		Total Passage	31	11	233334	54762	24	288163	302433	233600		0.7724

5.9.Year 2006

2006	Brood-Year 2004	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer
		10378.78						
Wild	Prosser Wild Tally	788	400	21517	9248	45	41588	41588
	American	WDFW Percent	7.36%	0.00%	5.52%	5.45%	2.27%	
		Estimated Prosser Tally	764	0	1187	504	1	2456
		WDFW Percent	39.88%	25.96%	35.95%	39.11%	15.91%	
	Naches	Estimated Prosser Tally	4139	104	7736	3617	7	15602

		WDFW Percent	52.76%	74.04%	58.53%	55.45%	81.82%					
Upper Yakima		Estimated Prosser Tally	5475.924	893	296	12593	5127	37	23530	23530		
		Yakima Passage Wild Tally	10379	400	21517	9248	45	41588	Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	21.0%	21.0%	21.0%	23.7%	23.7%					
		Total Passage	49335	1901	102278	38999	191	192705	192705	126524		0.6566
		American Passage	3632	0	5644	2124	4	11404	11404	7488		
		Naches Passage	19673	494	36772	15252	30	72222	72222	47419		
		American & Naches Passage	23305	494	42416	17376	35	83626	83626	54906		
		Upper Yakima Passage	26029	1408	59862	21623	156	109079	109079	71618		
McN UnStr Wild	Estimate b.	Detection Efficiency	20.5%	20.5%	20.5%	20.5%	20.5%					
		Total Passage	50510	1947	104715	45005	220	202397	202397	131973		0.6520
		American Passage	3719	0	5779	2451	5	11953	11953	7794		
		Naches Passage	20142	505	37648	17601	35	75932	75932	49511		
		American & Naches Passage	23861	505	43427	20052	40	87885	87885	57305		
		Upper Yakima Passage	26650	1441	61288	24953	180	114512	114512	74667		
Pooled Str Wild	Estimate c.	Detection Efficiency	20.1%	20.1%	20.1%	22.0%	22.0%					
		Total Passage	51735	1994	107254	42031	206	203220	203220	133218		0.6555
		American Passage	3809	0	5919	2289	5	12021	12021	7880		
		Naches Passage	20631	518	38561	16438	33	76180	76180	49939		
		American & Naches Passage	24439	518	44480	18727	37	88201	88201	57819		
		Upper Yakima Passage	27296	1476	62774	23304	168	115019	115019	75399		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	20.7%	20.7%	20.7%	20.7%	20.7%					
		Total Passage	50065	1930	103791	44608	218	200612	200612	130809		0.6520
		American Passage	3686	0	5728	2429	5	11847	11847	7725		
		Naches Passage	19964	501	37316	17446	35	75262	75262	49075		
		American & Naches Passage	23650	501	43044	19875	40	87110	87110	56800		
		Upper Yakima Passage	26415	1429	60747	24733	179	113502	113502	74009		
Hatchery		Prosser Hatchery Tally	3	9	46130	45561	19	91722	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	14	43	219277	192140	81	411555	431559	283348	0.0464	0.6566

McN-UnStr Hatch	Estimate b.	Total Passage	15	44	224500	221728	93	446380	468077	305209	0.6520
Pooled Str Hatch	Estimate c.	Total Passage	15	45	229944	207074	87	437166	458415	300508	0.6555
Pooled UnStr Hatch	Estimate e.	Total Passage	15	44	222520	219773	92	442444	463950	302518	0.6520

5.10. Year 2007

2007		Brood-Year 2005	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer			
			541.5116									
Wild		Prosser Wild Tally	347	523	17147	11159	189	29559	29559			
	American	WDFW Percent	9.10%	14.50%	6.81%	16.75%	11.54%					
		Estimated Prosser Tally	49	76	1167	1869	22	3183	3183			
	Naches	WDFW Percent	18.20%	32.30%	24.72%	29.78%	26.07%					
		Estimated Prosser Tally	99	169	4239	3323	49	7879	7879			
	Upper Yakima	WDFW Percent	72.70%	53.20%	68.47%	53.47%	62.39%					
		Estimated Prosser Tally	584	278	11740	5967	118	18497	18497			
		Yakima Passage Wild Tally	542	523	17147	11159	189	29559	Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	30.2%	30.2%	30.2%	21.9%	21.9%					
		Total Passage	1791	1728	56711	51048	866	112144	112144	99769		0.8897
		American Passage	163	251	3860	8550	100	12924	12924	11498		
		Naches Passage	326	558	14022	15200	226	30332	30332	26985		
		American & Naches Passage	489	809	17882	23750	326	43256	43256	38483		
		Upper Yakima Passage	1302	920	38829	27297	540	68888	68888	61287		
McN UnStr Wild	Estimate b.	Detection Efficiency	26.3%	26.3%	26.3%	26.3%	26.3%					
		Total Passage	2058	1986	65172	42413	719	112349	112349	98319		0.8751
		American Passage	187	288	4436	7104	83	12098	12098	10588		
		Naches Passage	375	642	16114	12629	188	29946	29946	26207		
		American & Naches Passage	562	930	20550	19733	271	42045	42045	36794		
		Upper Yakima Passage	1496	1057	44622	22680	449	70304	70304	61525		
Pooled Str Wild	Estimate c.	Detection Efficiency	28.3%	28.3%	28.3%	23.7%	23.7%					
		Total Passage	1916	1849	60674	47178	800	112417	112417	99265		0.8830
		American Passage	174	268	4130	7902	92	12567	12567	11097		
		Naches Passage	349	597	15001	14048	209	30204	30204	26670		

		American & Naches Passage	523	865	19131	21950	301	42771	42771	37767		
		Upper Yakima Passage	1393	984	41543	25228	499	69646	69646	61498		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	26.2%	26.2%	26.2%	26.2%	26.2%					
		Total Passage	2068	1996	65477	42611	723	112874	112874	98779	0.8751	
		American Passage	188	289	4457	7137	83	12155	12155	10637		
		Naches Passage	376	645	16189	12688	188	30087	30087	26329		
		American & Naches Passage	565	934	20646	19825	272	42241	42241	36967		
		Upper Yakima Passage	1503	1062	44831	22786	451	70633	70633	61813		
Hatchery		Prosser Hatchery Tally	0	629	61236	37776	281	99922	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	0	2079	202534	172814	1285	378712	396759	352979	0.0455	0.8897
McN-UnStr Hatch	Estimate b.	Total Passage	0	2389	232752	143581	1068	379790	397889	348202		0.8751
Pooled Str Hatch	Estimate c.	Total Passage	0	2224	216687	159714	1188	379813	397912	351359		0.8830
Pooled UnStr Hatch	Estimate e.	Total Passage	0	2400	233841	144253	1073	381568	399751	349831		0.8751

5.11. Year 2008

2008	Brood-Year 2006	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer				
		7037.374										
Wild	Prosser Wild Tally	779	1052	44603	16505	443	69641	69641				
	American	WDFW Percent	8.33%	0.00%	5.22%	5.00%	14.81%					
		Estimated Prosser Tally	586	0	2327	825	66	3804	3804			
	Naches	WDFW Percent	8.33%	14.29%	25.22%	31.11%	51.85%					
		Estimated Prosser Tally	586	150	11248	5135	230	17349	17349			
	Upper Yakima	WDFW Percent	83.33%	85.71%	69.57%	63.89%	33.33%					
		Estimated Prosser Tally	983	902	31028	10545	148	48487	48487			
		Yakima Passage Wild Tally	7037	1052	44603	16505	443	69641	Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	71.4%	71.4%	71.4%	35.6%	10.8%					
		Total Passage	9857	1473	62485	46346	4094	124254	124254	107901		0.8684
		American Passage	821	0	3260	2317	606	7005	7005	6083		
		Naches Passage	821	210	15757	14419	2123	33330	33330	28944		
		American & Naches Passage	1643	210	19017	16736	2729	40335	40335	35027		

		Upper Yakima Passage	8214	1263	43468	29610	1365	83919	83919	72874		
McN UnStr Wild	Estimate b.	Detection Efficiency	46.1%	46.1%	46.1%	46.1%	46.1%					
		Total Passage	15257	2281	96703	35784	961	150986	150986	130742	0.8659	
		American Passage	1271	0	5045	1789	142	8248	8248	7142		
		Naches Passage	1271	326	24386	11133	498	37614	37614	32571		
		American & Naches Passage	2543	326	29431	12922	641	45863	45863	39714		
		Upper Yakima Passage	12715	1955	67272	22862	320	105123	105123	91029		
Pooled Str Wild	Estimate c.	Detection Efficiency	48.8%	48.8%	66.7%	31.2%	7.9%					
		Total Passage	14422	2156	66892	52920	5644	142034	142034	123735	0.8712	
		American Passage	1202	0	3490	2646	836	8174	8174	7121		
		Naches Passage	1202	308	16868	16464	2927	37769	37769	32903		
		American & Naches Passage	2404	308	20358	19110	3763	45943	45943	40024		
		Upper Yakima Passage	12018	1848	46534	33810	1881	96091	96091	83711		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	41.4%	41.4%	41.4%	41.4%	41.4%					
		Total Passage	16979	2538	107612	39821	1069	168019	168019	145492	0.8659	
		American Passage	1415	0	5615	1991	158	9179	9179	7948		
		Naches Passage	1415	363	27137	12389	554	41858	41858	36246		
		American & Naches Passage	2830	363	32752	14380	713	51037	51037	44194		
		Upper Yakima Passage	14149	2175	74861	25441	356	116983	116983	101298		
Hatchery		Prosser Hatchery Tally	0	233	43465	65164	930	109793	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	0	326	60890	182980	8595	252791	268938	233543	0.0600	0.8684
McN-UnStr Hatch	Estimate b.	Total Passage	0	505	94235	141281	2017	238037	253242	219289		0.8659
Pooled Str Hatch	Estimate c.	Total Passage	0	477	65185	208936	11851	286449	304746	265485		0.8712
Pooled UnStr Hatch	Estimate e.	Total Passage	0	561	104866	157219	2245	264891	281812	244028		0.8659

5.12. Year 2009

2009	Brood-Year 2007	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer
Wild	Prosser Wild Tally	14956	543	27585	9394	2450	54927	54927
American	WDFW Percent	9.80%	10.93%	12.06%	10.95%	36.29%		
	Estimated Prosser Tally	1466	59	3327	1029	889	6769	6769
Naches	WDFW Percent	35.60%	32.43%	29.25%	40.78%	28.23%		

		Estimated Prosser Tally	5324	176	8068	3831	691	18090	18090			
		WDFW Percent	54.60%	56.64%	58.69%	48.27%	35.48%					
Upper Yakima		Estimated Prosser Tally	8166.224	368	307	16191	4534	869	30067	30067		
		Yakima Passage Wild Tally	14956	543	27585	9394	2450	54927	Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	28.4%	28.4%	21.2%	12.5%	12.5%					
		Total Passage	52671	1911	130062	75334	19645	279622	279622	240827		0.8613
		American Passage	5162	209	15686	8249	7129	36434	36434	31379		
		Naches Passage	18751	620	38038	30723	5545	93676	93676	80680		
		American & Naches Passage	23912	828	53724	38972	12674	130111	130111	112059		
		Upper Yakima Passage	28758	1082	76338	36362	6971	149512	149512	128768		
McN UnStr Wild	Estimate b.	Detection Efficiency	15.3%	15.3%	15.3%	15.3%	15.3%					
		Total Passage	98002	3555	180751	61551	16051	359910	359910	318180		0.8841
		American Passage	9604	388	21799	6740	5825	44356	44356	39213		
		Naches Passage	34889	1153	52863	25102	4530	118537	118537	104793		
		American & Naches Passage	44493	1541	74662	31842	10355	162893	162893	144006		
		Upper Yakima Passage	53509	2014	106089	29710	5695	197017	197017	174173		
Pooled Str Wild	Estimate c.	Detection Efficiency	26.2%	26.2%	21.3%	11.4%	11.4%					
		Total Passage	57137	2073	129580	82196	21434	292419	292419	250846		0.8578
		American Passage	5599	226	15628	9000	7778	38232	38232	32797		
		Naches Passage	20341	672	37897	33521	6050	98481	98481	84480		
		American & Naches Passage	25940	899	53525	42521	13828	136713	136713	117277		
		Upper Yakima Passage	31197	1174	76055	39674	7606	155705	155705	133569		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	14.6%	14.6%	14.6%	14.6%	14.6%					
		Total Passage	102487	3718	189022	64368	16785	376379	376379	332739		0.8841
		American Passage	10044	406	22797	7048	6091	46386	46386	41008		
		Naches Passage	36485	1206	55282	26251	4738	123961	123961	109588		
		American & Naches Passage	46529	1612	78078	33299	10829	170347	170347	150596		
		Upper Yakima Passage	55958	2106	110943	31069	5956	206032	206032	182143		
Hatchery		Prosser Hatchery Tally	31	42	23787	39531	303	63695	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index

McN-Str Hatch	Estimate a.	Total Passage	111	148	112155	317029	2431	431874	454638	391561	0.0501	0.8613
McN-UnStr Hatch	Estimate b.	Total Passage	206	276	155865	259027	1986	417360	439358	388416		0.8841
Pooled Str Hatch	Estimate c.	Total Passage	120	161	111739	345905	2653	460577	484854	415923		0.8578
Pooled UnStr Hatch	Estimate e.	Total Passage	216	288	162997	270879	2077	436457	459463	406189		0.8841

5.13. Year 2010

2010	Brood-Year 2008	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer				
Wild	Prosser Wild Tally	3862	3204	70483	24871	637	103056	103056				
	American	WDFW Percent	30.31%	0.00%	14.16%	11.88%	0.00%					
		Estimated Prosser Tally	1170	0	9981	2955	0	14106	14106			
	Naches	WDFW Percent	7.35%	19.50%	37.13%	33.63%	75.49%					
		Estimated Prosser Tally	284	625	26167	8364	481	35921	35921			
	Upper Yakima	WDFW Percent	62.34%	80.50%	48.71%	54.49%	24.51%					
		Estimated Prosser Tally	2407.390	06	2579	34334	13552	156	53029	53029		
		Yakima Passage Wild Tally	3862	3204	70483	24871	637	103056	Expanded Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	45.0%	45.0%	45.0%	59.2%	43.6%					
		Total Passage	8584	7122	156665	42045	1459	215875	215875	221188		1.0246
		American Passage	2602	0	22186	4995	0	29782	29782	30515		
		Naches Passage	631	1389	58163	14140	1101	75424	75424	77281		
		American & Naches Passage	3233	1389	80349	19135	1101	105206	105206	107796		
		Upper Yakima Passage	5351	5733	76316	22910	358	110668	110668	113392		
McN UnStr Wild	Estimate b.	Detection Efficiency	52.2%	52.2%	52.2%	52.2%	52.2%					
		Total Passage	7396	6137	134998	47635	1219	197386	197386	201737		1.0220
		American Passage	2242	0	19117	5659	0	27018	27018	27614		
		Naches Passage	544	1197	50119	16020	921	68800	68800	70316		
		American & Naches Passage	2785	1197	69236	21679	921	95818	95818	97930		
		Upper Yakima Passage	4611	4940	65761	25956	299	101568	101568	103807		
Pooled Str Wild	Estimate c.	Detection Efficiency	45.4%	45.4%	45.4%	57.4%	35.4%					
		Total Passage	8507	7058	155261	43333	1796	215955	215955	221228		1.0244
		American Passage	2578	0	21987	5148	0	29713	29713	30439		
		Naches Passage	625	1377	57642	14573	1356	75572	75572	77418		

		American & Naches Passage	3204	1377	79629	19721	1356	105285	105285	107856		
		Upper Yakima Passage	5303	5682	75632	23612	440	110669	110669	113372		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	51.3%	51.3%	51.3%	51.3%	51.3%					
		Total Passage	7530	6248	137440	48497	1241	200957	200957	205387	1.0220	
		American Passage	2282	0	19463	5761	0	27507	27507	28113		
		Naches Passage	553	1219	51026	16310	937	70044	70044	71588		
		American & Naches Passage	2836	1219	70489	22071	937	97551	97551	99702		
		Upper Yakima Passage	4694	5030	66951	26426	304	103406	103406	105685		
Hatchery		Prosser Hatchery Tally	0	204	58305	129493	737	188739	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	0	453	129598	218915	1688	350653	367535	376582	0.0459	1.0246
McN-UnStr Hatch	Estimate b.	Total Passage	0	390	111674	248021	1411	361496	378900	387253		1.0220
Pooled Str Hatch	Estimate c.	Total Passage	0	449	128436	225621	2078	356584	373751	382878		1.0244
Pooled UnStr Hatch	Estimate e.	Total Passage	0	397	113694	252508	1436	368036	385755	394259		1.0220

5.14. Year 2011

2011	Brood-Year 2009	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer				
Wild	Prosser Wild Tally	24773	4142	30530	15792	91	75328	75328				
	American	WDFW Percent	8.64%	0.00%	3.49%	5.92%	16.65%					
		Estimated Prosser Tally	2140	0	1066	935	15	4156	4156			
		WDFW Percent	18.19%	19.75%	23.96%	13.10%	0.00%					
	Naches	Estimated Prosser Tally	4506	818	7316	2069	0	14709	14709			
		WDFW Percent	73.17%	80.25%	72.55%	80.98%	83.35%					
	Upper Yakima	Estimated Prosser Tally	18126.20	455	3324	22149	12788	75	56463	56463		
		Yakima Passage Wild Tally	24773	4142	30530	15792	91	75328	Expanded Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	17.5%	17.5%	28.7%	30.9%	30.9%					
		Total Passage	141442	23652	106452	51115	293	322954	322954	299949		0.9288
		American Passage	12221	0	3716	3027	49	19012	19012	17657		
		Naches Passage	25728	4671	25508	6697	0	62605	62605	58146		
		American & Naches Passage	37949	4671	29224	9724	49	81617	81617	75803		

		Upper Yakima Passage	103493	18980	77228	41391	244	241337	241337	224146		
McN UnStr Wild	Estimate b.	Detection Efficiency	27.9%	27.9%	27.9%	27.9%	27.9%					
		Total Passage	88870	14861	109524	56652	325	270231	270231	254125	0.9404	
		American Passage	7678	0	3823	3355	54	14910	14910	14021		
		Naches Passage	16165	2935	26245	7423	0	52768	52768	49623		
		American & Naches Passage	23844	2935	30067	10777	54	67678	67678	63644		
		Upper Yakima Passage	65026	11926	79457	45875	271	202554	202554	190481		
Pooled Str Wild	Estimate c.	Detection Efficiency	17.6%	17.6%	28.3%	29.5%	29.5%					
		Total Passage	140705	23528	107826	53479	307	325846	325846	303711	0.9321	
		American Passage	12157	0	3764	3167	51	19138	19138	17838		
		Naches Passage	25594	4647	25838	7007	0	63086	63086	58800		
		American & Naches Passage	37751	4647	29601	10174	51	82224	82224	76639		
		Upper Yakima Passage	102954	18882	78225	43306	256	243622	243622	227072		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	27.3%	27.3%	27.3%	27.3%	27.3%					
		Total Passage	90699	15166	111779	57819	332	275795	275795	259357	0.9404	
		American Passage	7836	0	3901	3424	55	15217	15217	14310		
		Naches Passage	16498	2995	26785	7576	0	53854	53854	50644		
		American & Naches Passage	24335	2995	30686	10999	55	69071	69071	64954		
		Upper Yakima Passage	66365	12171	81093	46819	276	206724	206724	194403		
Hatchery		Prosser Hatchery Tally	70	4100	57391	66684	580	128824	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	398	23409	200108	215843	1877	441635	461721	428831	0.0435	0.9288
McN-UnStr Hatch	Estimate b.	Total Passage	250	14708	205884	239222	2080	462144	483164	454365		0.9404
Pooled Str Hatch	Estimate c.	Total Passage	396	23287	202692	225825	1963	454164	474820	442564		0.9321
Pooled UnStr Hatch	Estimate e.	Total Passage	255	15011	210123	244147	2123	471659	493111	463720		0.9404

5.15. Year 2012

2012	Brood-Year 2010	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer
Wild	Prosser Wild Tally	15922	6786	14719	5327	993	43746	43746
	American	WDFW Percent	10.99%	5.31%	6.17%	13.65%	23.46%	
		Estimated Prosser Tally	1750	360	908	727	3978	3978
	Naches	WDFW Percent	31.62%	29.60%	29.32%	38.48%	29.45%	

		Estimated Prosser Tally	5034	2009	4316	2050	292	13700	13700			
		WDFW Percent	57.39%	65.09%	64.51%	47.87%	47.09%					
Upper Yakima		Estimated Prosser Tally	9138.041									
		Estimated Prosser Tally	429	4416	9495	2550	468	26067	26067			
		Yakima Passage Wild Tally	15922	6786	14719	5327	993	43746	Expanded Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	10.6%	10.6%	6.8%	6.4%	6.4%					
		Total Passage	149599	63757	215132	82800	15434	526721	526721	301173		0.5718
		American Passage	16439	3386	13274	11299	3621	48019	48019	27456		
		Naches Passage	47298	18874	63077	31863	4545	165658	165658	94721		
		American & Naches Passage	63738	22260	76350	43162	8166	213676	213676	122178		
		Upper Yakima Passage	85861	41497	138782	39638	7267	313045	313045	178995		
McN UnStr Wild	Estimate b.	Detection Efficiency	6.8%	6.8%	6.8%	6.8%	6.8%					
		Total Passage	233096	99343	215485	77987	14537	640449	640449	368824		0.5759
		American Passage	25615	5276	13295	10642	3411	58239	58239	33539		
		Naches Passage	73698	29408	63180	30011	4281	200579	200579	115510		
		American & Naches Passage	99312	34684	76476	40654	7692	258818	258818	149049		
		Upper Yakima Passage	133784	64659	139010	37334	6845	381631	381631	219775		
Pooled Str Wild	Estimate c.	Detection Efficiency	17.2%	12.0%	8.0%	6.2%	6.2%					
		Total Passage	92790	56530	184609	86385	16102	436417	436417	252029		0.5775
		American Passage	10197	3002	11390	11788	3778	40155	40155	23189		
		Naches Passage	29337	16735	54127	33243	4742	138184	138184	79801		
		American & Naches Passage	39534	19737	65518	45031	8520	178339	178339	102990		
		Upper Yakima Passage	53256	36794	119091	41354	7582	258077	258077	149038		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	7.4%	7.4%	7.4%	7.4%	7.4%					
		Total Passage	216431	92241	200080	72412	13497	594661	594661	342455		0.5759
		American Passage	23783	4898	12345	9881	3167	54075	54075	31141		
		Naches Passage	68429	27306	58663	27866	3975	186239	186239	107252		
		American & Naches Passage	92212	32204	71008	37747	7142	240314	240314	138393		
		Upper Yakima Passage	124219	60036	129071	34665	6356	354347	354347	204063		
Hatchery		Prosser Hatchery Tally	0	1485	20279	22395	919	45078	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index

McN-Str Hatch	Estimate a.	Total Passage	0	13952	296397	348103	14288	672740	707207	404372	0.0487	0.5718
McN-UnStr Hatch	Estimate b.	Total Passage	0	21739	296884	327872	13457	659952	693764	399527		0.5759
Pooled Str Hatch	Estimate c.	Total Passage	0	12370	254344	363177	14906	644798	677833	391446		0.5775
Pooled UnStr Hatch	Estimate e.	Total Passage	0	20185	275659	304431	12495	612770	644164	370963		0.5759

5.16. Year 2013

2013	Brood-Year 2011	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer				
Wild	Prosser Wild Tally	28502	18683	50994	8258	336	106774	106774				
	American	WDFW Percent	8.23%	2.30%	5.72%	16.96%	6.39%					
		Estimated Prosser Tally	2346	429	2916	1401	22	7113	7113			
	Naches	WDFW Percent	17.43%	20.59%	27.50%	29.53%	7.85%					
		Estimated Prosser Tally	4968	3847	14023	2439	26	25303	25303			
	Upper Yakima	WDFW Percent	74.34%	77.11%	66.78%	53.51%	85.76%					
		Estimated Prosser Tally	21188.49	724	14407	34055	4419	289	74358	74358		
		Yakima Passage Wild Tally	28502	18683	50994	8258	336	106774	Expanded Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	26.7%	26.7%	37.1%	23.4%	23.4%					
		Total Passage	106741	69970	137366	35270	1437	350785	350785	358055		1.0207
		American Passage	8785	1608	7855	5982	92	24321	24321	24826		
		Naches Passage	18605	14408	37774	10415	113	81314	81314	82999		
		American & Naches Passage	27390	16016	45628	16397	205	105636	105636	107825		
		Upper Yakima Passage	79352	53955	91738	18873	1232	245149	245149	250230		
McN UnStr Wild	Estimate b.	Detection Efficiency	32.6%	32.6%	32.6%	32.6%	32.6%					
		Total Passage	87352	57260	156284	25309	1031	327236	327236	333839		1.0202
		American Passage	7189	1316	8936	4293	66	21800	21800	22240		
		Naches Passage	15225	11791	42976	7474	81	77546	77546	79111		
		American & Naches Passage	22415	13106	51912	11766	147	99346	99346	101351		
		Upper Yakima Passage	64938	44154	104372	13543	884	227890	227890	232489		
Pooled Str Wild	Estimate c.	Detection Efficiency	27.5%	27.5%	35.1%	21.1%	21.1%					
		Total Passage	103702	67978	145428	39056	1591	357755	357755	365468		1.0216
		American Passage	8535	1562	8316	6624	102	25139	25139	25680		
		Naches Passage	18075	13997	39991	11533	125	83721	83721	85526		

		American & Naches Passage	26610	15560	48306	18157	227	108860	108860	111206		
		Upper Yakima Passage	77092	52418	97122	20898	1365	248896	248896	254261		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	30.5%	30.5%	30.5%	30.5%	30.5%					
		Total Passage	93410	61231	167121	27064	1103	349929	349929	356990	1.0202	
		American Passage	7688	1407	9556	4590	70	23312	23312	23782		
		Naches Passage	16281	12608	45956	7992	87	82924	82924	84597		
		American & Naches Passage	23969	14015	55512	12582	157	106235	106235	108379		
		Upper Yakima Passage	69441	47216	111609	14482	946	243693	243693	248611		
Hatchery		Prosser Hatchery Tally	0	13014	69719	20263	879	103874	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	0	48738	187807	86542	3753	326839	343892	351019	0.0496	1.0207
McN-UnStr Hatch	Estimate b.	Total Passage	0	39885	213671	62100	2693	318349	334959	341718		1.0202
Pooled Str Hatch	Estimate c.	Total Passage	0	47350	198830	95831	4155	346166	364227	372079		1.0216
Pooled UnStr Hatch	Estimate e.	Total Passage	0	42651	228489	66406	2879	340425	358187	365415		1.0202

5.17.Year 2014

2014	Brood-Year 2012	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer				
Wild	Prosser Wild Tally	1589	4340	14949	11897	959	33735	33735				
	American	WDFW Percent	11.65%	12.03%	9.09%	11.95%	13.86%					
		Estimated Prosser Tally	185	522	1360	1421	133	3621	3621			
	Naches	WDFW Percent	41.19%	21.74%	30.16%	38.12%	0.00%					
		Estimated Prosser Tally	655	944	4509	4535	0	10643	10643			
	Upper Yakima	WDFW Percent	47.16%	66.23%	60.74%	49.93%	86.14%					
		Estimated Prosser Tally	749.6015 614	2874	9080	5940	826	19471	19471			
		Yakima Passage Wild Tally	1589	4340	14949	11897	959	33735	Expanded Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	13.9%	13.9%	13.9%	13.9%	6.0%					
		Total Passage	11447	31257	107660	85679	15923	251966	251966	250881		0.9957
		American Passage	1334	3760	9791	10236	2208	27329	27329	27211		
		Naches Passage	4715	6795	32474	32662	0	76646	76646	76317		
		American & Naches Passage	6049	10555	42266	42898	2208	103975	103975	103528		
		Upper Yakima Passage	5398	20701	65395	42781	13715	147991	147991	147354		
McN UnStr Wild	Estimate b.	Detection Efficiency	13.8%	13.8%	13.8%	13.8%	13.8%					
		Total Passage	11481	31349	107976	85931	6930	243667	243667	241676		0.9918
		American Passage	1338	3771	9820	10266	961	26156	26156	25942		
		Naches Passage	4729	6815	32570	32758	0	76872	76872	76244		
		American & Naches Passage	6066	10586	42390	43024	961	103027	103027	102186		
		Upper Yakima Passage	5414	20762	65587	42907	5969	140639	140639	139490		
Pooled Str Wild	Estimate c.	Detection Efficiency	13.1%	13.1%	13.1%	13.1%	5.0%					
		Total Passage	12091	33016	113718	90500	19031	268355	268355	267433		0.9966
		American Passage	1409	3972	10342	10812	2638	29173	29173	29073		
		Naches Passage	4980	7178	34302	34500	0	80959	80959	80681		
		American & Naches Passage	6389	11149	44644	45312	2638	110132	110132	109754		
		Upper Yakima Passage	5702	21866	69074	45188	16392	158223	158223	157679		
Pooled UnStr Wild	Estimate e.	Total Passage	13.0%	13.0%	13.0%	13.0%	13.0%					

		Total Passage	12197	33306	114717	91295	7363	258877	258877	256762	0.9918	
		American Passage	1421	4007	10433	10907	1021	27788	27788	27561		
		Naches Passage	5024	7241	34603	34803	0	81670	81670	81003		
		American & Naches Passage	6445	11247	45036	45710	1021	109459	109459	108564		
		Upper Yakima Passage	5752	22058	69681	45585	6342	149419	149419	148198		
Hatchery		Prosser Hatchery Tally	0	1493	16126	30753	1114	49486	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	0	10749	116139	221480	18480	366847	385256	383598	0.0478	0.9957
McN-UnStr Hatch	Estimate b.	Total Passage	0	10781	116480	222131	8043	357434	375371	372304		0.9918
Pooled Str Hatch	Estimate c.	Total Passage	0	11354	122673	233942	22087	390056	409630	408222		0.9966
Pooled UnStr Hatch	Estimate e.	Total Passage	0	11454	123751	235997	8545	379747	398803	395545		0.9918

5.18. Year 2015

2015	Brood-Year 2013	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer				
Wild	Prosser Wild Tally	2658	13541	35320	11639	4	63162	63162				
	American	WDFW Percent	13.86%	11.62%	8.92%	14.74%	14.74%					
		Estimated Prosser Tally	368	1573	3149	1716	1	6807	6807			
		WDFW Percent	16.80%	26.32%	23.13%	24.09%	24.09%					
	Naches	Estimated Prosser Tally	447	3564	8169	2804	1	14985	14985			
	Upper Yakima	WDFW Percent	69.34%	62.06%	67.96%	61.17%	61.17%					
		Estimated Prosser Tally	1842.998005	8404	24002	7119	2	41370	41370			
		Yakima Passage Wild Tally	2658	13541	35320	11639	4	63162	Expanded Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	52.9%	52.9%	52.9%	56.3%	56.3%					
		Total Passage	5028	25614	66809	20689	6	118146	118146	120848		1.0229
		American Passage	697	2976	5956	3050	1	12680	12680	12970		
		Naches Passage	845	6742	15451	4985	2	28024	28024	28665		
		American & Naches Passage	1541	9718	21408	8035	3	40704	40704	41635		

		Upper Yakima Passage	3486	15897	45401	12655	4	77442	77442	79213		
McN UnStr Wild	Estimate b.	Detection Efficiency	53.2%	53.2%	53.2%	53.2%	53.2%					
		Total Passage	4999	25468	66427	21890	7	118791	118791	121334	1.0214	
		American Passage	693	2959	5922	3227	1	12802	12802	13076		
		Naches Passage	840	6703	15363	5274	2	28182	28182	28786		
		American & Naches Passage	1533	9662	21285	8501	3	40984	40984	41861		
		Upper Yakima Passage	3466	15806	45141	13389	4	77807	77807	79472		
Pooled Str Wild	Estimate c.	Detection Efficiency	37.1%	37.1%	62.1%	57.6%	57.6%					
		Total Passage	7170	36531	56858	20221	6	120786	120786	123289	1.0207	
		American Passage	994	4244	5069	2981	1	13289	13289	13564		
		Naches Passage	1205	9615	13150	4872	2	28843	28843	29441		
		American & Naches Passage	2198	13859	18219	7853	2	42132	42132	43005		
		Upper Yakima Passage	4972	22671	38639	12368	4	78654	78654	80284		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	51.4%	51.4%	51.4%	51.4%	51.4%					
		Total Passage	5173	26355	68741	22653	7	122930	122930	125561	1.0214	
		American Passage	717	3062	6129	3339	1	13248	13248	13531		
		Naches Passage	869	6937	15898	5458	2	29164	29164	29788		
		American & Naches Passage	1586	9999	22027	8797	3	42412	42412	43320		
		Upper Yakima Passage	3587	16356	46714	13856	4	80518	80518	82241		
Hatchery		Prosser Hatchery Tally	0	43016	90070	26254	11	159351	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	0	81366	170371	46668	19	298424	317197	324451	0.0592	1.0229
McN-UnStr Hatch	Estimate b.	Total Passage	0	80901	169397	49377	21	299696	318550	325368		1.0214
Pooled Str Hatch	Estimate c.	Total Passage	0	116043	144995	45612	19	306669	325961	332715		1.0207
Pooled UnStr Hatch	Estimate e.	Total Passage	0	83720	175300	51098	21	310139	329649	336705		1.0214

5.19. Year 2016

2016	Brood-Year 2014	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer
------	-----------------	-----------	-------	-------	-----	----------	-------	--------------------

YKFP Project Year 2020 M&E Annual Report, Appendix C, Chandler Certification

Wild		Prosser Wild Tally	2900	3922	4227	3478	73	14599	14599			
American	WDFW Percent	5.69%	7.42%	9.44%	13.00%	3.71%						
	Estimated Prosser Tally	165	291	399	452	3	1310	1310				
Naches	WDFW Percent	26.41%	23.18%	38.42%	34.52%	0.00%						
	Estimated Prosser Tally	766	909	1624	1200	0	4500	4500				
Upper Yakima	WDFW Percent	67.90%	69.40%	52.13%	52.49%	96.29%						
	Estimated Prosser Tally	1968.880324	2722	2204	1825	70	8790	8790				
		Yakima Passage Wild Tally	2900	3922	4227	3478	73	14599	Expanded Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	5.5%	5.5%	5.5%	22.8%	22.8%					
		Total Passage	52843	71469	77035	15257	320	216925	216925	51305		0.2365
		American Passage	3007	5304	7273	1983	12	17578	17578	4157		
		Naches Passage	13956	16568	29600	5266	0	65391	65391	15465		
		American & Naches Passage	16963	21872	36873	7250	12	82969	82969	19623		
		Upper Yakima Passage	35881	49598	40162	8008	308	133956	133956	31682		
McN UnStr Wild	Estimate b.	Detection Efficiency	9.6%	9.6%	9.6%	9.6%	9.6%					
		Total Passage	30115	40730	43902	36116	757	151620	151620	39037		0.2575
		American Passage	1714	3022	4145	4694	28	13603	13603	3502		
		Naches Passage	7953	9442	16869	12466	0	46731	46731	12031		
		American & Naches Passage	9667	12465	21014	17161	28	60334	60334	15534		
		Upper Yakima Passage	20448	28265	22888	18956	729	91286	91286	23503		
Pooled Str Wild	Estimate c.	Detection Efficiency	5.9%	5.9%	4.4%	21.5%	21.5%					
		Total Passage	49149	66473	96748	16177	339	228887	228887	53478		0.2336
		American Passage	2797	4933	9134	2103	13	18979	18979	4434		
		Naches Passage	12980	15410	37175	5584	0	71149	71149	16624		
		American & Naches Passage	15777	20343	46309	7687	13	90128	90128	21058		
		Upper Yakima Passage	33372	46131	50439	8491	326	138759	138759	32420		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	8.4%	8.4%	8.4%	8.4%	8.4%					
		Total Passage	34538	46712	50350	41421	868	173890	173890	44770		0.2575
		American Passage	1965	3466	4754	5384	32	15601	15601	4017		
		Naches Passage	9122	10829	19347	14297	0	53594	53594	13799		

			American & Naches Passage	11087	14295	24100	19681	32	69196	69196	17815	
			Upper Yakima Passage	23451	32417	26250	21740	836	104694	104694	26955	
Hatchery		Prosser Hatchery Tally	0	9155	14039	20515	66	136488	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	0	166846	255836	90006	289	1499037	1587340	375419	0.0556	0.2365
McN-UnStr Hatch	Estimate b.	Total Passage	0	95085	145799	213058	685	1417512	1501013	386455		0.2575
Pooled Str Hatch	Estimate c.	Total Passage	0	155183	321302	95434	307	1632683	1728859	403938		0.2336
Pooled UnStr Hatch	Estimate e.	Total Passage	0	109051	167214	244352	785	1625716	1721481	443217		0.2575

5.20.Year 2017

2017	Brood-Year 2015	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer				
Wild	Prosser Wild Tally	2542	458	993	1352	24	5369	5369				
	American	WDFW Percent	10.20%	11.21%	15.80%	10.78%	37.16%					
		Estimated Prosser Tally	296	440	668	375	27	1805	1805			
		WDFW Percent	31.70%	27.73%	27.10%	29.57%	11.47%					
	Naches	Estimated Prosser Tally	919	1087	1146	1028	8	4189	4189			
		WDFW Percent	58.10%	61.06%	57.10%	59.65%	51.37%					
	Upper Yakima	Estimated Prosser Tally	1684.712029	2395	2414	2074	37	8605	8605			
		Yakima Passage Wild Tally	2900	3922	4227	3478	73	14599	Expanded Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	5.5%	5.5%	5.5%	9.3%	9.3%					
		Total Passage	45879	8257	17922	14554	258	86871	86871	60411		0.6954
		American Passage	4680	926	2832	1569	96	10102	10102	7025		
		Naches Passage	14544	2289	4857	4304	30	26024	26024	18097		
		American & Naches Passage	19223	3215	7688	5873	126	36125	36125	25122		
		Upper Yakima Passage	26656	5042	10233	8682	133	50745	50745	35289		
McN UnStr Wild	Estimate b.	Detection Efficiency	7.2%	7.2%	7.2%	7.2%	7.2%					
		Total Passage	35465	6383	13854	18862	335	74899	74899	49700		0.6636
		American Passage	3617	716	2189	2033	124	8679	8679	5759		

		Naches Passage	11242	1770	3754	5578	38	22383	22383	14853		
		American & Naches Passage	14860	2485	5943	7611	163	31062	31062	20612		
		Upper Yakima Passage	20605	3897	7910	11251	172	43836	43836	29088		
Pooled Str Wild	Estimate c.	Detection Efficiency	5.9%	5.9%	5.9%	9.7%	9.7%					
		Total Passage	43257	7785	16897	14009	249	82198	82198	57051	0.6941	
		American Passage	4412	873	2670	1510	92	9557	9557	6633		
		Naches Passage	13712	2159	4579	4143	29	24622	24622	17089		
		American & Naches Passage	18125	3031	7249	5653	121	34179	34179	23723		
		Upper Yakima Passage	25132	4754	9648	8357	128	48019	48019	33328		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	7.6%	7.6%	7.6%	7.6%	7.6%					
		Total Passage	33442	6019	13064	17786	316	70627	70627	46866	0.6636	
		American Passage	3411	675	2064	1917	117	8184	8184	5431		
		Naches Passage	10601	1669	3540	5260	36	21107	21107	14006		
		American & Naches Passage	14012	2344	5604	7177	154	29291	29291	19436		
		Upper Yakima Passage	19430	3675	7459	10609	162	41336	41336	27429		
Hatchery		Prosser Hatchery Tally	1	235	1943	5727	41	7947	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	18	4241	35067	61646	441	386839	412204	286652	0.061	0.6954
McN-UnStr Hatch	Estimate b.	Total Passage	9	3279	27108	79893	572	425176	453055	300633	0.1029	0.6636
Pooled Str Hatch	Estimate c.	Total Passage	12	3999	33063	59338	425	369465	393691	273248	0.1029	0.6941
Pooled UnStr Hatch	Estimate e.	Total Passage	9	3092	25561	75336	539	400926	427215	283486	0.1029	0.6636

5.21.Year 2018

2018	Brood-Year 2016	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer
Wild	Prosser Wild Tally	6091	1173	8517	1374	96	17251	17251
	American	WDFW Percent	8.80%	3.30%	5.82%	10.40%	25.00%	0.00
		Estimated Prosser Tally	255	129	246	362	18	1010
	Naches	WDFW Percent	31.70%	27.73%	27.10%	29.57%	11.47%	0.00

		Estimated Prosser Tally	919	1087	1146	1028	8	4189	4189			
Upper Yakima	WDFW Percent	58.10%	61.06%	57.10%	59.65%	51.37%	0.00					
	Estimated Prosser Tally	1684.712029	2395	2414	2074	37	8605	8605				
		Yakima Passage Wild Tally	2859	3612	3805	3464	64	13804	Expanded Elastomer	Calibrated Total	PIT- Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	9.8%	9.8%	9.8%	4.9%	4.9%					
		Total Passage	62211	11978	86996	27928	1951	191064	191064	128380		0.6719
		American Passage	5475	395	5061	2904	488	14323	14323	9624		
		Naches Passage American & Naches Passage	19721	3321	23576	8259	224	55101	55101	37024		
		Upper Yakima Passage	25196	3716	28637	11164	712	69424	69424	46647		
		Upper Yakima Passage	36145	7314	49674	16659	1002	110794	110794	74445		
McN UnStr Wild	Estimate b.	Detection Efficiency	8.4%	8.4%	8.4%	8.4%	8.4%					
		Total Passage	72640	13986	101579	16386	1145	205735	205735	122910		0.5974
		American Passage	6392	462	5909	1704	286	14753	14753	8814		
		Naches Passage American & Naches Passage	23027	3878	27528	4846	131	59410	59410	35493		
		Upper Yakima Passage	29419	4339	33437	6550	418	74163	74163	44307		
		Upper Yakima Passage	42204	8540	58001	9774	588	119107	119107	71157		
Pooled Str Wild	Estimate c.	Detection Efficiency	13.7%	13.7%	9.3%	4.4%	4.4%					
		Total Passage	44443	8557	91787	30928	2161	177875	177875	131489		0.7392
		American Passage	3911	282	5340	3216	540	13289	13289	9824		
		Naches Passage American & Naches Passage	14088	2373	24874	9147	248	50730	50730	37500		
		Upper Yakima Passage	17999	2655	30214	12363	788	64019	64019	47324		
		Upper Yakima Passage	25821	5225	52410	18448	1110	103015	103015	76150		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	8.2%	8.2%	8.2%	8.2%	8.2%					
		Total Passage	74408	14326	104052	16785	1173	210744	210744	136769		0.6490
		American Passage	6548	473	6053	1745	293	15112	15112	9808		
		Naches Passage American & Naches Passage	23587	3972	28198	4964	135	60856	60856	39495		
		Upper Yakima Passage	30135	4445	34251	6709	428	75969	75969	49302		
		Upper Yakima Passage	43231	8748	59413	10012	602	122007	122007	79180		
Hatchery		Prosser Hatchery Tally	0	1470	15058	2640	392	19560	Expanded Elastomer	Expanded PIT	PIT- Tag/Total	Calibration Index

McN-Str Hatch	Estimate a.	Total Passage	0	15011	153802	53661	7968	386839	411667	276607	0.0603	0.6719
McN-UnStr Hatch	Estimate b.	Total Passage	0	17527	179584	31484	4675	425176	452465	270311		0.5974
Pooled Str Hatch	Estimate c.	Total Passage	0	10724	162273	59425	8824	369465	393178	290644		0.7392
Pooled UnStr Hatch	Estimate e.	Total Passage	0	17954	183956	32251	4789	400926	426658	276892		0.6490

5.22.Year 2019

2019	Brood-Year 2017	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer				
Wild	Prosser Wild Tally	15489	3937	10596	23290	63	53374	53374				
	American	WDFW Percent	9.90%	12.44%	14.70%	14.71%	0.00%	0.00%				
		Estimated Prosser Tally	287	488	621	511	0	1908	1908			
	Naches	WDFW Percent	20.00%	20.33%	22.70%	30.22%	0.00%	0.00%				
		Estimated Prosser Tally	580	797	959	1051	0	3387	3387			
	Upper Yakima	WDFW Percent	76.22%	73.17%	74.47%	66.19%	100.0%	0.00%				
		Estimated Prosser Tally	2,210	2,870	3,148	2,302	73	10,602	10,602			
		Yakima Passage Wild Tally	3077	4154	4729	3864	73	15897	Expanded Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	18.5%	18.5%	18.5%	39.6%	39.6%					
		Total Passage	83,879	21,319	57,385	58,761	158	221,503	221,503	168,119		0.7590
		American Passage	8,305	2,652	8,434	8,641	-	28,032	28,032	21,276		
		Naches Passage	16,776	4,333	13,024	17,755	-	51,888	51,888	39,382		
		American & Naches Passage	25,081	6,985	21,457	26,397	-	79,919	79,919	60,658		
		Upper Yakima Passage	63,930	15,600	42,734	38,892	158	161,313	161,313	122,435		
McN UnStr Wild	Estimate b.	Detection Efficiency	27.1%	27.1%	27.1%	27.1%	27.1%					

		Total Passage	57,169	14,530	39,111	85,963	231	197,005	197,005	154,848	0.7860		
		American Passage	5,660	1,807	5,748	12,642	-	25,857	25,857	20,324			
		Naches Passage	11,434	2,953	8,876	25,974	-	49,238	49,238	38,701			
		American & Naches Passage	17,094	4,761	14,624	38,616	-	75,095	75,095	59,025			
		Upper Yakima Passage	43,572	10,632	29,126	56,896	231	140,457	140,457	110,401			
Pooled Str	Wild	Estimate c.	Detection Efficiency	20.1%	20.1%	20.1%	35.9%	35.9%					
			Total Passage	77,184	19,618	52,827	64,908	175	214,712	214,712	175,427	0.8170	
			American Passage	7,642	2,440	7,764	9,545	-	27,391	27,391	22,379		
			Naches Passage	15,437	3,987	11,989	19,613	-	51,026	51,026	41,690		
			American & Naches Passage	23,079	6,427	19,753	29,158	-	78,417	78,417	64,069		
			Upper Yakima Passage	58,827	14,354	39,340	42,961	175	155,656	155,656	127,176		
Pooled UnStr	Wild	Estimate e.	Detection Efficiency	27.9%	27.9%	27.9%	27.9%	27.9%					
			Total Passage	55,458	14,095	37,941	83,390	224	191,108	191,108	154,530	0.8086	
			American Passage	5,491	1,753	5,576	12,263	-	25,083	25,083	20,282		
			Naches Passage	11,092	2,865	8,611	25,197	-	47,764	47,764	38,622		
			American & Naches Passage	16,582	4,618	14,187	37,460	-	72,847	72,847	58,904		
			Upper Yakima Passage	42,268	10,314	28,254	55,193	224	136,253	136,253	110,174		
Hatchery			Prosser Hatchery Tally	-	904	24,775	76,824	198	102,701	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	-	4,897	134,169	193,833	500	386,839	409,539	310,836	0.0554	0.7590	
McN-UnStr Hatch	Estimate b.	Total Passage	-	3,337	91,444	283,561	732	425,176	450,126	353,803		0.7860	
Pooled Str Hatch	Estimate c.	Total Passage	-	4,506	123,513	214,108	552	369,465	391,145	319,579		0.8170	

Pooled UnStr Hatch	Estimate e.	Total Passage	-	3,237	88,707	275,073	710	400,926	424,452	343,212	0.8086
--------------------	-------------	---------------	---	-------	--------	---------	-----	---------	---------	---------	--------

5.23. Year 2020

2020	Brood-Year 2017	Pre-March	March	April	May	Post-May	Total	Expanded Elastomer			
Wild	Prosser Wild Tally	8843	2602	30737	10851	58	53092	53092			
American	WDFW Percent	3.78%	6.50%	2.84%	3.60%	0.00%	0.00%				
	Estimated Prosser Tally	110	255	120	125	0	610	610			
Naches	WDFW Percent	20.00%	20.33%	22.70%	30.22%	0.00%	0.00%				
	Estimated Prosser Tally	580	797	959	1051	0	3387	3387			
Upper Yakima	WDFW Percent	76.22%	76.22%	76.22%	76.22%	76.2%	76.22%				
	Estimated Prosser Tally	2,210	2,989	3,222	2,650	56	11,127	11,127			
	Yakima Passage Wild Tally	2900	4041	4301	3826	56	15124	Expanded Elastomer	Calibrated Total	PIT-Tag/Total	Calibration Index
McN Str Wild	Estimate a.	Detection Efficiency	23.7%	23.7%	23.7%	58.0%	58.0%				
		Total Passage	37,350	10,991	129,819	18,722	101	196,983	196,983	201,313	1.0220
		American Passage	1,413	715	3,683	673	-	6,484	6,484	6,627	
		Naches Passage	7,470	2,234	29,463	5,657	-	44,824	44,824	45,809	
		American & Naches Passage	8,883	2,949	33,145	6,331	-	51,308	51,308	52,436	
		Upper Yakima Passage	28,467	8,377	98,943	14,269	77	150,133	150,133	153,433	
McN UnStr Wild	Estimate b.	Detection Efficiency	33.4%	33.4%	33.4%	33.4%	33.4%				
		Total Passage	26,445	7,782	91,916	32,450	174	158,767	158,767	168,133	1.0590
		American Passage	1,001	506	2,608	1,167	-	5,282	5,282	5,593	
		Naches Passage	5,289	1,582	20,860	9,805	-	37,536	37,536	39,750	
		American & Naches Passage	6,290	2,088	23,468	10,972	-	42,818	42,818	45,344	

		Upper Yakima Passage	20,155	5,931	70,055	24,732	133	121,007	121,007	128,145		
Pooled Str Wild	Estimate c.	Detection Efficiency	32.3%	20.1%	20.1%	35.9%	35.9%					
		Total Passage	27,409	8,065	92,297	18,321	98	146,190	146,190	151,265	1.0347	
		American Passage	1,037	525	2,618	659	-	4,839	4,839	5,007		
		Naches Passage	5,482	1,639	20,947	5,536	-	33,604	33,604	34,770		
		American & Naches Passage	6,519	2,164	23,565	6,195	-	38,443	38,443	39,777		
		Upper Yakima Passage	20,890	6,147	70,345	13,963	75	111,420	111,420	115,288		
Pooled UnStr Wild	Estimate e.	Detection Efficiency	44.0%	44.0%	44.0%	44.0%	44.0%					
		Total Passage	20,117	5,919	69,920	24,685	133	120,773	120,773	115,300	0.9547	
		American Passage	761	385	1,984	888	-	4,018	4,018	3,836		
		Naches Passage	4,023	1,203	15,868	7,459	-	28,553	28,553	27,259		
		American & Naches Passage	4,784	1,588	17,852	8,347	-	32,571	32,571	31,095		
		Upper Yakima Passage	15,332	4,512	53,290	18,814	101	92,049	92,049	87,877		
Hatchery		Prosser Hatchery Tally	8	1,419	64,446	82,305	789	148,967	Expanded Elastomer	Expanded PIT	PIT-Tag/Total	Calibration Index
McN-Str Hatch	Estimate a.	Total Passage	32	5,995	272,195	142,004	1,361	421,586	447,027	456,852	0.0569	1.0220
McN-UnStr Hatch	Estimate b.	Total Passage	24	4,245	192,723	246,127	2,358	445,452	472,332	500,195		0.7860
Pooled Str Hatch	Estimate c.	Total Passage	24	4,399	193,521	138,959	1,331	338,210	358,619	371,069		0.8170
Pooled UnStr Hatch	Estimate e.	Total Passage	17	3,229	146,602	187,226	1,794	375,875	398,556	380,494		0.8086

Appendix D

Survival to McNary Dam for PIT-tagged Spring Chinook Salmon smolts released at Roza Dam from 1999-2020



Prepared by:

Shubha Pandit, Bill Bosch and Mark Johnston
Yakima-Klickitat Fisheries Project
Yakama Nation Fisheries
P.O. Box 151, Toppenish,
WA 98948, USA

July 15, 2021

Table of Contents

EXECUTIVE SUMMARY	3
1.0 INTRODUCTION	6
2.0 METHODOLOGY	8
3.0 RESULTS AND DISCUSSION	14
3.1 FISH SIZES	14
3.2 YAKIMA RIVER FLOW BELOW PROSSER DAM	15
3.3. TRAVEL TIME FROM THE RELEASE SITE (ROZA DAM) TO McNARY DAM	16
3.4. SURVIVAL RATE OF HATCHERY- AND NATURAL-ORIGIN SMOLTS	19
3.5. EFFECT OF RIVER FLOW ON SURVIVAL RATE	22
3.6. COMPARISON OF NATURAL- AND HATCHERY-ORIGIN SMOLT SURVIVAL TO McNARY DAM OF ROZA RELEASES DURING THE "LATE" RELEASE PERIOD	24
3.7. COMPARISON OF EARLY AND LATE NATURAL-ORIGIN SMOLT SURVIVAL TO McNARY DAM	26
3.8. WEEKLY SURVIVAL RATE OF NATURAL- AND HATCHERY-ORIGIN SMOLT	26
4.0 ACKNOWLEDGMENT	33
5.0 REFERENCES	33

Executive Summary

This report summarizes the results of an evaluation to estimate survival rate and travel time of juvenile Spring Chinook Salmon (*Oncorhynchus tshawytscha*) released into the Roza Dam bypass during 2020. This evaluation is part of an ongoing study that was initiated in 1999. Differences between natural and hatchery rearing environments have a significant influence over the demographic attributes of naturally-spawned and hatchery-origin Chinook salmon beginning in early developmental stages. Moreover, hatchery-origin smolts released into the natural environment experience in-stream conditions that are dramatically different from a controlled hatchery rearing environment. Therefore, attempts to infer the survival rate for natural-origin smolts based on survival of hatchery-reared smolts (or vice versa) can be biased by relative differences in fish size, behaviors such as outmigration timing, fitness, and environmental conditions encountered during outmigration. Our investigation of interannual variation in survival rate and travel time for both natural- and hatchery-origin outmigrating smolts will help managers implement effective strategies for maintaining abundance and viability of the natural spring-run Chinook salmon population in the Upper Yakima River Basin.

In 2020, we tagged 2386 hatchery-origin smolts and 253 natural-origin smolts with passive integrated transponder (PIT) tags at Roza Dam. Tagged fish were released from March 13th through April 30th into the Roza Dam bypass system (ROZ - Release into the Roza Facility Bypass Flume/Pipe). The size of tagged and released hatchery-origin smolts ranged from 82 mm to 169 mm (average 121mm). Hatchery fish were significantly larger than PIT-tagged natural-origin fish, which ranged in size from 78 mm to 135 mm (average 105 mm for Natural-origin smolt).

Our results indicated variable travel times for hatchery- and natural-origin smolts, based on travel between the Roza Diversion Dam's bypass (about 206 kilometers upstream from the mouth of the Yakima River) and the downstream detection site at McNary Dam on the Columbia River 64 rkm downstream from the Yakima River. Fish generally exhibited immediate outmigration behavior after release. In 2020 the travel time from Roza Dam to McNary Dam for natural-origin smolts ranged from 6 to 50 days (mean \pm SE 21.23 \pm 1.18 days), whereas the travel time for hatchery-origin smolts ranged from 10 to 23 days (18.25 \pm 2.95 days). It indicates that

hatchery smolt took less time to travel to McNary Dam than the natural fish. Mean travel times in 2020 appeared to be shorter for smolts of hatchery origin compared to the 2018 and 2019 outmigration year but natural-origin smolt took about 3 days more on average than in 2019 to arrive at McNary Dam. Travel time was positively related with rate of river flow during outmigration, and results of analysis of variance (ANOVA) showed an interaction effect between fish size and origin (hatchery and natural) on travel time. Specifically, hatchery fish which were larger on average exhibited shorter travel times (days to reach McNary Dam) compared to travel times of smaller, natural-origin fish. These results are consistent with Melnychuk et al. (2010) who found that in small rivers, downstream travel speed increased with increasing body length. In addition, downstream outmigration timing and travel days are believed to be affected by many environmental variables, including photoperiod, river discharge, precipitation, lunar phase, air and water temperature, and fish size (Duston and Saunders 1995; McCormick et al. 1995; McCormick 2012; Sykes and Shrimpton 2009; Zydlewski et al. 2014).

In this study, the survival rates from release location to downstream detection at McNary Dam were estimated using the Cormack-Jolly-Seber (CJS) mark-recapture model, which has been commonly used within the Federal Columbia River Power System (FCRPS) to estimate survival rates for juvenile anadromous fish species (salmon and steelhead, see Tuomikoski et al. 2012). The model uses multiple detections of individual marked fish at several dams with PIT-tag detection capabilities. However, survival estimates in prior years (1999 to 2018) were based on a logistic regression model described in the 2018 annual report. We further evaluated the effects of river flow on survival rate by introducing flow as a covariate in the CJS model. Results indicated that survival rate between the Roza Dam release site and the McNary Dam in 2020 was about $38.3\% \pm 21.0\%$, but it was different between natural- and hatchery-origin smolts. The survival rate of the natural smolt was found to be $14.2\% \pm 11.8\%$ (mean \pm SE), where hatchery smolt had $50.2\% \pm 34.0\%$ survival rate. Very few smolts were also found to be detected at McNary Dam for both groups (natural origin-hatchery origin), which resulted in larger error bounds from the mean survival rate.

We further evaluated whether the survival rates of early (tagged and released on or before March 18th) and late (after March 18th) natural-origin and hatchery-origin Spring Chinook were different. The survival rate to McNary Dam of the early release group of natural-origin

(29.7%±2.4%) was lower than that of the late release group (33.8%±3.4%), but this difference was not significant. Within the late period, when both hatchery- and natural-origin smolts were migrating, the natural-origin survival rate to McNary Dam of (33.8%±3.4%), was significantly higher than the hatchery-origin survival rate (27.5±2.4%), suggesting that the superior adaptation of the natural-origin group to outmigration conditions outweighed any negative interactions with their hatchery-origin counterparts.

Results also revealed that survival rate increased with increasing river flow during the downstream migration time but the effect was not significantly different between hatchery- and natural-origin smolts. However, when using mean survival rate of all previous years (21 years, 1999-2020), the survival rate was relatively higher for the natural-origin fish ($F_{1,40} = 3.22$, $p = 0.084$). Further, we were unable to estimate juvenile survival rates by weekly basis for some of the groups because of high standard errors (SE) in some weeks or because the model failed to converge, indicating insufficient sample sizes for weekly estimates, especially for natural-origin juveniles.

1.0 Introduction

In recent years, naturally spawning Pacific salmon populations have declined relative to historical abundances, resulting in many ESA listings and heightened conservation concerns (Prince et al. 2019; Rand et al. 2012; Ford 2011; Gustafson et al. 2007). The recovery of depressed stocks is contingent on obtaining accurate and precise estimates of survival through the hydro system. Juvenile salmon emigrating from the Yakima Basin must navigate downstream through several dams in the Yakima and Columbia rivers during migration to the ocean. For over a decade, hatchery production in Yakima basin has been used to supplement natural salmon populations in order to benefit fisheries opportunities, and to boost declining natural populations. These hatchery programs are likely to continue and possibly increase significantly within the Columbia River Basin (WDFW 2019).

Since 1999, the Yakama Nation has been studying downstream survival of juvenile salmon, with a focus on understanding whether or not survival, and the factors affecting survival, are similar between hatchery-origin and natural-origin groups. The study involves annual releases of hatchery-origin and natural-origin spring Chinook salmon smolts with inserted passive integrated transponder (PIT) tags. There is some evidence to suggest that captive rearing of salmon under certain hatchery protocols (e.g. segregated programs) confers a genetic fitness deficit (domestication) to hatchery fish released into the natural environment compared to naturally reared salmon (Lynch and O' Hely 2001; Ford 2002; Frankham et al. 2002). While the CESRF program has attempted to minimize differences between hatchery- and natural-origin smolts, artificial rearing environments are still starkly different from the natural in-river conditions fish experience after release. Differences between natural and hatchery rearing environments have a significant influence over the demographic attributes of natural- and hatchery-origin Chinook salmon beginning early in their development. Inferring survival rates for one rearing history based on survival rates for smolts with a different history can be misleading due to differences in fish size, behavior, fitness, and acclimation to environmental conditions encountered during outmigration. In the CESRF program, the desired acclimation window as well as water permit limitations force an emigration window that is different than that of natural-origin fish (YN unpublished data), and this in turn will affect survival rates as flow conditions may vary considerably even on a daily basis depending on weather events. Hatchery-origin smolts are

often larger, likely due to temperature and feed regimens and nutrition content that differ from the temperatures and diet that natural-origin fish experience.

Survival rate may also vary with river flow and fish size during the outmigration period (Zabel and Achord, 2004). Juvenile outmigration is a critical phase in the overall life history of salmon (NPPC, 1992). Mortality rate is likely to increase as a function of migration distance (often hundreds of kilometers), where risk is compounded by exposure to several factors, including predation, extreme temperatures and diseases (Miller et al., 2014), and entrainment at diversions or dams. Furthermore, outmigration is concurrent with the smoltification process, where a fish undergoes physiological, behavioral and biochemical changes in preparation for saltwater habitat (Hoar 1976). Thus, it is important that the coordination between smoltification, outmigration, and arrival time to the estuary be preserved (Folmar and Dickhoff 1980) and remain on schedule with physiological readiness for saltwater.

The timing and duration of outmigration are believed to be affected by many factors, such as fish size, photoperiod, discharge, precipitation, lunar phase, water temperature and type of origin (natural vs. hatchery) (Duston and Saunders 1995; McCormick et al. 1995; McCormick 2012; Sykes and Shrimpton 2009; Zydlewski et al. 2014). In order to determine whether or not downstream survival rate and downstream migration dynamics of Spring Chinook smolts differ between natural and hatchery populations, our study focused on the following objectives:

- 1) evaluate the survival rate from the release location (Roza Dam) to McNary Dam (McN) between hatchery- and natural-origin smolts based on PIT-tag detections,
- 2) determine whether, for natural-origin Spring Chinook smolts, there is a significant difference in downstream survival between early outmigrants (sampled and PIT-tagged before the first hatchery-origin smolts appeared in the sample) and late outmigrants (captured during the hatchery smolt outmigration),
- 3) determine the effect of river flow on survival rate for both groups (hatchery- and natural-origin), and
- 4) determine whether or not travel time in the Yakima River differs between natural and hatchery smolts.

2.0 Methodology

The Roza Diversion Dam north of Yakima, Washington (Figure 1) withdraws water from the upper Yakima River for irrigation and hydroelectric power. Rotary drum screens and a bypass system return entrained fish to the Yakima River and provide an opportunity to sample and mark fish before they reenter the river. We queried the PTAGIS database (<https://www.ptagis.org/>) in February 2021 to retrieve available PIT-tag detection information for all Spring Chinook salmon smolts (hatchery- and natural-origin) released at Roza Dam from 2015 through 2020 (Roza bypass; Fig. 1). A total of 2386 hatchery-origin smolts and 253 natural-origin smolts with passive integrated transponder (PIT) tags were released from March 13 through April 30th, 2020 into the Roza bypass system (Fig. 2).

Hatchery-origin juveniles were acclimated at three sites upstream of Roza Dam: Jack Creek in the Teanaway River system, and Easton and Clark Flat on the Yakima River (Figure 1). Natural-origin and hatchery origin-smolts captured in the Roza bypass were PIT-tagged if not among the 40,000 (about 6% of the total hatchery release) tagged earlier at the hatchery. Previously-tagged hatchery Spring Chinook were noted as recaptures and included in the Roza release group.

Travel times and survival estimates of PIT-tagged hatchery-origin smolts were compared with those of PIT-tagged natural-origin smolts from the point of release into the Roza Dam bypass to the juvenile detection facilities at McNary Dam. Natural-origin smolts were identified as “early” for those sampled and PIT-tagged before the first hatchery-origin smolts appeared in the sample after their mid-March volitional release from acclimation sites. Natural-origin smolts captured during the hatchery smolt outmigration were assigned to the “late” group. In each release year, survival-estimate comparisons were made between late and early natural smolts, and travel time was measured as the difference between the release date at the Roza bypass and recovery/detection date at the downstream detection facilities at McNary Dam.

Although the survival rate from Roza Dam to McNary Dam in each year from 1999⁺ through 2018 was estimated using weighted logistic regression (see Neeley, 2018), the survival rates for both groups (natural- and hatchery-origin smolts) for the last 6 years (2015-2020) were estimated

¹ The first outmigration year of Upper Yakima River hatchery-reared Spring Chinook

using the Cormack-Jolly-Seber (CJS) mark-recapture model (see, White and Burnham 1999; Lebreton et al. 1992; Williams et al. 2002; Conner et al. 2015), in accordance with Federal Columbia River Power System (FCRPS) methodology (Tuomikoski 2012). The CJS model uses multiple detections of individual marked fish at several dams equipped with PIT-tag detection capabilities. The assumption of the CJS model is that there is no immigration or emigration during capture and recapture intervals, which is valid in the Columbia Basin hydrosystem (where smolts must pass several hydroelectric dams to reach the ocean) because fish behavior is relatively consistent (all fish are moving in one direction and over a relatively short period; (Conner et al. 2015). The CJS model was originally conceived to calculate time-interval survival of tagged animals by recapturing individuals and estimating survival and recapture probabilities using maximum likelihood. A spatial form of the CJS model can be used for species that migrate uni-directionally and are recaptured/detected within a discrete migratory corridor (Burnham 1987; Henderson et al. 2018). We used individual fish encounter histories to estimate the likelihood that a fish would survive and be detected at the tag detection facility of each hydroelectric dams (Lebreton et al. 1992).

The CJS model was run for different groups by year based on an encounter history constructed from the number of fish released at Roza dam and subsequent detection events at McNary and below McNary Dam (John Day and Bonneville dams). Similar to previous studies (Neeley 2018), all smolt releases were grouped into seven-day periods for analyses. For example, smolts released during ordinal days 1-7 and 8-14 were treated as two distinct release group based on Julian/ordinal period. The estimated survival rates were compared among release groups where the sample sizes were sufficient to provide statistical confidence. In general, every year the volitional exit period from the three Spring Chinook acclimation sites begins March 15, after the natural outmigration is well underway. Natural-origin Spring Chinook smolts captured, PIT-tagged and released at Roza Dam on March 18th or earlier (Julian date: <78) were categorized as early releases, and natural-origin captures after March 18 as late releases. This demarcation was based on Spring Chinook counts at the Prosser Dam juvenile monitoring facility downstream from Roza Dam during the 2020 outmigration year, where about 35% of the natural-origin smolts and none of the hatchery smolts passed Prosser dam on or before March 18th.

Analysis of variance (ANOVA) was performed to evaluate differential survival between hatchery- and natural-origin smolts, using group (hatchery-origin vs. natural-origin) and release period (early and late) as factors and years as replicates. There were no PIT-tagged smolts released for this study in 2014 because of a radio-tagging study conducted at Roza Dam. A radio-tagging study was also conducted in 2016, but PIT-tag releases continued on a reduced schedule that year.

Several environmental factors are known to influence downstream smolt survival, and river flow is among the most impactful (Raymond 1968; Connor et al. 2003; Tiffan et al. 2009). We introduced flow rate as a covariate in the CJS model to study its effects. Bureau of Reclamation (BOR) flow data were accessed at:

<https://www.usbr.gov/pn/hydromet/yakima/yakwebaread.html>. The average travel time from Roza Dam to McNary Dam averaged about 20 days (combining hatchery-origin and natural-origin smolts) so each day's release was assigned the 20-day average river flow commencing on the day of release. For example, a fish released on April 1st would reach McNary Dam by about April 20th, and the 20-day average flow rate for that time period would be assigned for that fish to determine the effect of river flow on survival.

Several CJS candidate models were built and compared using every possible combination of variables with river flow. Examples of candidate models included (1) two types of temporal variation in survival probability: time variation, which assumed that survival probability (ψ) varies by year; and no time variation which assumed that ψ remains constant for all years; (2) two types of temporal variation in detection probability: time variation, which assumed that detection probability varies by year; and no time variation which assumed that detection remains constant for all years; (3) variation in survival and detection probabilities between natural- and Hatchery-origin, and (4) influence of river flow on the survival and detection probabilities. Altogether, 49 models were built using these combinations (Table 4).

To determine the rank of the different models (49 models), we used the difference in QAICc score relative to the top model. For models with the difference of QAICc (QAICc) < 2 , we selected the model with the lowest QAICc and fewest parameters as the best model (Burnham

and Anderson 2002). We tested the Goodness of Fit (GOF) of competing models using the Bootstrapping Goodness of Fit Approach (“Bootstrap GOF”) in program MARK (Cooch and White 2012) to estimate the variance inflation factor for the model constructed to have the most parameters while remaining biologically meaningful (hereafter referred to as the “global model”). All subsequent models were then corrected for over-dispersion using \hat{c} . Using the best selected model, we estimated the effect of river flow on downstream survival rate (Roza Dam to McNary Dam) for both groups (hatchery- and natural-origin smolts). The CJS models and program MARK (White and Burnham 1999) were run within the RMark package (Laake and Rexstad 2019) in R statistical software, version 3.3.6 (R Core Team 2019).

Figure 1. The Yakima River Basin showing the three acclimation sites and Roza Dam where natural- and hatchery-origin Spring Chinook smolts were captured, tagged and released. Adapted from Fast et al. (2015).




Figure 2. Number of spring Chinook tagged and released into the Roza dam fish bypass (Hatchery-origin smolt, red; and natural-origin smolt, blue) for each year from 2015-2020. The value on the top of each bar represents the total number of tagged and released smolts on that specific date and year. Hatchery and natural-origin totals are also shown for each year with red for hatchery-origin and green for natural-origin (also see table 1 by groups of 7-days Julian date).




Table 1: Total release number of smolt with PitTags (hatchery-Origin and Natural-Origin) by seven-day period (Group by Julian period) for the 2015 -2020.

Group by Julian periods	# Hatchery-Origin smolt release						# Natural-Origin smolt release					
	2015	2016	2017	2018	2019	2020	2015	2016	2017	2018	2019	2020
42	0	0	0	0	0	0	19	0	0	0	0	0
49	0	0	0	0	0	0	40	0	0	0	0	0
56	0	0	0	0	0	0	107	0	0	0	0	0
63	0	0	0	0	0	0	0	0	0	0	0	0
70	264	0	0	0	0	0	143	0	0	0	0	0
77	0	0	0	0	0	0	29	97	0	110	0	73
84	265	177	431	714	0	388	31	32	48	159	0	16
91	86	167	286	547	0	825	26	3	58	47	0	35
98	417	198	294	305	508	469	84	4	31	30	19	72
105	70	100	0	579	739	136	24	0	0	5	52	37
112	0	160	252	359	373	308	0	0	21	29	6	18
119	211	171	275	74	388	191	6	0	12	2	211	2
126	119	84	245	0	128	69	15	0	11	0	0	0
133	0	47	0	0	102	0	0	0	0	0	4	0
140	0	0	0	0	0	0	0	0	0	0	0	0
147	0	0	0	0	0	0	0	0	0	0	0	0
Total	1432	1104	1783	2578	2238	2386	524	136	181	382	292	253

Note: All smolt releases were also grouped into seven-day periods: i.e., smolt released between Julian dates 1 and 7 were treated as one release period (Group by Julian periods:1), those released between Julian dates 42 and 48 were treated as another release group (Group by Julian periods: 42).

3.0 Results and Discussion

3.1 Fish sizes

During the last six years (2015-2020), fish with passive integrated transponder (PIT) tags were released for this study from as early as February to as late as May into the Roza bypass system (Figure 2).

In 2020, 2386 hatchery-origin smolts and 253 natural-origin smolts with PIT tags were released. The fork lengths of the released hatchery-origin smolts that were PIT-tagged at the dam ranged from 82 mm to 169 mm (average 121mm). Hatchery fish were significantly larger than PIT-tagged natural-origin fish ($F_{1,8}=16.87$, $p<0.01$), and ranged in fork length from 78mm to 135 mm (average 105 mm; figure 3, table 2).

Figure 3. Frequency distribution of fork lengths of hatchery and natural-origin smolts PIT-tagged and released at Roza Dam.




Table 2. Annual totals of smolts released by origin, their mean and median fork lengths (FL), and standard error (se) around each mean value.

Released Year	Hatchery-Origin				Natural-Origin			
	Total # fish (N)	Mean FL	Median FL	se	Total # fish (N)	Mean FL	Median FL	se
2015	1432	117.93	118.00	0.31	524	100.42	100.00	0.48
2016	1104	121.83	122.00	0.32	136	98.21	98.00	0.77
2017	1783	115.79	116.00	0.24	181	100.13	100.00	0.78
2018	2578	117.88	118.00	0.19	381	101.80	102.00	0.51
2019	2237	121.17	121.00	0.22	292	117.64	118.50	0.74
2020	2382	121.37	121.00	0.21	250	105.00	104.00	0.66


3.2 Yakima River flow below Prosser Dam

Yakima River flow below Prosser dam (gaging station YRPW, which represents the 18-kilometer reach between Prosser Dam and the Chandler Power Plant outfall) for monthly 2020 averaged approx. 1338 cubic feet per second (cfs), which was lower than the monthly flow from 2015 - 2018 but higher than April 2019 (Figure 4, Table 3). However average monthly during spring and summer month was lowest in 2015. For all years, the river flow from June to August was generally less than 800 cfs.

Table 3. Average monthly flow (cfs) and during spring and summer months of the Yakama River at the YRPW gage below Prosser Dam for the years 2015-2020.

	Months												Monthl y Average	Spring & summer (March-Aug) Average
	1	2	3	4	5	6	7	8	9	10	11	12		
2015	4793	4420	2523	1043	895	420	387	526	581	892	3549	5237	2106	966
2016	2632	8603	7982	8600	3437	983	822	519	517	2086	2582	1920	3390	3724
2017	1696	3460	9492	8778	6959	2697	640	666	657	1463	3585	2434	3544	4872
2018	3038	4138	2632	5183	6183	994	574	604	542	1054	1489	1775	2351	2695
2019	1389	1536	3066	4444	1860	563	560	749	568	1041	1458	2122	1613	1874
2020	3145	4411	1470	2050	2107	1241	577	584	693	1407	2195	2120	1833	1338

Figure 4. Average daily flow (cfs, blue line) and 20-day moving average flow (yellow line) of the Yakama River at the YRPW gage below Prosser Dam for calendar years 2015-2020. The red boxes represent the period in which natural- and hatchery-origin Spring Chinook smolts were tagged/released during that year.



3.3. Travel time from the release site (Roza Dam) to McNary Dam

The study showed that the travel time (days) of smolts from the release site (Roza Dam) to McNary (McN) dam during the 2020 migration year varied between hatchery- and natural-origin smolts (Figure 5). Most of the smolts were released during the month of April, which is a typical peak outmigration period. As a result, most of the fish generally exhibited immediate outmigration behavior after release. In 2020, one of the hatchery-origin smolts was detected at McNary Dam only 6 days from the date of release at Roza Dam. The travel time from Roza Dam

to McNary Dam for hatchery-origin smolts ranged from 6 to 50 days (mean±SE 18.13±0.9 days, see table 4); whereas the travel time for natural-origin smolts ranged from 9 to 37 days (21±1.19 days).

Since the hatchery-origin smolts were larger than natural-origin smolts (figure 6A), size might have played a role in the travel time difference between the hatchery and natural groups. The hatchery smolts took less time to reach McNary Dam than the natural-origin smolts. Previous results have showed that travel time varied among years and the variation might have been related to the variation in river flow among years. In general, the travel time between Roza and McNary dams decreased as river flow increased (Figure 6B).

Survival of juvenile salmon and travel days have been positively related to river discharge (Perry et al., 2018). Travel time was negatively related with river flow during outmigration, and analysis of variance (ANOVA) showed an interaction effect between fish size and origin (hatchery and natural) on travel time. Specifically, hatchery fish, which were larger on average, had shorter travel times to McNary Dam than the smaller, natural-origin fish. These results are found to be consistent with Melnychuk et al. (2010) who found that in small rivers, downstream travel speed increased with increasing body length. In addition, downstream outmigration timing and travel days are believed to be affected by many environmental variables, including photoperiod, river discharge, precipitation, lunar phase, air and water temperature, and fish size (Duston and Saunders 1995; McCormick et al. 1995; McCormick et al. 2000; McCormick 2013; Sykes and Shrimpton 2009; Zydlewski et al. 2013; Zydlewski et al. 2014).

Figure 5. Number of detections of PIT-tagged smolts released at Roza dam from 2015 through 2020 at McNary Dam by release date.





Table 3. The number of observations at McNary Dam (N), mean travel days from the release location to McNary Dam, its standard error (se), and maximum and minimum travel days (range) for both hatchery-origin and natural-origin Spring Chinook smolts by year from 2015-2020.

Release Year	Natural-Origin				Hatchery -Origin			
	Total # fish (N)	Mean	se	range	Total # fish (N)	Mean	se	range
2015	70	30.33	1.33	9-53	44	41.41	2.02	16-76
2016	132	13.42	0.83	3-45	12	41.00	3.06	24-62
2017	85	19.28	1.12	4-53	3	11.00	3.06	7-17
2018	108	24.63	0.88	5-43	17	36.71	2.90	20-60
2019	95	17.93	0.88	4-40	14	19.00	2.49	9-37
2020	43	21.23	1.18	6-50	4	18.25	2.95	10-23

Figure 6. The relationship between travel days from Roza dam to McNary (McN) dam and fork length at the time of tagging for hatchery- and natural-origin Spring Chinook smolts from 2015-2020 ([A.]); and the relationship between the average travel time (days) from release site (Roza Dam bypass) to McNary Dam and the average river flow below Prosser Dam during the months in which the fish were released ([B.])



3.4. Survival rate of hatchery- and natural-origin smolts

Based on the CJS model, the average survival probability from Roza Dam to McNary Dam for the pooled populations (hatchery- and natural-origin smolts combined) released at Roza Dam during 2020 was $38.30 \pm 21.0\%$ (mean \pm SE), however the hatchery-origin survival rate was $50.2 \pm 34.0\%$, which was higher than that of the natural-origin smolts ($14.2 \pm 11.8\%$; see Table 3). A similar result was observed in 2017 through 2019. The results further showed the standard error of the survival rate in 2020 was larger because fewer fish were detected at downstream dams in 2020 (Table 4). In fact, only 3 smolts were detected at all 3 dams with juvenile detection capability McNary, John Day and Bonneville.

We further evaluated whether the survival rates of early (tagged and released on or before March 18th) and late (after March 18th) natural-origin and hatchery-origin Spring Chinook were different. The survival rate to McNary Dam of the early release group of natural-origin ($29.7\% \pm 2.4\%$) was lower than that of the late release group ($33.8\% \pm 3.4\%$), but this difference was not significant (Table 7 and Figure 7B). Within the late period, when both hatchery- and natural-origin smolts were migrating, the natural-origin survival rate to McNary Dam of ($33.8\% \pm 3.4\%$),) was significantly higher than the hatchery-origin survival rate ($27.5 \pm 2.4\%$; Figure 7A and Table 7), suggesting that the superior adaptation of the natural-origin group to outmigration conditions outweighed any negative interactions with their hatchery-origin counterparts.


Table 4. Released and detected populations at McNary (McN) and Bonneville (BON) Dams and Roza-to-McNary survival rate for all (“All”: pooled hatchery- and natural-origin), hatchery-origin, and natural-origin smolts for the years 2015-2020. Note: there were 5 detection-history combinations and each group (Detection histories) is represented by the release-detection sequence and the number of fish detected in that sequence. “1-0-0” corresponds to release at Roza dam (1) but no detections at McNary (0) or Bonneville (0) dams. Similarly, “1-0-1” represents the number of fish released at Roza dam (1), not detected at McNary (0), but detected at Bonneville. The remaining two sequences for each year represent detection only at McNary, and detection at both McNary and Bonneville.

Year	Detection histories	Number of PitTags by Type		
		All (H+W)	Hatchery-origin (H)	Natural-origin (W)
2020	1-0-0	2529	2288	241
	1-0-1	66	58	8
	1-1-0	41	38	3
	1-1-1	3	2	1

	Survival rate	0.383±0.21	0.502±0.34	0.142±0.118
2019	1-0-0	2312	2044	268
	1-0-1	115	105	10
	1-1-0	87	74	12
	1-1-1	17	15	2
	Survival rate	0.316±0.066	0.318±0.070	0.288±0.174
2018	1-0-0	2799	2435	364
	1-0-1	41	39	2
	1-1-0	108	93	15
	1-1-1	12	11	1
	Survival rate	0.179±0.043	0.183±0.047	0.125±0.10
2017	1-0-0	1848	1674	174
	1-0-1	32	28	4
	1-1-0	79	76	3
	1-1-1	5	5	0
	Survival rate	0.316±0.128	0.299±0.12	***
2016	1-0-0	1070	946	124
	1-0-1	31	31	0
	1-1-0	125	113	12
	1-1-1	14	14	0
	Survival rate	0.360±0.077	0.370±0.079	***
2015	1-0-0	1807	1334	473
	1-0-1	37	28	9
	1-1-0	101	62	39
	1-1-1	11	8	3
	Survival rate	0.249±0.064	0.22±0.065	0.321±0.154

*** indicates the models failed to converge so that the survival rate could not be estimated.

Figure 7. The box plot showing the 22-year average survival probabilities of natural-origin (Natural) and hatchery-origin (Hatchery) Spring Chinook smolts (see Table 7 for the data). A. is the comparison of Late hatchery- and natural-origin smolts; and B. is the comparison between Early and Late natural-origin smolts.




3.5. Effect of river flow on survival rate

We further evaluated whether river flow affects the outmigration survival rate for hatchery- and natural-origin smolts. Among the 49 models shown in Table 6, the model that included an effect of river flow on the survival rate for the groups but varied by years had the lowest QAICs and therefore was selected to illustrate the effects of river flow on survival. Based on the best model, the survival rate between Roza and McNary dams was positively related with the river flows for the years 2015-2020 (Table 6 and Figure 8). This result is consistent with the previous results as that smolt migration and dam passage survival were positively correlated with stream flow because higher flows increase migration rates, potentially reducing exposure to predation, and reduce delays in reservoirs (see, Courter et al., 2016).

Table 5. The top 20 candidate models of the 49 candidate models, and associated statistical parameters. The models are ranked based on Quasi-likelihood Akaike's Information Criterion adjusted for over-dispersion (QAIC_c). The model with the lowest QAIC_c value was considered 'best'. "Wt" represents the weight of the model. S and p represent survival and capture probability, respectively, while "npar" represents the number of parameters used in the model. The models were built using 2015-2020 data.

SN	Models	npar	QAIC _c	Delta(Δ)	Wt
1	S(~Year + riverFlow) p(~Year:RearType + riverFlow)	20	6993.68	0.00	0.46
2	S(~Year:RearType + riverFlow) p(~Year + riverFlow)	24	6995.34	1.67	0.20
3	S(~Year:RearType + riverFlow) p(~RearType + riverFlow)	21	6996.75	3.07	0.10
4	S(~Year:RearType + riverFlow) p(~Year:RearType + riverFlow)	27	6997.03	3.35	0.09
5	S(~Year + riverFlow) p(~Year + riverFlow)	15	6998.96	5.28	0.03
6	S(~Year * RearType) p(~Year:RearType + riverFlow)	21	6999.58	5.90	0.02
7	S(~RearType + riverFlow) p(~Year * RearType)	21	7000.06	6.38	0.02
8	S(~Year * RearType) p(~Year + riverFlow)	21	7000.27	6.59	0.02
9	S(~1) p(~Year * RearType)	20	7000.91	7.23	0.01
10	S(~Year + riverFlow) p(~Year * RearType)	26	7001.16	7.48	0.01
11	S(~Year:RearType + riverFlow) p(~RearType)	20	7002.44	8.77	0.01
12	S(~ReleasedYear) p(~Year:RearType + riverFlow)	15	7002.58	8.90	0.01
13	S(~RearType) p(~Year * RearType)	21	7002.91	9.23	0.00
14	S(~Year:RearType + riverFlow) p(~1)	19	7003.56	9.88	0.00
15	S(~Year:RearType + riverFlow) p(~ReleasedYear)	23	7003.71	10.03	0.00
16	S(~ReleasedYear) p(~Year * RearType)	24	7003.84	10.16	0.00
17	S(~Year * RearType) p(~1)	18	7004.69	11.01	0.00
18	S(~Year * RearType) p(~ReleasedYear)	22	7005.63	11.95	0.00
19	S(~Year + riverFlow) p(~RearType + riverFlow)	15	7006.76	13.08	0.00
20	S(~Year * RearType) p(~RearType + riverFlow)	23	7007.70	14.02	0.00

Figure 8. The predicted survival rate as a function of river flow based on the best CJS models (“S(~Year + riverFlow) p(~Year:RearType + riverFlow)”; Table 5). The shaded area is the standard error of the predicted mean.



3.6. Comparison of Natural- and Hatchery-Origin Smolt Survival to McNary Dam of Roza releases during the “Late” release period

Yearly survival estimates based on all contemporaneous late-period smolt are given in Table 7 and Figure 9A (top panel). Because natural-origin smolts have spent more time in the natural habitat than hatchery-origin smolts by the time fish pass Roza Dam, it has always been hypothesized that, for smolt contemporaneously released at Roza, the survival to McNary of natural-origin smolt would be greater than that of hatchery-spawned smolt even though the hatchery-origin fish tend to be larger. However, in 2020, the survival rate of hatchery-origin smolts was greater than that of natural-origin smolts

(fig. 9A) and a similar result was observed in 2017 and 2018. However, when using mean survival rate of all previous years (21 years, 1999-2020), the survival rate was higher for the natural-origin fish ($F_{1,40} = 3.218$, $p = 0.08$) (Figure 7A).

Table 7. Survival of Spring Chinook smolts from Roza Dam to McNary Dam by origin, release year and release period, showing the number released (N), the survival probability, and the standard error of the survival probability (SE; 2020 only) for each release group.

Year	Natural-Origin						Hatchery-Origin					
	Early			Late			Early			Late		
	N	Surv.	SE	N	Surv.	SE	N	Surv.	SE	N	Surv.	SE
1999				312	0.739		1082	0.591		1082	0.591	
2000	3013	0.331		3196	0.498		2999	0.279		2999	0.279	
2001	755	0.475		1424	0.133		1744	0.175		1744	0.175	
2002	6130	0.216		2588	0.342		1503	0.263		1503	0.263	
2003	6614	0.314		1190	0.309		2146	0.246		2146	0.246	
2004	3699	0.354		232	0.375		1509	0.204		1509	0.204	
2005	1688	0.268		25	0.195		701	0.118		701	0.118	
2006	1833	0.197		500	0.513		3689	0.250		3689	0.250	
2007	1072	0.319		336	0.183		2477	0.406		2477	0.406	
2008	735	0.283		498	0.396		4911	0.260		4911	0.260	
2009	1804	0.430		239	0.484		3931	0.204		3931	0.204	
2010	0			105	0.540		1130	0.320		1130	0.320	
2011	1040	0.231		904	0.311		3051	0.331		3051	0.331	
2012	2482	0.301		191	0.241		4424	0.153		4424	0.153	
2013	2435	0.277		38	0.578		550	0.264		550	0.264	
2014												
2015	167	0.363		358	0.420		1503	0.243		1503	0.243	
2016	97	0.228		39	0.567		575	0.216		575	0.216	
2017				181	0.111		1869	0.216		1869	0.216	
2018	110	0.415		274	0.118		2550	0.214		2550	0.214	
2019	0			292	0.288	0.174				2238	0.318	0.07
2020	73	0.055	1.00*	180	0.155	0.127				2386	0.503	0.34
Mean		0.297	0.024		0.338	0.034		0.261	0.24		0.275	4

Note: estimates for the years 1999- 2018 are from Neeley 2019. * SE is very high indicating high uncertainty of the estimated value (low precision).

3.7. Comparison of Early and Late Natural-Origin Smolt Survival to McNary Dam

There were no early natural-origin fish releases at Roza prior to passage of hatchery-origin smolt in 1999, 2010, 2017, 2019, 2020; and, as stated earlier, there were no PIT-tagged releases at Roza Dam in 2014. Table 6 and Figure 9B present the natural-origin early and late smolt survivals from Roza to McNary for all years. Of the 18 years with early releases, late releases had greater Roza-to-McNary survival than early releases but the difference was not statistically significant ($F_{1,36}=1.524$, $p=0.22$, Fig 7B). In general, earlier outmigrants are believed to have a greater survival rate. However, the results showed that later releases had higher survival rates, although not significantly higher. A lower survival rate for earlier releases could be due to a lower proportion of out-migrants entering juvenile bypass systems where PIT tags can be detected. Generally, McNary Dam's bypass is watered up after Julian date 90 (March 30th), so fish passing earlier would be spilled rather than bypassed, resulting in a lower detection rate, consequently survival rate is also lower. It may also be that some of the early natural-origin releases pass McNary Dam before they could be detected in McNary's bypass, in which case the early-release natural survival estimates presented here may be underestimated.

3.8. Weekly survival rate of natural- and hatchery-origin Smolt

The survival rate (Roza-McNary Dam) varied by week for both groups (natural- and hatchery-origin), however the number of natural-origin releases were not sufficient to estimate the weekly survival rate with statistical confidence. In general, the hatchery-origin smolts that were released early [Julian date 91, which was the week of April 1st to 7th, 2020) had higher survival rate ($78.21\% \pm 7.4\%$) than the smolts released during the week [Julian date 126] between May 7 and May 12th, 2020 ($27.2 \pm 14.88\%$, see table 7 and figure 10).

Figure 9. Bar-diagram of Upper-Yakima Spring-Chinook Roza to-McNary Smolt-to-Smolt Survival for Late Natural- and Hatchery-Origin juvenile for each release year (1999-2020). **A.** is the comparison of Late hatchery- and Late Natural-origin smolt; and **B.** is the comparison between Early and Late Natural-origin Smolt.




Table 8. Roza-Dam to McNary-Detection Smolt-to-Smolt Survival probability with respect to Julian week. “Sur” and “N” represent survival probability and the number of smolts tagged and released, respectively.

Origin	Parameter	Julian Date																								Early	Late	Over All									
		345	351	359	365	7	14	21	28	35	42	49	56	63	70	77	84	91	98	105	112	119	126	133	140												
1999																																					
Natural	Sur																											0.47	0.88	0.64	0.85	0.78			0.74	0.74	
	N																												34	37	62	34	145			312	312
Hatchery	Sur																												0.53	0.70	0.65	0.60	0.55			0.59	0.59
	N																												266	103	306	100	307			1082	1082
2000																																					
Natural	Sur	0.44		0.20	0.40	0.34	0.34	0.28	0.31		0.49		0.52	0.54	0.42	0.69	0.52	0.65	0.55	0.56	0.17	0.33	0.40								0.33	0.50	0.42				
	N	56		47	55	1575	845	435	243		847		506	723	235	46	248	156	92	17	19	23	41								3013	3196	6209				
Hatchery	Sur								0.40		0.48		0.51	0.21	0.24	0.43	0.27	0.23	0.26	0.34	0.32	0.35	0.23								0.28	0.28					
	N								8		20		20	83	152	103	689	547	346	115	365	272	279							2999	2999						
2001																																					
Natural	Sur								0.40	0.48	0.39	0.40	0.50	0.64	0.60	0.29	0.33	0.15	0.09	0.09	0.05										0.47	0.13	0.25				
	N								32	121	159	145	144	85	69	85	150	155	583	396	55									755	1424	2179					
Hatchery	Sur															0.30	0.25	0.15	0.10	0.17	0.16										0.18	0.18					
	N															132	465	288	500	293	66									1744	1744						
2002																																					
Natural	Sur			0.17	0.14	0.29	0.20	0.18	0.16	0.32	0.16	0.25	0.28	0.23	0.32	0.32	0.36	0.34	0.33	0.41		0.35								0.22	0.34	0.25					
	N			500	501	295	761	960	533	178	388	328	804	398	484	617	665	277	750	47		232								6130	2588	8718					
Hatchery	Sur															0.51	0.35	0.21	0.24	0.20	0.14									0.26	0.26						
	N															89	428	144	444	108	290									1503	1503						
2003																																					
Natural	Sur								0.27	0.29	0.28	0.31	0.32	0.33	0.33	0.51	0.28		0.39	0.37	0.27	0.37	0.19						0.31	0.31	0.31						
	N								515	1188	1600	639	794	1284	256	338	441		284	110	85	115	155						6614	1190	7804						
Hatchery	Sur															0.34		0.28	0.25	0.25	0.12	0.13							0.25	0.25							
	N															431		574	221	411	332	177							2146	2146							
2004																																					
Natural	Sur		0.22	0.20	0.11			0.29	0.31	0.16	0.37	0.33	0.48	0.45	0.51	0.41		0.40			0.00								0.35	0.37	0.36						
	N		184	156	153			301	603	43	889	276	352	398	344	195		19			18								3699	232	3931						
Hatchery	Sur															0.28		0.22	0.12	0.11	0.09								0.18	0.18							
	N															220		1036	439	220	253								2168	2168							

		2005																						
Natural	Sur	0.23	0.25	0.32	0.21	0.41	0.63												0.19	0.27	0.19	0.27		
	N	831	300	335	110	77	35												25	1688	25	1713		
Hatchery	Sur									0.25	0.08	0.16							0.12		0.14	0.14		
	N									205	187	327							701		1420	1420		
		2006																						
Natural	Sur		0.15	0.17	0.13	0.21	0.29	0.14		0.58	0.47	0.32		0.62	0.33	0.35	0.65	0.39	0.34	0.27		0.20	0.51	0.26
	N		351	551	215	250	200	125		18	67	56		269	21	32	31	70	41	36		1833	500	2333
Hatchery	Sur													0.36	0.25	0.22	0.26	0.27	0.13	0.26		0.25	0.25	
	N													450	686	827	601	639	356	130		3689	3689	
		2007																						
Natural	Sur					0.27	0.37		0.32							0.06		0.53	0.42	0.52	0.32	0.18	0.29	
	N					453	476		143							233		31	62	10	1072	336	1408	
Hatchery	Sur															0.23		0.33	0.40	0.56		0.41	0.41	
	N															622		393	571	891		2477	2477	
		2008																						
Natural	Sur								0.22	0.33		0.41	0.46	0.28	0.64	0.41	0.11			0.40		0.28	0.40	0.33
	N								332	403		77	48	157	88	77	28			23		735	498	1233
Hatchery	Sur											0.27	0.24	0.21	0.18	0.27	0.16			0.40		0.26	0.26	
	N											505	467	879	316	505	1013			1226		4911	4911	
		2009																						
Natural	Sur						0.36	0.44	0.43	0.37	0.41	0.57	0.40		0.46		0.52	0.55		0.78	0.43	0.48	0.44	
	N						450	321	160	179	379	315	81		39		74	37		8	1804	239	2043	
Hatchery	Sur												0.34		0.25		0.18	0.27		0.13		0.20	0.20	
	N												413		712		920	448		1438		3931	3931	
		2010																						
Natural	Sur												0.70		0.45				0.51			0.54	0.54	
	N												33		57				15			105	105	
Hatchery	Sur													0.36		0.32			0.18			0.32	0.32	
	N												318		707				105			1130	1130	
		2011																						
Natural	Sur							0.18	0.23	0.52	0.31	0.26	0.13		0.41		0.83		0.64		0.23	0.31	0.27	
	N							430	538	72	113	473	126		109		58		25		1040	904	1944	
Hatchery	Sur										0.23	0.28	0.26	0.21	0.27		0.25		0.93			0.33	0.33	
	N										521	710	465	63	381		634		340			3114	3114	

		2012																
Natural	Sur	0.22	0.27	0.31	0.33	0.46	0.35	0.25		0.22	0.36	0.00		0.75	0.30	0.24	0.30	
	N	469	650	383	548	202	230	106		35	24	22		4	2482	191	2673	
Hatchery	Sur							0.16		0.18	0.14	0.03		0.64		0.15	0.15	
	N							839		1790	772	900		123		4424	4424	
		2013																
Natural	Sur	0.20	0.25	0.28	0.30	0.49					0.35	0.86		0.28	0.58	0.28		
	N	608	436	538	631	222					21	17		2435	38	2473		
Hatchery	Sur										0.18	0.31			0.26	0.26		
	N										182	368			550	550		
		2014																
Natural	Sur		0.17	0.47		0.42	0.74	0.58	0.24	0.21			0.24	0.36	0.42	0.40		
	N		60	107		143	60	26	84	24			21	167	358	525		
Hatchery	Sur					0.28	0.43	0.17	0.26	0.21			0.08		0.24	0.24		
	N					272	271	89	451	73			347		1503	1503		
		2016																
Natural	Sur					0.23			0.57				0.23	0.57	0.32			
	N					97			39				97	39	136			
Hatchery	Sur								0.22			0.70		0.24	0.24			
	N								575			35		610	610			
		2017																
Natural	Sur						0.00	0.00	0.00		0.59	0.00	0.70		0.11	0.11		
	N						48	58	31		21	12	11		181	181		
Hatchery	Sur						0.10	0.07	0.23		0.65	0.38	0.00		0.23	0.23		
	N						449	299	306		271	286	258		1869	1869		
		2018																
Natural	Sur					0.41	0.09	0.00	0.14			0.37		0.41	0.12	0.20		
	N					110	160	47	31			36		110	274	384		
Hatchery	Sur					0.05	0.18	0.15			0.40			0.21	0.21			
	N					753	576	317			904			2550	2550			
		2019																
Natural	Sur									0.1	NA	0.17	0.34		0.287	0.29		
	N														292	292		
Hatchery	Sur									0.34	0.2	0.71	0.61	NA	0.27		0.301	0.3
	N									508	739	373	388	128	102		2238	2238
		2020																
Natural	Sur					0.03	NA	NA	0.083	NA	NA	NA		0.06	0.14	0.14		
	N					73	16	35	72	37	18	2		73	180	253		
Hatchery	Sur					0.14	NA	0.31	NA	NA	NA	NA			0.5	0.5		
	N					388	825	469	136	308	191	69			2386	2386		

NA*indicates the model failed to converge so that the estimate was not reported.

Figure 10. Roza-dam to McNary-detection Smolt Survival Rate with respect to Julian Week grouping. Note: All smolt releases were also grouped into seven-day periods: i.e., smolt released between Julian dates 1 and 7 were treated as one release period (Group by Julian periods:1), those released between Julian dates 42 and 48 were treated as another release group (Group by Julian periods: 42).



Figure 10 (continued) Roza-dam to McNary-detection Smolt Survival Rate with respect to Julian Week grouping. Note: All smolt releases were also grouped into seven-day periods: i.e., smolt released between Julian dates 1 and 7 were treated as one release period (Group by Julian periods: 1), those released between Julian dates 42 and 48 were treated as another release group (Group by Julian periods: 42).



4.0 Acknowledgment

We thank all of the crews whose collective fish-tagging efforts over the years made this study possible. We are also grateful to Andrew Matala and David Lind who reviewed and provided valuable comments in the draft report. We would also like to thank Daylen Isaac who provided the hydrological data.

5.0 References

- Buckland ST, Burnham KP, Augustin NH. 1997. Model selection: An integral part of inference. *Biometrics*. 53:603–618.
- Conner, M. M., S. N. Bennett, W. C. Saunders, and N. Bouwes. 2015. Comparison of tributary survival estimates of steelhead using Cormack-Jolly-Seber and Barker models: Implications for sampling effort and designs. *Transactions of the American Fisheries Society*, 144:1, 34–47.
- Connor, WP, Steinhorst, RK, and Burge. HL. 2003. Migrational behavior and seaward movement of wild subyearling fall chinook salmon in the Snake River. *North American Journal of Fisheries Management* 23:414–430.
- Courter I., Garrison T and TJ Kock. 2015. Evaluation of stream flow effects on smolt survival in the Yakima River Basin, Washington, 2012-2014. <https://pubs.er.usgs.gov/publication/70176445>
- Duston, J., Saunders, RL, 1995. Increased winter temperature did not affect completion of smolting in Atlantic salmon. *Aquac. Int.* 3, 196–204.
- Fast, D. E., Bosch, W. J., Johnston, M. V., Strom, C. R., Knudsen, C. M., Fritts, A. L., & May, D. 2015. A synthesis of findings from an integrated hatchery program after three generations of spawning in the natural environment. *North American Journal of Aquaculture*, 77, 377–395.
- Folmar, CF., and Dickhoff, WW. 1980. The parr-smolt transformation (smoltification) and seawater adaptation in salmonids. *Aquaculture* 21: 1-37
- Ford MJ. 2002. Selection in captivity during supportive breeding may reduce fitness in the wild. *Conservation Biology*. 16:815–825.
- Ford, MJ. 2011. Status review update for Pacific salmon and steelhead listed under the Endangered Species Act: Pacific Northwest. U.S. Dept. Commerce. Available: http://www.nwr.noaa.gov/species/pacific_salmon_steelhead/
- Frankham R, Briscoe DA, Ballou JD. 2002. *Introduction to Conservation Genetics*. Cambridge

- Gustafson, R. G., Waples, R. S., Myers, J. M., Weitkamp, L. A., Bryant, G. J., Johnson, O. W., & Hard, J. J. 2007. Pacific salmon extinctions: Quantifying lost and remaining diversity. *Conservation Biology*, 21(4), 1009–1020.
- Gustafson, RG., Waples, RS., Myers, J. M., Weitkamp, LA., Bryant, GJ, Johnson, OW, and Hard, JJ. 2007. Pacific salmon extinctions: quantifying lost and remaining diversity. *Conservation Biology* 21: 1009– 1020.
- Henderson MJ, Iglesias IS, Michel CJ, Ammann AJ, Huff DD. 2018. Estimating spatial-temporal differences in Chinook salmon outmigration survival with habitat and predation related covariates. *Can J Fish Aquat Sci.* <https://doi.org/10.1139/cjfas-2018-0212>
- Hoar, W.S. 1976. Smolt transformation: evolution, behavior, and physiology. *J. Fish. Res. Board Can.* 33:1234-1252
- Jolly, GM. 1965. ‘Explicit Estimates From Capture-Recapture Data With Both Death and Immigration Stochastic Model.’, *Biometrika*, 52(1), pp. 225–47.
- Kock, T.J., Perry, R.W., and Hansen, A.C., 2016, Survival of juvenile Chinook salmon and coho salmon in the Roza
- Dam fish bypass and in downstream reaches of the Yakima River, Washington, (ver. 1.1, April 2017): U.S. Geological Survey Open-File Report 2016–1210, 32 p., <https://doi.org/10.3133/ofr20161210>.
- Laake, J. (2013) ‘RMark: an R interface for analysis of capture-recapture data with MARK’. Seattle WA: US Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Alaska Fisheries Science Center.
- Lebreton, JD, Burnham, KP, Clobert, J & Anderson, DR. 1992. Modeling survival and testing biological hypotheses using marked animals: a unified approach with case studies, *Ecological Monographs*, 62, pp. 67-118.
- Lynch M, O’ Hely M. 2001 Captive breeding and the genetic fitness of natural populations. *Conservation Genetics*.2:363–378.
- McCormick, MI. 2012. Lethal effects of habitat degradation on fishes through changing competitive advantage. *Proc. R. Soc. Lond. b* 279, 3899–3904.
- McCormick, SD., Björnsson, BTh., Sheridan, M, Eilertson, C., Carey, JB. & O’Dea, M. 1995. Increased daylength stimulates plasma growth hormone and gill Na⁺,K⁺-ATPase in Atlantic salmon (*Salmo salar*). *Journal of Comparative Physiology* 165, 245–254.

- Melnychuk MC., Welch DW., Walters CJ. 2010 Spatio-temporal migration patterns of Pacific salmon smolts in rivers and coastal marine waters. PLoS ONE. 5:e12916. doi: 10.1371
- Miller KM, Teffer A, Tucker S et al. 2014. Infectious disease, shifting climates, and opportunistic predators: cumulative factors potentially impacting wild salmon declines. *Evol Appl* 7:812–855
- Murray, C., Wood, C., 2002. Status of Sakinaw lake Sockeye salmon (*Oncorhynchus nerka*). Canadian Science Advisory Secretariat. Research document 2002-088.
- NPPC (Northwest Power Planning Council). 1992. Strategy for Salmon. Northwest Power Planning Council, Portland, Oregon.
- Perry, R.W., Pope, A.C., Romine, J.G., Brandes, P.L., Burau, J.R., Blake, A., Ammann, A., & Michel, C. (2018). Flow-mediated effects on travel time, routing, and survival of juvenile Chinook salmon in a spatially complex, tidally forced river delta. *Canadian Journal of Fisheries and Aquatic Sciences*, 75, 1886-1901.
- Price, MH., Connors, BM., Candy, JR., McIntosh, B., Beacham, TD., Moore, JW., and Reynolds, JD. 2019. Genetics of century-old fish scales reveal population patterns of decline. *Conservation Letters*.
- Rand, PS., BA. Berejikian, A. Bidlack, D. Bottom, J. Gardner, M. Kaeriyama, R. Lincoln, M. Nagata, TN. Pearsons, M. Schmidt, W. W. Smoker, L. A. Weitkamp, and L. A. Zhivotovsky. 2012. Eco-logical interactions between wild and hatchery salmonids and key recommendations for research and management actions in selected regions of the North Pacific. *Environmental Biology of Fishes* 94:343–358
- Raymond, HL. 1968. Migration rates of yearling chinook salmon in relation to flows and impoundments in the Columbia and Snake Rivers. *Transactions of the American Fisheries Society* 97(4): 356-359.
- Sykes, GE. and Shrimpton, JM. 2010. Effect of temperature and current manipulation on smolting in Chinook salmon (*Oncorhynchus tshawytscha*): the relationship between migratory behaviour and physiological development. *Canadian Journal of Fisheries and Aquatic Sciences* 67, 191–201.
- Tiffan KF, Kock TJ, Haskell CA, Connor WP and Steinhorst RK. 2009. Water velocity, Turbulence, and Migration Rate of Subyearling Fall Chinook Salmon in the Free-Flowing and Impounded Snake River. *Trans Am Fish Soc* 138:373–384.
- Tuomikoski, J., J. McCann, B. Chockley, H. Schaller, S. Haeseker, J. Fryer, R. Lessard, C. Petrosky, E. Tinus, T. Dalton, and R. Ehlke. 2012. Comparative survival study (CSS) of PIT-tagged spring/summer Chinook and Summer Steelhead 2012 Annual report. (<http://www.psmfc.org/wp-content/uploads/2013/08/ComparativeSurvivalStudyofPIT-TaggedChinook.pdf>)

- WDFW. 2019. Proposal to increase hatchery production to benefit southern resident killer whales. Washington Department of Fish and Wildlife, Olympia, Washington, USA.
- White, GC., and KP. Burnham. Program MARK: survival estimation from populations of marked animals. *Bird Study* 46(Supplement): 120—138.
- Zabel RW, Achord S. 2004. Relating size of juveniles to survival within and among populations of Chinook salmon. *Ecology* 85:795–806
- Zydlevski GB., Stich DS. & McCormick SD. 2014. Photoperiod control of downstream movement of Atlantic salmon *Salmo salar* smolts. *Journal of Fish Biology* 85: 1023–1041.

Appendix E

Juvenile Coho outmigration survival and adult Coho returns to the Yakima Basin, 1999-2020



Prepared by:
Shubha Pandit, Todd Newsome, Bill Bosch
Yakima-Klickitat Fisheries Project
Yakama Nation Fisheries
P.O. Box 151, Toppenish, WA 98948, USA

July 16, 2021

EXECUTIVE SUMMARY 1

1. INTRODUCTION 6

2. METHODOLOGY 8

 2.1 GEOGRAPHICAL DISTRIBUTION: HISTORICAL AND CURRENT8

 2.2 FISH PIT-TAG DATA9

 2.3 DATA ANALYSES11

 2.4 THE SMOLT-TO-ADULT RETURNS (SAR)13

 2.5 AGE COMPOSITION OF ADULT RETURNS14

3. RESULTS AND DISCUSSION15

 3.1 FISH SIZE (FORK LENGTH) AT THE TIME OF TAGGING AND RELEASE15

 3.2 TRAVEL TIME FROM RELEASE LOCATIONS TO McNARY DAM.....16

 3.3 DETECTION RATE OF SMOLT RELEASES AT McNARY DAM.....18

 3.4 SURVIVAL PROBABILITY (RELEASE SITE TO McNARY DAM)20

A. Survival probability of smolt and parr releases by migration year.....20

B. Survival probability of smolt releases by broodstock22

C. Survival probability of smolt and parr releases by release location23

 C.1 Smolt releases23

 C.1.1. Annual comparison of survival rates for Prosser releases.....25

 C.1.2. Annual comparison of survival rates for Stiles releases25

 C.2 Parr releases27

 C.2.1. Annual comparison of survival rates for parr releases in Yakima Basin streams29

 C.2.2. Effect of release month on parr survival rate.....31

 C.3 Effect of river flow and release month on survival rate33

 3.5 THE SMOLT-TO-ADULT RETURNS (McNARY JUVENILE TO BONNEVILLE ADULT).....36

 3.6 AGE- DISTRIBUTION AT RETURN37

4. ACKNOWLEDGMENT.....39

5. REFERENCES39

Executive Summary

Coho salmon (*Oncorhynchus kisutch*) in the Yakima basin were extirpated in the early 1980s but reintroduction efforts initiated in the mid-1980s have resulted in hatchery-produced coho naturally reproducing in both the Yakima and Naches rivers. In 1984 there was no escapement (n=0) of adult Coho Salmon returning to the Yakima Basin, but the return of hatchery-produced origin fish peaked in 2014 (> 25,000 adults). Several release strategies for outplanting fish have been implemented in the reintroduction program to evaluate and compare relative survival and escapement. Outplants have been released at both the parr and smolt life stages, including different size classes, released at multiple locations, including different release dates, and outplanted from different broodstock sources. A diverse strategic approach has been utilized to maximize the likelihood of achieving stable and abundant returns of natural-origin Coho salmon to the Yakima River and to enhance the stability and resiliency of the population against potential environmental changes.

An ongoing, long-term monitoring program is being conducted with the aim of improving project objectives and strategies by applying what is learned from the project experiments, monitoring and evaluation, and literature reviews in an adaptive management framework. This evaluation is an annual update of ongoing monitoring that began with the first reintroduction efforts in 1996. The report summarizes the estimated survival rate and travel time of juvenile (smolt and parr) Coho salmon releases from multiple locations in the Yakima basin with a focus on the following objectives:

- Determining survival rate and travel time of smolts released in 2020 and parr released in 2019 (migration year 2020) and quantifying annual trends
- Comparing survival rates between outplants from different broodstock sources (Yakima-origin vs. outplants from Eagle Creek National Fish Hatchery and Washhoulgal Hatchery)
- Identifying watershed-specific survival rates between the upper Yakima River and Naches River release locations for out-migrating juveniles, and identifying whether survival rate differs as a function of release month (February, March or April)
- Evaluating the effects of Yakima River flow (using average flow at Prosser Dam for the summer months) on outmigration survival rate
- Evaluating Smolt-Adult return (SAR) percentage for Coho released as parr and as smolts each year for outmigration years 2004-2020

- Determining the age composition of the coho returns from ocean at Bonneville Dam for Coho released as parr and those released as smolts

Sample size and fish size

In 2020, a total 13845 PIT-tagged smolts were released in the Yakima Basin. Among the PIT-tagged smolts, the brood source for 9954 tagged smolts was Eagle Creek NFH with 2952 tagged smolts originating from Yakima Basin returns. Both stocks were reared in Prosser Hatchery and released in the Yakima River at Prosser Dam on March 27, 2020. Another 939 tagged Yakima-origin smolts were reared at Prosser Hatchery and released in the lower Yakima Basin at Ahtanum Creek on Feb 18, 2020. Regarding parr for migration year 2020, 1289 tagged parr were released into the Naches River near the Tieton River confluence in 2019 (8/9/2019). During 2020, 1249 tagged parr were released in the upper mainstem Yakima River near the Holmes acclimation site near Ellensburg WA in August of 2020, but this parr release group is not included in this report because they start to outmigrate only in 2021. Although a total 13845 smolts and 2538 parr were PIT-tagged for this (2020) migration year, fish length information was available only 79 fish and all of which were released at Prosser Dam. Based on the available data, the length of the smolts released at Prosser was 105.35 ± 0.89 mm at the time of tagging.

Travel time to McNary Dam

Most of the 2020 smolts were released at Prosser Dam. The average mean travel time from Prosser Dam to McNary Dam was about 32.96 ± 8.93 days; whereas in 2019 it was 21.32 ± 8.54 days. The variation in travel time among years can be associated with several in-river conditions including variation in river flow, water temperature or release time. Besides the Prosser Dam release, relatively few smolts (939 smolts with PIT tags) were released from Ahtanum Creek, and none of the released fish were detected at McNary Dam so that it was not possible to compare the travel times among release locations for the 2020 migration year. For the parr release group, the 2019 Upper Yakima River parr outmigrating in 2020 were detected at McNary Dam $288 \text{ days} \pm 12.5 \text{ days}$ (mean \pm SE; range 276 days to 301 days) after release. In previous years (2015-2019), fish released at Prosser Dam exhibited the shortest travel time to McNary Dam, whereas the Jack Creek release group from the upper Yakima Basin had the longest travel time (mean 47.14 ± 4.59 days). Travel times for the

groups released at the Easton, Holmes and Stiles ponds, Wenas Lake, and Ahtanum Creek ranged from 33 to 39 days.

McNary Dam detection rate

The overall smolt detection rate at McNary Dam was $4.31\% \pm 0.58\%$ in migration year 2020, which was slightly lower than the detection rate of $6.23\% \pm 0.9\%$ in 2019. In general McNary detection rates varied among years. The highest detection rate was in 2016 ($24.63 \pm 1.51\%$) and the lowest was in 2020 ($4.31\% \pm 0.58\%$). Variation in the detection rate at McNary Dam might be due to river flow, spill percentage and how surface-passage structures were operated. Similar to smolt releases, detection rates at McNary Dam for parr releases were also variable among years.

Release-McNary Juvenile survival rate

The average survival probability of juvenile Coho Salmon smolts from the release sites to McNary Dam in 2020 was $47.14 \pm 5.78\%$, which was higher than the 2019 estimate ($14.27 \pm 2.64\%$), the 2018 estimate ($24.51 \pm 3.2\%$) and the 2017 estimate ($29.06 \pm 3.4\%$). The higher survival rate for 2020 migration is at least partly due to release location because most of the tagged smolts in 2020 were released at Prosser Dam in the lower Yakima River. In other years, smolts were also released from several locations upstream from Prosser Dam, with correspondingly lower survival rates.

The survival rate of smolts to McNary Dam was higher for the Eagle Creek-stock releases ($52.6 \pm 7.4\%$) than for Yakima-origin release ($41.1 \pm 10.1\%$). In 2019 and 2020, there was no release of the Washougal stock, but survival rate for the data pooled from 2015 to 2020 by stocks the highest survival rate was for Eagle Creek smolts and the lowest was for Washougal smolts, with Yakima-origin smolts in the middle of the survival range.

We further evaluated whether there was an effect of hatchery environment on survival rate. Yakima-origin smolts were released in two groups from Stiles Pond on the same date in 2016. One group had been reared at Eagle Creek Hatchery and the other at Prosser Hatchery. Survival rate (Stiles to McNary Dam) was higher for the smolts reared at Eagle Creek Hatchery (35.26 ± 4.14 , mean \pm SE) than for those reared at Prosser hatchery (13.28 ± 1.49 , mean \pm SE), which suggests a hatchery effect, which could arise from many factors including water temperature and water quality. Fish sizes would have aided the comparison but were not available.

Since smolts were released over a three-month period (February, March and April), release date might also have affected survival because of the variation of the habitat quality or quantity including water temperature and river flow. The effects of river flow and release month were introduced as covariates in the CJS model using survival data from 2015 – 2020 releases. Smolts released in March had a higher survival rate compared to February and April, and higher flow during the 20-day period following release improved survival in all release months.

For parr released during the 2020 migration year, releases occurred only in the lower Yakima River at Ahtanum Creek in 2019 but the very low number of fish detected at McNary Dam indicated that the survival rate for parr migrating in 2020 was also very low. For the last 6 migration years (2015-2020), Coho parr were released at different locations from May to October. Release site-to-McNary survival of the parr releases was higher for the population released in August ($14\% \pm 0.20$) and followed by the group of July releases ($3.1\% \pm 0.40$) and then June releases ($1\% \pm 0.4$).

The Smolt-to-Adult Return (SAR) percentage

The percentage of smolts that survive and return (SAR) to spawn is important because it incorporates cumulative impacts of the freshwater habitat, hydrosystem and ocean conditions that determine the sustainability of adult returns over time. In this study, SARs were based on the percentage of smolts detected at McNary Dam that returned as adults to Bonneville Dam. Since Coho can spend as many as 3 years in the ocean, we estimated SAR for the populations that out-migrated from 2004 through 2018 for both parr and smolt releases.

The results showed that the (McNary-to-Bonneville) SAR estimates varied by year and life stage at release during the 15-year study period. On average, the SAR was slightly higher for the group released as parr (SAR: $3.98 \pm 1.06\%$) than the SAR for the group released as smolts (SAR: $3.79 \pm 1.06\%$). The highest SAR for the group released as parr was in migration year 2018 ($8.75\% \pm 1.66\%$), followed by the groups released in 2013 ($7.56 \pm 1.23\%$) and 2008 ($6.85 \pm 1.04\%$), but lowest in migration year 2006 (0%). For smolts, the highest SAR was for the group released in 2008 ($8.28 \pm 0.25\%$), followed by 2013 ($8.08 \pm 0.91\%$) and 2010 ($6.48 \pm 1.14\%$), while the lowest SAR was in 2011 ($0.85 \pm 0.25\%$). The variation in SARs among years can be associated with many factors such as smolt size, release and ocean entry timing, and ocean conditions.

Returning adult age composition

For outmigration years 2004 through 2019, a total of 3194 returning PIT-tagged Coho released as smolts and 1459 returning Coho released as parr in the Yakima Basin were detected at Bonneville Dam. For the adult returning group released as smolts, ~85% of were age 3 (ocean age 1), 9% were age 2 (ocean age 0), and 7% were age 4 (ocean age 2). For the group released as parr, 90% of returning Coho were age 3, 10% were age 2, and none were age 4.

1. Introduction

Prior to their extirpation in the early 1980's, Yakima Basin Coho salmon (*Oncorhynchus kisutch*) were once widely distributed among tributaries of the Yakima and Naches rivers (Fulton 1970; Chapman 1986), with annual adult returns numbering from 44,000 to 150,000 (Kreeger and McNeil 1993).

Releases of hatchery reared Coho salmon in the Yakima Basin began in 1983 with the first release of 324,000 smolts originating from the Little White Salmon Hatchery (YN 1997). In 1988, the Yakama Nation (YN) and the Washington Department of Fish and Wildlife (WDFW) developed and implemented a reintroduction program that has successfully shown evidence of natural production in both the Yakima and Naches rivers. The highest return of adults (2014) from hatchery releases and natural production was greater than 25,000 fish.

Several alternative release strategies have been utilized in the reintroduction program over time in response to observations in long-term monitoring. Smolts were initially released in the mainstem of the Yakima River (Dunnigan et al. 2002), but subsequent releases have explored a range of different locations to understand how geographically and hydrologically diverse habitats within the Yakima Basin affect outmigration survival and adult returns. Habitat capacity and quality have a significant impact on growth rate and survival, and within the Yakima River Basin human alterations to the environment continue to exacerbate naturally limiting conditions by reducing the quality and quantity of available spawning and rearing habitat. On the other hand, broad habitat restoration programs are concurrently being implemented to improve habitat conditions in many Yakima Basin streams. Other exploratory release strategies have included variable life stages (parr vs. smolts) at release, different release times, and use of multiple outplant sources. In past years, the primary sources of Coho outplants have been Yakima Basin returns, Eagle Creek National Fish Hatchery and WDFW's Washhoughal Hatchery. In total, about 500,000 juvenile coho have been released each year from permanent acclimation sites or from temporary mobile acclimation facilities operated in upstream locations in tributary streams of the Naches and upper Yakima rivers.

Columbia River Coho typically spend one year in freshwater before out-migrating as yearling smolts (typically in April and May), then spend two growing seasons (about 18 months) in the ocean before returning as 3-year-old adults (Hassler 1987) to spawn in their natal streams (Beamish et al. 2004). Precocious, sexually mature males (jacks) may also return to spawn after a summer at ocean. Adult Coho generally migrate upstream at water temperature ranging from 7.2°C to 15.6°C (Reiser and

Bjornn 1979 cited in Laufle et al. 1986) and spawn from late October to November, sometimes as late as December or January.

Spawning normally occurs in transitions from pools or runs to riffles, in minimum water depth of 0.18 m, at water temperatures ranging from 4.4°C to 9.4°C, and velocities ranging from 0.3 to 0.91 m/sec (Thompson 1972, BOR 2007). The optimum temperature for coho salmon egg incubation was 4°C to 11°C (Davidson and Hutchinson 1938, cited in Sandercock 1991). Juvenile coho salmon survive best in low-gradient habitats (generally less than four percent; Jones and Moore 1999) and tributaries with a stream gradient less than 3% with complex and deep pools or beaver ponds (Bradford et al. 1997 and Reeves et al. 1989).

An ongoing, long-term monitoring program is being conducted with the aim of monitoring progress towards project objectives and improving strategies by applying what is learned from the project experiments, monitoring and evaluation, and literature reviews in the Yakima-Klickitat Fisheries Project adaptive management policy. This report is an annual update of an ongoing monitoring effort that began in 2001. It summarizes survival rate and travel time estimates for juvenile Coho parr and smolts released from multiple locations in the Yakima basin, with a focus on the following objectives:


- ❖ Determining survival rate and travel time of smolts released in 2020 and parr released in 2019 (migration year 2020)
- ❖ Comparing survival rates between outplants from different broodstock sources: Yakima returns vs. outplants either from Eagle Creek National Fish Hatchery or Washougal Hatchery
- ❖ Identifying watershed-specific survival rates among upper Yakima basin and Naches basin locations for out-migrating juveniles, and identifying whether survival differs as a function of release month (February, March, April)
- ❖ Evaluating the effects of river flow on outmigration survival rate
- ❖ Determining the annual Smolt-Adult return (SAR) from 2004-2020 and age compositions of the adult returns

2. Methodology


2.1 Geographical distribution: historical and current

Coho salmon were native to the Yakima River basin and its spawning area was quite widespread in the Yakima River basin, including the Bumping River (Wydoski and Whitney 2003; Tuck 1995). Historically, it was assumed that Coho were present in low-gradient streams in the Yakima Basin prior to extensive habitat alteration and were widely distributed among tributaries of the Yakima and Naches rivers (Haring 2001; Berg and Fast 2001; Figure 1A). Acclimation and release sites designated in the reintroduction program overlap this historical geographical distribution (Figure 1B).

Figure 1. Historical Coho geographical distribution and recent reintroductions.



B. *Coho smolt and parr release sites, 2008-2020*



2.2 Fish PIT-tag Data

We queried the PTAGIS database (<https://www.ptagis.org/>) in May 2021 to retrieve available PIT-tag detection information for all Coho Salmon smolts released at the different locations in the Yakima Basin from 2015 to 2020 (Figure 1). Numbers of PIT-tagged fish released each year among sites in the Yakima Basin ranged from 13,865 in 2020 to 20,305 in 2019 (Figure 1, Table 1).

Two stocks (Yakima and Eagle Creek) were released from Prosser and Ahtanum creek on Feb 18 and March 27, 2020 for outmigration year 2020. Fish were released at only two sites because of restrictions associated with the COVID pandemic. A total of 2952 and 939 PIT-tagged smolts of the

Yakima stock were released in 2020 at Prosser Dam and Ahtanum Creek on the La Salle High School grounds, respectively. A total of 9874 PIT-tagged smolts from Eagle Creek Hatchery were released at Prosser Dam in 2020 (Table 1).

Table 1: Broodstock sources, juvenile rearing facilities, release sites and counts of PIT-tagged smolts released in outmigration years 2015 to 2020.

Broodstock	Rearing Facility	Release site	Release Year					
			2015	2016	2017	2018	2019	2020
1. EagleCr	Eagle Cr. Hatchery	Easton Pond	3751			4994	5088	
		Holmes Pond	2501				2495	
		Stiles Pond	2498			5008	5011	
		Prosser Hatchery						9974
2. Washougal	Prosser Hatchery	Prosser hatchery				4254		
3. Yakima (In-basin)	a. Eagle Cr. Hatchery	Easton Pond			5098			
		Holmes Pond			5050	5002		
		Stiles Pond			1253	5007		
	b. Prosser Hatchery	Wenas Cr. above Wenas Lake					814	
		Wenas Cr. below Wenas Lake					819	
		Wenas Lake at Upper Boat Launch					567	
		Ahtanum	6	869	1527		1705	939
		Prosser Hatchery	1265	2501	2876	2509	2533	2952
		Lost Creek Pond	2506	2502				
		Stiles Pond	2520	2503				
		Buckskin slough				1250		
		*South Fork Cowiche Cr.				1251		
		Mobile Acclimaiton below Roza dam		2500				
		Mobile Acclimaiton into Buckin Slgh	1247	2501				
		Mobile Acclimaiton into Rattlesnake Cr.	1249					
		Mobile Acclimaiton into Cowiche Cr.	1250					
Jack creek Acclimation site					1273			
		Total	18793	24777	14412	19266	20305	13865

Unlike smolts, which begin emigration immediately after release, parr typically outmigrate as yearling smolts in the spring following their release, so Coho parr released in 2019 were evaluated for migration year 2020. A total of 1289 PIT-tagged Coho parr were released into the Tieton River of the Naches Subbasin in 2019 (migration year 2020, Table 2). The total release and the number of release sites were much smaller and fewer than in previous years as shown in Table 2. There were no parr releases for migration year 2017.

Table 2: Coho parr releases by migration year from 2015-2020. Parr were released one year earlier than the designated migration year.

Sub-basin	Release Location	Migration Year				
		2015	2016	2018	2019	2020
Lower Yakima	AHTANC - Ahtanum Creek, Yakima River	1349	1648	3009	4453	
Naches	COWICC - Cowiche Creek- Naches Subbasin	3017	3005	3035	3013	
	INOUYE.SIDE.CHANNEL	1606	0	0	0	
	Little Naches	6036	3008	3042	3006	
	Little Naches_SF	3004	0	0	0	
	NATCHR - Natches River	0	3017	0	3549	
	Quartz Cr.	3012	0	0	0	
	Rattlesnake Cr	0	3032	0	3049	
	TIETNR - Tieton River	0	0	0	3010	1289
Upper Yakima	HundleyPonds_nearNelsonSiding	1531	0	0	0	
	Big cr	3003	3013	0	3056	
	Lake.Cle.Elum	0	3015	0	0	
	Mercer Cr	0	1543	0	0	
	Mercer Cr. Upstream	0	1523	0	0	
	Reecer Creek	3026	0	3069	3005	
	SWAUKC - Swauk Creek	0	0	3024	3041	
	Wilson Cr	3027	3011	3019	6082	
	YAKIM2 - Yakima River - above Naches River	0	0	3046	6011	
Total		28611	25815	21244	41275	1289

2.3 Data analyses

Travel times and survival rates for both parr and smolt releases from the different release locations to McNary Dam were estimated each year from 2015 to 2020. Travel time was estimated as the difference between the date of release and the date of detection at McNary Dam.

For outmigration years 2007 through 2018 a logistic regression model (Neeley 2012) was used to estimate survival probability of the groups. Beginning in 2019 and in this report, survival probability from release locations to McNary Dam and detection rate of the released PIT-tagged Coho smolts at McNary Dam were estimated using the Cormack-Jolly-Seber (CJS) mark-recapture model (see, White and Burnham 1999; Lebreton et al. 1992; Williams, et al. 2002, Conner et al. 2015), which has been commonly used within the Federal Columbia River Power System (FCRPS) to estimate survival

rates for juvenile anadromous fish species (Tuomikoski et al. 2013). The model uses multiple detections of individually marked fish at several dams with PIT-tag detection capabilities.

Among the several assumptions of the CJS model, one assumption is no immigration or emigration during capture (tagging) and recapture (detection) intervals, which is valid in the hydrosystem because of necessary passage at several hydroelectric dams, and where fish behavior is relatively consistent as fish are moving in one direction over a relatively short period of time (see; Conner et al. 2015). The CJS model was originally formulated to calculate time-interval survival of tagged animals by recapturing individuals and estimating their survival and recapture probabilities using maximum likelihood. A spatial form of the CJS model can be used for species that migrate unidirectionally, and are recaptured/detected within a discrete migratory corridor (Henderson et al. 2018, Burnham 1987). We used individual fish encounter histories to estimate the likelihood that a fish would survive and be detected at each tag receiver facility (dams in this study; see Lebreton et al. 1992). The CJS model was run for all smolts released at each location based on an encounter history constructed from the number of fish released at the different locations and subsequent detection events at McNary, John Day and Bonneville dams on the Columbia River. Similar to previous studies (Neeley 2018), we estimated the survival rate and detection efficiencies for each release group and broodstock source.

Several environmental factors are known to influence downstream smolt survival, and river flow is among the most impactful (Raymond 1968; Connor et al. 2003; Tiffan et al. 2009). Since early and late release groups presumably experience variable flow regimes in the Yakima River, each is likely to incur a different rate of survival associated with temporal river conditions. Therefore, it was necessary to introduce both river flow and release month as covariates in the CJS model to estimate the survival rate of the releases. In the model we used the last six years of data (2015-2020) to increase the overall sample size and confidence around our estimates. Fish were released from February through May with multiple releases in each year (2015-2020); however, in 2015, a drought year, a negligible number (6) of PIT-tagged Coho were released in May, and we excluded this release and evaluated release month effects for February through April in 2015.

Flow data for the Yakima River below Prosser Dam (YRPW) were accessed from the Bureau of Reclamation website at: <https://www.usbr.gov/pn/hydromet/yakima/yakwebarcread.html>. The average travel time from Prosser to McNary Dam was approximately 20 days. Accordingly, a 20-day

moving average of river flow data was assigned to each tag PIT-tagged fish to determine the effect of river flow on survival rate of the release group.

Several candidate CJS models were built using every possible combination of river flow and release month, with varying or constant survival and detection probabilities at dams in the CJS models. To determine the rank of the different candidate models we used the difference in the QAICc (Δ QAICc: Quasi-likelihood AICc Akaike's information criterion difference) relative to the top model. For models with Δ QAICc <2, we selected the model with the lowest QAIC and fewest parameters as the best model (Burnham and Anderson 2002). Selecting the best model, we estimated the effect of river flow on downstream survival rate for each release group. The CJS models were run within the RMark package (Laake and Rexstad 2019) in R statistical software, version 3.3.6 (R Core Team 2019). More information about the model is available in Pandit et al. 2020 (Appendix D of Bosch et al., 2020).

2.4 The Smolt-to-Adult Returns (SAR)

SAR, which is the percentage of smolts that survive and return as an adult to spawn, is a metric that captures most of the cumulative impacts of the hydro-system and ocean conditions on anadromous fish, indicating how sustainable the returns of adults are over time. The SAR was estimated as the percentage of smolts detected at McNary Dam returning as adults to Bonneville Dam using the following equation for each year and release group:

$$U_{at\ McN\ \&\ BON} / J_{at\ McN}$$

Where, $U_{at\ McN\ \&\ BON}$ is a total number of PIT tagged fish which were detected at McNary Dam (McN) as a juvenile and also detected at Bonneville Dam (BON) as a returning adult (joint detection). $J_{at\ McN}$ is the total number of fish detected at McNary Dam as juveniles. Since Coho can spend as many as 3 years in the ocean, we estimated SAR for the populations that out-migrated from 2004 through 2018 for both groups (released as parr and smolt). Nonparametric 90% confidence intervals were computed around the estimated annual overall SARs for each group as described by McCann et al. (2020). The nonparametric bootstrapping approach of Efron and Tibshirani (1993) was used where first, the point estimates were calculated from the sample for each population, and then the data were re-sampled, with replacement, to create 1,000 simulated samples (Berggren et al. 2002, Chapter 4). These 1,000 iterations are used to produce a distribution of annual SARs from

which the value in the 50th ranking is the lower limit and value in the 950th ranking is the upper limit of the resulting 95% nonparametric confidence interval.

2.5 Age composition of adult returns

The ocean age of each returning Coho was estimated by subtracting the date of detection at the Bonneville Adult passage from the date of release. Coho smolt and parr releases naturally show different outmigration behavior after release. Coho smolts start to migrate downstream immediately after release, while parr typically outmigrate as yearling smolts in the spring following release in summer/Fall. Therefore, for parr release groups, ocean age was estimated as:

1. Ocean age of parr = date of detection of returning adult at Bonneville Dam - release date - 365 days;

whereas the ocean age of the smolt was estimated as:

2. Ocean age of smolt = date of detection of returning adult at Bonneville Dam - release date

Return age composition was estimated as the proportion of each age class of adult return detected at Bonneville Dam for each brood year and life stage.

3. Results and Discussion

3.1 Fish size (Fork length) at the time of tagging and release

Among the 111,418 Coho smolts released with PIT-tags from outmigration years 2015 through 2020, lengths at the time of tagging were available for only 8605 fish (7%; Table 3). The broodstock sources of these smolts were Yakima returns, Eagle Creek outplants and Washougal outplants released in March and April of each outmigration year. Overall, there was no significant difference in mean smolt fork length among release groups in different months, but fish released in March tended to be larger at tagging than fish in the April release groups. This was contrary to expectations since fish should be growing larger over time. This was most likely a hatchery effect, as March releases were largely comprised of fish reared at the Prosser hatchery where water temperatures are higher than at the other hatcheries used to rear Coho juveniles for this study.

For the 2020 migration year only, a total 13845 smolts (9954+2952+939) and 1289 parr were released but fish length information was available only for 79 of the PIT-tagged Coho smolts released at Prosser. The mean length of the smolts in this small sample was 105.35 ± 0.89 mm at the time of tagging. Since the fish of the Ahtanum releases had not been measured, the length at tagging could not be evaluated.

Table 3: Smolt fork length by year, release location and release month, with sample size (n). Data are based on the limited data available from PITAGIS (n= 8605 out of 111,418 total tags).

Year	Release Location	Release Month	n	Mean (mm)	SE	Range	
						min	max
2015	Easton	March	431	133.76	0.47	94	166
2015	Holmes	March	377	126.15	0.48	95	157
2015	Stiles	March	585	119.78	0.60	72	168
2016	Easton	April	521	114.49	0.44	63	155
2016	Holmes	April	1074	112.82	0.29	63	144
2016	Stiles	April	558	122.07	0.54	82	160
2016	Prosser	April	303	133.06	0.46	104	155
2016	Ahtanum	March	520	127.28	0.62	75	220
2016	LostCr	April	85	129.96	0.79	110	150
2017	Holmes	March	292	115.83	0.48	85	136
2017	Stiles	April	600	116.08	0.35	88	140
2017	Prosser	March	414	126.72	0.52	91	160
2018	Easton	April	1108	108.56	0.23	83	140
2018	Stiles	April	800	107.40	0.25	83	151

2019	Easton	April	206	100.20	0.62	71	118
2019	Holmes	April	204	101.31	0.75	67	126
2019	Stiles	April	442	100.22	0.52	67	126
2020	Prosser	March	79	105.35	0.89	80	123

3.2 Travel Time from Release Locations to McNary Dam

Most of the 2020 migration-year Coho smolts were released from Prosser Dam, and only 939 smolts (7%) were released from Ahtanum Creek. For the group released at Prosser, its mean travel time to McNary Dam was 32 days \pm 8.93 days (see table 4A), which was about 11 days more than 2019 (it was only 21.32 \pm 8.54 days in 2019 migration year). Variation in travel time among years can be associated with several in-river conditions including variation in river flow, water temperature, release timing, or fish size. Other in-river conditions can also affect movements, such as hydroelectric dams and their impoundments on the Columbia River, especially for smaller or not fully-smolted juveniles. The 2019 Prosser group was released on April 02, 2019 and the 2020 Prosser group was released March 27, 2020, which was about 5 days later during 2020 out-migration year than 2019 outmigration year.

None of the 939 smolts released from Ahtanum Creek were detected at McNary Dam, so it was not possible to compare travel times among release locations for the 2020 outmigration year. However, based on the 2015-2019 outmigration years for the evaluation of release sites, fish released at Prosser Dam exhibited the shortest travel time to McNary Dam (mean 21.32 \pm 8.54 days), whereas the group from Jack Creek, released 234 river kilometers upstream from Prosser Dam, had the longest travel time (mean 47.14 \pm 4.59 days). Travel times for the groups released at the Easton, Holmes and Stiles ponds, Wenas Lake, and Ahtanum Creek, all upstream from Prosser Dam, ranged from 33 to 39 days.

Coho parr released in 2019 and outmigrating in 2020 were detected at McNary Dam after 288 days \pm 12.5 days (mean \pm SE) following release in 2019, ranging from a minimum of 276 days to a maximum of 301 days (Table 4B).

Table 4. Travel time from release site to McNary Dam for [A] smolt releases, and [B] parr releases.

A. Average travel days from release location to McNary Dam for smolt

Broodstock	Rearing Facility	Release site	Release Year/Migration Year						
			2015	2016	2017	2018	2019	2020	
1. EagleCr	Eagle Cr. Hatchery	Easton Pond	56			45	38		
		Holmes Pond	54				33		
		Stiles Pond	52			17	51		
		Prosser Hatchery						33	
2. Washougal	Prosser Hatchery	Prosser hatchery				52			
3. Yakima (In-basin)	a. Eagle Cr. Hatchery	Easton Pond		35					
		Holmes Pond		31	65				
		Stiles Pond		34	31				
	b. Prosser Hatchery	Wenas Cr. above Wenas Lake						NA	
		Wenas Cr. below Wenas Lake						NA	
		Wenas Lake at Upper Boat Launch						34	
		Ahtanum	NA	33	31			36	NA
		Prosser Hatchery	19	19	34	48	21	34	
		Lost Creek Pond	57	42					
		Stiles Pond	52	36					
		Buckskin slough				53			
		*South Fork Cowiche Cr.				74			
		Mobile Acclimaiton below Roza dam		33					
		Mobile Acclimaiton into Buckin Slgh	27	32					
Mobile Acclimaiton into Rattlesnake Cr.	46								
Mobile Acclimaiton into Cowiche Cr.	54								
Jack creek Acclimation site						47			
		Average Travel Days	46	33	40	48	37	34	

B. Average travel days from release location to McNary Dam for parr

Sub-basin	Release Location	Migration Year				
		2015	2016	2018	2019	2020
Lower Yakima	AHTANC - Ahtanum Creek, Yakima River	NA	357	292	209	
Naches	COWICC - Cowiche Creek- Naches Subbasin	NA	298	294	316	
	INOUE.SIDE.CHANNEL	325				
	Little Naches	NA	290	294	304	
	Little Naches_SF	688				
	NATCHR - Natches River		285		308	
	Quartz Cr.	664				
	Rattlesnake Cr		291		300	
	TIETNR - Tieton River				297	288
Upper Yakima	HundleyPonds_nearNelsonSiding	NA				
	Big cr	320	290		316	
	Lake.Cle.Elum		528			
	Mercer Cr		290			
	Mercer Cr. Upstream		298			
	Reecer Creek	329		289	311	
	SWAUKC - Swauk Creek			301	312	
	Wilson Cr	326	288	288	309	
	YAKIM2 - Yakima River - above Naches River			290	302	
Total		442	322	293	299	288

3.3 Detection rate of smolt releases at McNary Dam

For 2020 outmigration year, a total of 13845 Coho were released as smolts from Prosser Dam in 2020 and 1289 were released as parr in the Naches River in 2019. Only a small proportion of smolts passing McNary Dam were detected at McNary Dam. The overall detection rate of the smolts at McNary Dam was $4.31\% \pm 0.58\%$, which indicates that with 95% confidence the true detection rate of can be found between 3.17% and 5.45% (Table 5a). A very few of the parr released in 2019 were detected at McNary Dam during migration year 2020, and the detection rate was $25.00\% \pm 21.65\%$, in which standard error was very high ($\pm 21.65\%$). It indicates that the uncertainty of the estimated detection rate is large, which might be due to a combination of factors such as lower sample size and higher mortality during the extended period between release and outmigration (Table 5b).

When evaluating detection rate at McNary Dam from 2016 through 2020, it was found that the detection rate varied by year. The highest rate was in 2016 and the lowest detection rates were in 2016 and 2020 (see Table 5). The variation in the detection rate at McNary Dam can be due to how

surface-passage structures are operated. In recent years, increasing spill and the use of surface-passage structures (spillway weirs) at dams are a primary management strategy to increase survival of juvenile fish passing dams within the Federal Columbia River Power System. Greater use of spillways results in a lower proportion of fish entering juvenile bypass systems where PIT tags can be detected (Widener et al. 2018), and fluctuations in spill and flow can produce variable detection rates among years or within a migration season.

Table 5: Detection history (number of juvenile Coho detected/not detected at McNary and Bonneville dams) and detection rate during out-migration of smolt release groups (A) and parr release groups (B) over migration years 2015-2020. Enumeration of fish fate (Release/detection histories) is coded by detection (1) and no detection (0) such that “1.0.0.” = no juvenile detection after release, “1.0.1” = not detected at McNary Dam but detected at Bonneville Dam, “1.1.0” = detected at McNary Dam but not at Bonneville Dam, and “1.1.1” = detected at both dams.

A. Smolt releases

Detection History	2015	2016	2017	2018	2019	2020
No detection after release (1.0.0)	18167	23128	13601	18356	19775	12430
Detected at BON Dam but not at McNary Dam (1.0.1)	392	621	337	483	338	1153
Detected at McNary Dam but not at BON Dam (1.1.0)	179	825	431	379	168	230
Detected at all Dams (1.1.1)	55	203	43	48	24	52
Detection rate (%)	12.30	24.63	11.31	9.03	6.23	4.31
Standard Error (\pm SE)	1.51	1.51	1.62	1.24	1.31	0.58

B. Parr releases (released parr typically outmigrate as yearling smolts). The year is the migration year. For example, number of fish in 2015 is the number of parr released in 2014.

Detection History	2015	2016	2017	2018	2019	2020
No detection after release (1.0.0)	28547	25473		20614	41175	1283
Detected at BON Dam but not at McNary Dam (1.0.1)	19	41		333	30	4
Detected at McNary Dam but not at BON Dam (1.1.0)	41	283		260	69	4
Detected at all Dams (1.1.1)	4	18		37	1	1
Detection Rate	0.90	3.82		13.98	5.26	25.0
Standard error (\pm SE)	0.39	0.74		2.05	5.13	21.65

Note: there was no parr release in 2016 (migration year 2017)

3.4 Survival Probability (Release Site to McNary Dam)

A. Survival probability of smolt and parr releases by migration year

The average survival probability of juvenile Coho smolts from tagging to McNary Dam in 2020 was $47.14 \pm 5.78\%$, which was higher than the averages in the previous 5 years (Figure 2). Obtaining the higher survival rate for 2020 migration can be due to an effect of release location because more than 90% of the smolts in 2020 were released from Prosser Dam compared to all other years: when smolts were released from several other locations in the Yakima Basin, all upstream from Prosser Dam. The previous studies showed that the survival rates were lower for the groups released from other locations compared to the group released at Prosser Dam. Outmigration survival rates might have been also influenced by in-river conditions such as water temperature and river flow in addition to outmigration distance as reported by Scheuerell et al. (2009), Petrosky and Schaller (2010) and Haeseker et al. (2012). For example, in 2015 there was an extremely low snow pack and an early snowmelt, which would have affected flow rate and water temperature. Not only for Coho, Summer chinook had also higher survival rate in 2020 compared to previous years. So other important environmental variables not included in the study may have a positive effect on the survival rate during the 2020 migration year.

The higher survival rate in the 2020 migration year can be also associated with either early fish release timing or favorable river flow or a combination of both. In 2020, the majority of the fish were released in the early period (end of the March), whereas in 2019 they were released in the middle period (April-May, Figure 7, Table 11). Since these fish were released early they were exposed to more pulse flow events (reservoir releases specifically for fish outmigration, Figure 9), compared to number of such events in 2019, which might have facilitated their migration down the Yakima River. Previous results in the Yakima Basin have shown that juvenile downstream survival increases as river flow increases, similar to the results cited above. Although releases from Prosser are advantageous from the standpoint of juvenile survival, adult harvest augmentation and assuring sufficient number of local broodstock for future juvenile releases, upstream releases are more likely to result in adult homing to viable spawning and rearing habitat and the ultimate development of self-sustaining natural coho populations in the Yakima Basin.


For the 2019 parr release migrating in 2020, very few PIT-tagged fish were detected downstream from McNary Dam, and consequently the survival rate was very low. In previous years, despite being

lower, survival rates of parr releases varied from year to year in a similar manner to those of smolt releases (Figure 2).

Because of their known effects on survival from other studies, river flow and release month were introduced as covariates in the CJS model using PIT-tag data from 2015 – 2020 releases as mentioned in methodology. The results also showed that effect of river flows on outmigration survival rate depend on the release months (February, March and April). Coho smolts released in March had a higher survival rate compared to February and April releases.

The only lower Yakima Basin parr releases were in Ahtanum Creek in 2019 but survival rate for Ahtanum Creek was very low. Pooling all other parr releases over the migration years 2015-2020, comparisons could be made among the parr release months of May through October. Release site-to-McNary survival of the parr-releases was highest for the population released in August ($14\% \pm 0.020$), followed by the groups of released in July ($3.1\% \pm 0.40$) and June ($1\% \pm 0.4$).


Figure 2. Overall smolt survival rate (\pm SE) from release site to McNary Dam for smolt and parr releases in migration years 2015-2020. The asterisk for 2020 indicates that the survival rate of parr could not be estimated because of no detection at dams below McNary Dam.



B. Survival probability of smolt releases by broodstock

During the 2020 migration year, the survival rate was higher for the Eagle Creek-stock releases ($52.6 \pm 7.4\%$) than for the Yakima Basin-stock release ($41.1 \pm 10.1\%$). In 2019 and 2020, there was no release of the Washougal stock. However, when we estimated the survival rate by pooling the data for 6 years from 2015 to 2020 by stocks (Yakima-stock releases and out-of-basin Eagle Creek-stock and Washougal-stock), we found that the average survival rate (2015-2020) for smolt releases differed among the stocks (Figure 3). The highest survival rate was for Eagle Creek smolts and the lowest was for Washougal smolts.

Figure 3. Average Coho smolt survival rate (release to McNary Dam) and 95% confidence intervals by broodstock origin for the migration years 2015 through 2020.



We had an expectation that survival rate of the smolts of the in-basin broodstock (Yakima broodstock) should have higher survival rate compared to Eagle Creek outplants, but that was not the case. We further evaluated whether there was an effect of hatchery environment on the survival rate. We were able to compare hatchery environments with two Stiles Pond releases in 2016 that varied only with respect to hatchery environment. Both were Yakima stock but one was reared at Prosser Hatchery and the other at Eagle Creek Hatchery: tag file session notes read: “*Yakima coho smolts reared at Prosser and released from Stiles pond*” and “*Yakima coho smolts reared at eagle cr NFH and*

released from Stiles pond'. The survival rate from Stiles Pond to McNary Dam was higher for the smolts reared in Eagle Creek Hatchery ($35.26 \pm 4.14\%$, mean \pm SE) than for those reared in Prosser hatchery ($13.28 \pm 1.49\%$, mean \pm SE). This variation in the survival rate indicates that hatchery environment seemed to have an effect. Variation in survival rate among hatcheries can be associated with many factors including water temperature and water quality of the hatchery. Previous studies outside of the Yakima basin showed that the size of the fish also affected the juvenile survival rate but we did not measure whether fish sizes of the smolts reared in two different hatcheries were different.

C. Survival probability of smolt and parr releases by release location

C.1 Smolt releases

In each year, smolts released at the Prosser site had the highest survival among all Yakima River sites (Table 6). Annual survival rates during 2015-2020 for all sites ranged from a low of $0.88 \pm 0.6\%$ in 2019 (Ahtanum Creek) to a high of 97.8% in 2018 for the Prosser release (Table 6, Figure 4). The high survival estimates for 2018 may be due to low estimated detection efficiencies for that release group, and this estimate was also based on the different method (see Neeley 2018) than the CJS model. For 2019 and 2020, we employed CJS models, which have been commonly used within the Federal Columbia River Power System (FCRPS) to estimate survival rates for juvenile anadromous fish species (Tuomikoski et al. 2013). In 2020 the Prosser release group had a 41.1% survival rate to McNary Dam. The survival rate of the smolts released into Ahtanum Creek could not be estimated because none of the Ahtanum tags were detected at McNary Dam (Table 6, Figure 4).


Table 6. Survival probability from the release location to McNary Dam for Coho smolt releases from 2015 through 2020. For 2019 and 2020 results, standard errors are also given (mean \pm SE). "NA" means that survival rate could not be estimated because there were not enough detections at downstream dams.

Stock	Release site	2015	2016	2017	2018	2019	2020
Yakima	• Stiles	8.2	24.7	27.40			
	• Prosser	37.2	22.9	66.50	97.9	25.2 \pm 2.9	41.1 \pm 10.1
	• Easton		13.3				
	• Buckskin Slough		20.4		24.1		
	• South Fk Cowiche Creek				25.3		
	• Ahtanum Creek					0.88 \pm 0.6	NA
	• Jack Creek					6.01 \pm 3.5	
	• Wenas Lake above Wenas Dam (Wenas wildlife area)					0.25 \pm 1.1	

• Wenas Creek below Wenas Dam		NA	
• Wenas Lake at Upper Boat Launch		1.6±1.1	
• Stiles	25.5	16.83±6.8	
• Prosser			52.6±7.4
Eagle Creek	• Easton	9.2	17.2±8.0
	• Holmes		6.5±4.0
Washougal	• Prosser		32.10

Note: Estimates for the years 2015-2018 were adopted from Neeley (2018).

Figure 4. Survival probability (release site to McNary Dam) of Coho released as smolts in outmigration years 2019 and 2020.



C.1.1. Annual comparison of survival rates for Prosser releases

As shown above, the juvenile outmigration survival rate to McNary Dam varied by release locations. Prosser releases had the highest survival rates among the groups released from the different locations, but it also varied among years. The highest estimated survival rate for a Prosser release was 97.9% in 2018 (Table 7), but as discussed above, the estimate is likely to be inaccurate, either because of a low detection rate at downstream dams or methodological errors. Ignoring 2018, the highest survival rate was in 2014 (78%) and the lowest was in 2016 (22.9%, Table 7).

Table 7. Survival to McNary Dam for Yakima-origin Coho released at the Prosser site. Standard errors are available only for the 2019 and 2020 releases.

Year	Number released	Release Date	Travel days (Mean \pm SE)	Survival Probability (Mean \pm SE)
2007	2499	4/15	15	62.7
2008				
2009	2506	4/2	41	65.7
2010	1371	4/4	24	52.5
2011	5036	4/15	30	37.6
2012	3811	3/5	58	33.9
2013	2520	4/15	8	67.2
2014	3004	4/14	18	78.0
2015	1265	3/23	21	37.2
2016	2501	4/4	19	22.9
2017	2876	3/19	34	66.5
2018	2509	3/14	48	97.9
2019	2533	4/2	21.32 \pm 8.54	25.19 \pm 2.98
2020	2952	3/27	33.78 \pm 1.14	41.06 \pm 10.09

Note: Estimates for the years prior to 2019 were adopted from Neeley (2018)

C.1.2. Annual comparison of survival rates for Stiles releases

Similar to Prosser, the survival rate to McNary dam of Stiles releases also varied by year. There were no Stiles releases in 2018, 2019 or 2020 (Table 8).

Table 8. Survival to McNary Dam for Yakima-origin released from Stiles Pond.


Year	Number released	Release Date	Travel days (Mean \pm SE)	Survival Probability (Mean \pm SE)
2001	1240	5/17	22	43.2
2002				

2003	1249	5/7	14	40.0
2004				
2005				
2006	2490	4/3	38	32.7
2007	2449	4/5	41	25.0
2008				
2009	2515	4/15	36	47.6
2010	2501	4/12	36	18.7
2011				
2012	2526	4/16	32	38.0
2013	2504	4/15	30	44.2
2014	2505	4/16	25	44.9
2015	2520	3/23	51	08.2
2016	3768	4/7	35	24.7
2017	5007	4/17	31	27.4
2018	NO RELEASE			
2019	NO RELEASE			
2020	NO RELEASE			

Note: Results were adopted from Neeley (2018)

Although the survival rates of Prosser and Stiles releases both varied by year, the Prosser release groups had higher survival rates to McNary Dam in most years than the Stiles groups. Only in 2012 and 2016 did Stiles releases survive better than Prosser releases (Figure 5).

Figure 5. Bar plot showing survival to McNary Dam for the Yakima-origin Coho released at Prosser Dam from 2007 through 2020 (red color) and from Stiles Pond (green color) from 2001 through 2020. The 2019 and 2020 results included 95% confidence intervals.



C.2 Parr releases

For the migration year 2020, the survival rate for the Coho group released as parr in 2019 from the release site to McNary Dam was very low. Fewer tags were released in 2019 than the average for prior parr releases (3000 tags is a typical release, but 2019 release was under 1300 tags), and small sample sizes may have combined with poor survival. Previous years' results have shown marked fluctuations in survival of parr releases (Table 9), and the survival rates for parr releases in the Yakima basin have been generally lower than the survival rates from smolt releases. For migration year 2019, the survival rate of parr releases was ~5% (Figure 6), with the highest survival rate observed among releases from the Rattlesnake Creek and the lowest measurable survival rate for the Big Creek and South Fork Cowiche Creek groups (less than 1%; Table 9). Survival rate from Swauk Creek to McNary Dam was also low (0.13%) but its standard error was very high (75.53%), because


only a few fish were detected at downstream dams (Table 9, Figure 6). Releasing more PIT tags per release site would help to reduce the error rates.

Table 9. Survival probability (from the release location to McNary Dam) for Coho parr releases in 2018 and 2019 (outmigration years 2019 and 2020). “NA” or “*” represent releases with too few downstream detections to estimate survival rate or to reduce estimation error.

Release Location	2019		2020	
	Mean (%)	SE (%)	Mean (%)	SE (%)
Ahtanum Creek	4.71	1.06		
Rattlesnake Creek	15.25	5.07		
Big Creek	0.4	0.15		
Naches River	4.78	4.42		
Easton Reach	NA			
SF Cowiche Creek	0.4	0.28		
Reecer Creek	2.56	1.1		
Swauk Creek	0.13	75.53*		
Tieton River	9.16	8.6	0.93	0.71
Coleman Creek	4.79	2.92		
Little Naches	NA			
Wilson Creek	2.14	0.87		
Yakima River (Thorp Boat Ramp)	NA			
All (Pooled)	5.26	5.31		

* There was an issue in model convergence because of low or no detections at downstream dams

Figure 6. Survival probability from release site to McNary Dam of the group released as parr in migration years 2019 and 2020 (release year 2018 and 2019). “NA” indicates no estimate of the survival rate due to lack of model convergence (not enough detections at downstream dams).



C.2.1. Annual comparison of survival rates for parr releases in Yakima Basin streams

Table 10 summarizes annual variations in survival rates of Coho parr released from seven locations in the Yakima Basin. There was substantial variation among years within sites, and among sites within years.

Table 10. Estimated survival from release to McNary Dam of Coho released as parr, by release location and migration year. For 2019 and 2020 results, average survival rate and its standard errors are also given (mean \pm SE) where applicable. An asterisk indicates that the survival rate could not be computed because of too few downstream detections.

Released river/tributary	Year	Released Pop ⁿ (N)	Survival rate (%)	SE	Stock	Notes
Cowiche Creek	2008	3001	30.7		Yakima	
	2009	6			Wild Parr	
	2009	3001	23.3		Yakima	
	2010	3004	16.9		Yakima	South Fork
	2011	3021	19.6		Yakima	
	2011	28	81.2		Wild Parr	
	2011	3049	20.1		Yakima	
	2012					South Fork
	2013	3003	11.3		Yakima	
	2013	2495	27.5		Yakima	
	2014	3014	3.6		Yakima	
	2014	1249	25.4		Yakima	Cowiche Cr from Mobile Site
	2015	3017			Yakima	
	2015	1250	15.4		Yakima	Cowiche Cr from Mobile Site
	2016					
	2017					
	2018	3035	16.6		Yakima	
	2019	3013	0.40	0.28	Yakima	
	2020	No release				
Reecer Creek	2008	3001	37.41		Yakima	
	2009	2965	25.21		Yakima	
	2010	3015	23.24		Yakima	
	2011	3004	29.24		Yakima	
	2012	3026	30.52		Yakima	
	2013	3032	13.35		Yakima	
	2014	3031	7.46		Yakima	
	2015	3026	3.26		Yakima	
	2016				Yakima	
	2017				Yakima	
	2018	3069	29.96		Yakima	
2019	3005	2.56	1.10	Yakima		

	2020	Not release			
Little Naches	2009	3000	16.6	Yakima	
	2010	3072	18.3	Yakima	
	2011	3022	9.6	Yakima	
	2012	3014	20.3	Yakima	
	2013	3019	7.6	Yakima	
	2014	3012	6.6	Yakima	
	2015	3026	0	Yakima	
	2015	3004	0	Yakima	
	2015	6030	0	Yakima	
	2016	3008	2.6	Yakima	
	2017			Yakima	
	2018	3042	12.3	Yakima	
	2019	3006	*	Yakima	
	2020	No release			
Wilson Creek	2008	3000	11.4	Yakima	
	2009	3007	15.5	Yakima	
	2010	3050	12.1	Yakima	
	2011	3008	13.8	Yakima	
	2012	3020	11.2	Yakima	
	2013	1518	4.9	Yakima	Above Buried Section
	2013	1502	10.2	Yakima	Below Buried Section
	2014	3024		Yakima	
	2015	3027	8.2	Yakima	
	2016	3011	7.1	Yakima	
	2017		11.6	Yakima	
	2018	3019	48.5	Yakima	
	2019	6082	2.14	0.87	Yakima
	2020	No release			
Swauk Creek	2018	3024	2.85	Yakima	
	2019	3041	0.13	75.5	Yakima
	2020	No release			
Tieton River	2019	3010	9.16	8.6	Yakima
	2020	1289	0.93	0.71	Yakima

C.2.2. Effect of release month on parr survival rate


Parr were released at different locations from May to October (Table 11, Figure 7). Survival from release to downstream detection at McNary Dam (outmigration years 2015-2020) was highest among parr released in August ($14\% \pm 0.020$), followed by parr releases in July ($3.1\% \pm 0.40$) and June ($1\% \pm 0.4$). Lower survival rates for groups released in May and June are likely due to mortality associated with longer exposure to summer conditions. Water temperature increases in most river sections during summer while parr are rearing. In the summer time, coho salmon fry reportedly

prefer water temperatures of 50°F to 59°F (10°C to 15°C; Reiser and Bjornn 1979), while higher temperatures may cause greater mortality in the parr life stage.

Table 11. Total number of PIT-tagged parr released by month from all locations in the Yakima Basin, and survival rate to McNary Dam for each month for pooled migration years 2015-2020.

Migration Year	Parr release months and number of parr with PIT Tags				
	May	June	July	August	October
2015	1349	27262	0	0	0
2016	1648	0	24167	0	0
2018	0	0	0	21244	0
2019	0	0	39837	0	1438
2020				1289	
Total released parr with PIT tagged	2997	27262	64004	22533	1438
Survival rate ± SE	0.7 ±0.05	1.0 ±0.04	3.10 ±0.4	14.0 ±2.20	0.00 ±0.00

Figure 7. Survival probability (release location-downstream to McNary Dam) of parr released in different months. The relationship was built using tag detections in the last five migration years (2015-2020).




C.3 Effect of river flow and release month on survival rate

One of our monitoring objectives was to evaluate the effects of river flow on outmigration survival rate, and to determine whether the effect differed as a function of smolt release month (February, March and April). Data showed that the average river flow measured below Prosser Dam during April, May and June of 2020 was approximately 2050, 2107 and 1241 cubic feet per second (cfs), respectively, which were higher than the average flow in April, May and June of 2015 but slightly lower than the average April flows for 2016 through 2019 (Figure 8, Table 12). However, river flow in June 2020 was higher than in previous years with the exception of 2017. River flow for the period from June through September in all years was considerably lower than in April and May, which is typical for Western rivers. Summer flow below Prosser Dam is maintained by reservoir releases to protect aquatic life, but target flows can vary according to how much water remains in storage.

Table 12. Average monthly Yakima River flow (cfs) measured below Prosser Dam (gaging station YRPW).

Year	Months											
	1	2	3	4	5	6	7	8	9	10	11	12
2015	4793	4420	2523	1043	895	420	387	526	581	892	3549	5237
2016	2632	8603	7982	8600	3437	983	822	519	517	2086	2582	1920
2017	1696	3460	9492	8778	6959	2697	640	666	657	1463	3585	2434
2018	3038	4138	2632	5183	6183	994	574	604	542	1054	1489	1775
2019	1389	1536	3066	4444	1860	563	560	749	568	1041	1458	2122
2020	3145	4411	1470	2050	2107	1241	577	584	693	1407	2195	2120


Figure 8. Average daily Yakima River flow (cfs; blue line) and 20-day average flow (smoothed yellow line) measured near Prosser Dam from January to December (2015-2020). The boxes (red border) highlight the time period when Coho smolts were released from different locations.



A CJS model was used to evaluate the effect of river flows on outmigration survival rate for each release month (February, March and April). Among several candidate models considered, the model with river flow and release month was the most parsimonious; the best competing model was φ (\sim Dam:Year:month + RF) $p(\sim$ Dam:Year:month + RF). Based on the best CJS models that included river flow and release months as covariates (the model with the lowest QAICs), we observed a positive correlation between flow and survival rate (survival increased as flow increased) for all three months. The highest survival rates over the range of flows were found for the March release groups,

followed by April releases, and lastly February releases (Figure 9). Since Prosser was the only location with releases in each month, we could not compare the effect of release time (months) for all release groups across all locations. Survival rates among years at the Prosser location (See Figure 9) were highest for the March release groups. However, the sample size for February releases was comparatively small (4% of total releases) compared to March releases (45%) and April releases (51%).

Figure 9. The relationship between survival probability from release location to McNary Dam and the river flow at Prosser Dam for the smolt release groups each month. The relationship was devolved using the 6 years of PIT-tag data (2015-2020).




3.5 The Smolt-to-Adult Returns (McNary juvenile to Bonneville adult)

The SAR estimates varied by year and life stage at release (parr and smolt) during the study period (Table 13, Figure 10). On an average, the SAR was slightly higher for the group released as parr (SAR: $3.98 \pm 1.06\%$) compared to the SAR of the group released as smolts (SAR: $3.79 \pm 1.06\%$). The highest SAR for the group released as parr was in 2018 ($8.75\% \pm 1.66\%$), followed by the groups released in 2013 and 2008, and lowest was in 2006 (0%). For smolts, the highest SAR was for the group released in 2008 ($8.28 \pm 0.25\%$) and followed by 2013 ($8.08 \pm 0.91\%$), 2010 ($6.48 \pm 1.14\%$), and the lowest was in 2011 ($0.85 \pm 0.25\%$). The variation in SARs among years can be associated with many factors such as smolt size, release and ocean entry timing, and ocean conditions.

Table 13. Smolt-adult returns (SAR, based on juvenile detection at McNary Dam and adult detection at Bonneville Dam) for each release over migration years 2004-2020. The values with yellow color indicates the value is subject to revision if 3-ocean adults may return in 2021 from the 2018/2019 releases. “N” represents the number of fish with PIT tags released; “SE” is the standard error.

Migration year	Parr			smolt			Both		
	N	SAR	SE	N	SAR	SE	N	SAR	SE
2004	NA	NA	NA	12412	2.85%	1.07%	12412	2.85%	1.07%
2005	9576	2.78%	2.70%	31246	4.84%	0.86%	40822	4.73%	0.79%
2006	8091	0.00%	0.00%	21260	2.76%	0.52%	29351	2.60%	0.47%
2007	11129	1.98%	0.88%	30681	3.12%	0.46%	41810	2.95%	0.41%
2008	20507	6.85%	1.04%	33668	8.28%	0.85%	54175	7.77%	0.64%
2009	29988	1.63%	0.44%	33146	3.53%	0.47%	63134	2.89%	0.34%
2010	27325	4.05%	0.72%	22845	6.48%	1.14%	50170	5.02%	0.60%
2011	27229	2.68%	0.60%	25286	0.85%	0.25%	52515	1.48%	0.25%
2012	33657	2.33%	0.61%	26705	1.93%	0.42%	60362	2.08%	0.34%
2013	31973	7.56%	1.23%	21023	8.08%	0.91%	52996	7.91%	0.73%
2014	28782	2.05%	0.82%	19970	1.62%	0.33%	48752	1.70%	0.31%
2015	28611	5.41%	3.57%	17544	4.08%	1.28%	46155	4.26%	1.19%
2016	25815	5.37%	1.33%	25069	2.66%	0.50%	50884	3.22%	0.47%
2017	NA	NA	NA	14469	3.38%	0.83%	14469	3.38%	0.85%
2018	21244	8.75%	1.66%	19696	2.79%	0.80%	40940	5.23%	0.83%
2019	41275	4.29%	2.51%	20305	2.00%	0.99%	61580	2.59%	0.96%
2020	1291			13865			15156		
Average		3.98%	1.06%		3.70%	0.93%		3.79%	1.93%

Figure 10. Smolt-Adult-Return (SAR) of coho (Hatchery) for the group released as parr and smolts from 2004 to 2019 in Yakima Basin based on juveniles detected at McNary Dam (downstream migration) and the subsequently detected as returning adults at Bonneville Dam. The SAR value for outmigration year 2019 is subject to revision if 3-ocean adults return in 2021 from the 2019 migration year.



3.6 Age- distribution at return

From outmigration year 2004 through 2019, a total of 3194 returning Coho with PIT tags that were released as smolt and 1459 returning Coho that were released as parr in the Yakima Basin were detected at Bonneville Dam. Among the tagged adults released as smolts, ~85% of the returning coho were age 3 (ocean age 1) while 9% of the returns were age 2 (year ocean age 0), and 7% were age 4 (ocean age 2). For the group released as Parr, 90% of returning coho were the age 3, 10% of the returns were the age 2, and no returns were age of 4.

Table 14. Total number of PIT-tagged Coho detected at return to Bonneville Dam by ocean age (years) for the group of fish released as a life stage “smolt” (A) and the group of fish released as “Parr” (B). Values shaded yellow are subject to change based on any 3-ocean returns.

A. Smolts			Number of returning adults			Percentage (%)		
Brood year	Release Year	Migration Year	Ocean Age 0	Ocean Age 1	Ocean Age 2	Ocean Age 0	Ocean Age 1	Ocean Age 2
2002	2004	2004	1	37	10	2.08	77.08	20.83
2003	2005	2005	7	134	38	3.91	74.86	21.23
2004	2006	2006	18	169	6	9.33	87.56	3.11
2005	2007	2007	5	187	1	2.59	96.89	0.52
2006	2008	2008	113	411	37	20.14	73.26	6.60
2007	2009	2009	17	252	3	6.25	92.65	1.10
2008	2010	2010	16	272	14	5.30	90.07	4.64
2009	2011	2011	3	111	1	2.61	96.52	0.87
2010	2012	2012	4	82	18	3.85	78.85	17.31
2011	2013	2013	17	504	20	3.14	93.16	3.70
2012	2014	2014	13	80	6	13.13	80.81	6.06
2013	2015	2015	5	64	6	6.67	85.33	8.00
2014	2016	2016	9	100	19	7.03	78.13	14.84
2015	2017	2017	12	118	17	8.16	80.27	11.56
2016	2018	2018	31	86	20	22.63	62.77	14.60
2017	2019	2019	4	96	0	4.00	96.00	0.00
2018	2020	2020						
Sum/Average			275	2703	216	8.61	84.63	6.76

B. Parr			Number of returning adults			Percentage (%)		
Brood year	Release Year	Migration Year	Ocean Age 0	Ocean Age 1	Ocean Age 2	Ocean Age 0	Ocean Age 1	Ocean Age 2
2002	2003	2004						
2003	2004	2005	0	3	0	0.00	100.00	0.00
2004	2005	2006	0	6	0	0.00	100.00	0.00
2005	2006	2007	1	20	0	4.76	95.24	0.00
2006	2007	2008	30	242	0	11.03	88.97	0.00
2007	2008	2009	4	73	0	5.19	94.81	0.00
2008	2009	2010	10	246	0	3.91	96.09	0.00
2009	2010	2011	8	161	0	4.73	95.27	0.00

2010	2011	2012	13	73	0	15.12	84.88	0.00
2011	2012	2013	13	197	0	6.19	93.81	0.00
2012	2013	2014	2	30	0	6.25	93.75	0.00
2013	2014	2015	0	7	0	0.00	100.00	0.00
2014	2015	2016	2	52	0	3.70	96.30	0.00
2015	2016	2017			0			
2016	2017	2018	60	154	0	28.04	71.96	0.00
2017	2018	2019	3	49	0	5.77	94.23	0.00
2018	2019	2020			0			
Average			146	1313	0	10.01	89.99	0.00

4. Acknowledgment

We thank all of the crews whose collective fish-tagging efforts over the years made this study possible. We are also grateful to David Lind, who provided valuable comments in the draft report. We would also like to thank Daylen Isaac, who provided the hydrological data.

5. References

- Berg, L. and Fast, D. 2001. Yakima Subbasin Summary. Prepared for the Northwest Power Planning Council. August 3, 2001.
- Bradford, M.J., R.A. Myers, and J.R. Irvine. 2000. Reference points for coho salmon (*Oncorhynchus kisutch*) harvest rates and escapement goals based on freshwater production. *Canadian Journal of Fisheries and Aquatic Sciences* 57:677-686.
- Chapman, D.W. 1986. Salmon and steelhead abundance in the Columbia River in the nineteenth century. *Transactions of the American Fisheries Society* 115:662-670.
- McCann et al., 2020. Comparative Survival Study of PIT-tagged Spring/Summer/Fall Chinook, Summer Steelhead, and Sockeye DRAFT 2020 Annual Report. (<https://www.fpc.org/documents/CSS/2020CSSDraftReport.pdf>)
- Conner, M. M., S. N. Bennett, W. C. Saunders, and N. Bouwes. 2015. Comparison of tributary survival estimates of steelhead using Cormack-Jolly-Seber and Barker models: Implications for sampling effort and designs. *Transactions of the American Fisheries Society*, 144:1, 34—47.
- Davidson, F.A. and S.J. Hutchinson. 1938. The geographic and environmental limitations of the Pacific salmon (genus *Oncorhynchus*). *Bulletin Bureau of Fisheries (U.S.)* 48:667-692.
- Dunnigan, J. L., W. J. Bosch, and J. D. Hubble. 2002. Preliminary results of an effort to re-introduce coho salmon in the Yakima River, Washington. Pp. 53-75 in “Hatchery Reform: the Science and the Practice”, Proceedings of the International Congress on the Biology of Fish, July, 2002, Don MacKinlay, editor, 555 West Hastings St., Vancouver BC V6B 5G3 Canada.

- Fulton, L.A. 1970. Spawning areas and abundance of steelhead trout and coho, sockeye, and chum salmon in the Columbia River Basin-past and present. United States Fish and Wildlife Service. Special scientific report-Fisheries Number 618. Washington, D.C.
- Haeseker, S. L., J. A. McCann, J. Tuomikoski, and B. Chockley. 2012. Assessing freshwater and marine environmental influences on life-stage-specific survival rates of Snake River spring–summer Chinook Salmon and steelhead. *Transactions of the American Fisheries Society* 141:121–138.
- Haring, D. 2001. Habitat limiting factors in the Yakima River Watershed WRIs 37-39 final report. Washington State Conservation Commission, Olympia, Washington. 364 pp.
- Hassler, T. J. 1987. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Pacific Southwest): coho salmon. United States Fish and Wildlife Service, National Wetlands Research Center. Biological Report 82(11.70).
- Jones, K.K. and K.M.S. Moore. 1999. Habitat Assessment in coastal Basins in Oregon: Implications For Coho Salmon Production and Habitat Restoration. 329-340. In Knudsen, E. E., C. R. Steward, D. D. MacDonald, J. E. Williams, and D. W. Reiser. *Sustainable Fisheries Management: Pacific Salmon*. Lewis Publishers, Boca Raton.
- Kreeger, K.E., and W.J. McNeil. 1993. Summary and estimation of the historic run-sizes of anadromous salmonids in the Columbia and Yakima rivers. Prepared for Yakima River Basin Coalition, Yakima, WA.
- Laake, J. (2019) 'RMark: an R interface for analysis of capture-recapture data with MARK'. Seattle WA: US Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Alaska Fisheries Science Center.
- Lebreton, J. D., K. P. Burnham, J. Clobert, and D. R. Anderson. 1992. Modeling survival and testing biological hypotheses using marked animals: a unified approach with case studies. *Ecological Monographs* 62:67–118.
- Moyle, P. B. 2002. *Inland Fishes of California*. Berkeley: University of California Press.
- Neeley, D. 2012. Prosser-Passage Estimation Issues. Appendix F in Sampson, Fast, and Bosch, Yakima/Klickitat Fisheries Project Monitoring and Evaluation, Final Report for the Performance Period May 1, 2011 through April 30, 2012. Yakama Fisheries, Toppenish, WA.
- Petrosky, C. E., and H. A. Schaller. 2010. Influence of river conditions during seaward migration and ocean conditions on survival rates of Snake River Chinook Salmon and steelhead. *Ecology of Freshwater Fish* 19:520–536
- R Core Team. 2020. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. Available: <https://www.R-project.org/>.
- Raymond, H. L. 1968. Migration rates of yearling chinook salmon in relation to flows and impoundments in the Columbia and Snake Rivers. *Transactions of the American Fisheries Society* 97(4): 356-359.

- Reeves, G.H., F.H. Everest, and T.E. Nickelson. 1989. Identification of Physical Habitats Limiting the Production of Coho Salmon in Western Oregon and Washington. Gen. Tech. Rep. PNW-GTR-245. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 18 p.
- Reiser, D. W. and T. C. Bjornn. 1979. Influence of Forest and Rangeland Management of Anadromous Fish Habitat in Western North America- Habitat Requirements of Anadromous Salmonids. USDA Forest Service General Technical Report PNW-96.
- Sandercock, F.K. 1991. Life History of Coho Salmon. In Groot, C. and L. Margolis (Eds.). Pacific Salmon Life Histories. UBC Press, Vancouver. 564 p.
- Scheuerell, M. D., R. W. Zabel, and B. P. Sandford. 2009. Relating juvenile migration timing and survival to adulthood in two species of threatened Pacific salmon (*Oncorhynchus* spp.). *Journal of Applied Ecology* 46:983–990.
- Thompson, K. 1972. Determining stream flows for fish life. Pages 31-50 in Proceedings, Instream flow requirements workshop. Pacific Northwest River Basins Commission. Vancouver, Washington.
- Tiffan KF, Kock TJ, Haskell CA, Connor WP, Steinhorst RK. 2009. Water velocity, Turbulence, and Migration Rate of Subyearling Fall Chinook Salmon in the Free-Flowing and Impounded Snake River. *Trans Am Fish Soc* 138:373–384.
- Tuck, R.L. 1995. Impacts of Irrigation Development on Anadromous Fish in the Yakima River Basin, Washington. M.S. Thesis, Central Washington University, Ellensburg. 246 p.
- Tuomikoski, J., J. McCann, B. Chockley, H. Schaller, S. Haeseker, J. Fryer, R. Lessard, C. Petrosky, E. Tinus, T. Dalton, and R. Ehlke. 2013. Comparative survival study (CSS) of PIT-tagged spring
- White, G. C., and K. P. Burnham. Program MARK: survival estimation from populations of marked animals. *Bird Study* 46(Supplement): 120-138.
- Williams, J. G. 2006. Central Valley salmon: a perspective on Chinook and steelhead in the Central Valley of California. *San Francisco Estuary and Watershed Science* [online serial] 4(3): article 2.
- Wydoski, R.S. and R.R. Whitney 2003. *Inland Fishes of Washington*. University of Washington Press. 322 pp. Seattle WA.

Appendix F
Juvenile Outmigration Survival of Yakima Basin
Summer Chinook Smolts to Prosser and McNary
Dams, 2009-2020

Prepared by
Shubha Pandit, Melinda Davis, Bill Bosch & Mark Johnston
Yakama Nation Fisheries
P.O. Box 151, Toppenish, WA 98948, USA
May 25, 2021

TABLE OF CONTENTS

<u>EXECUTIVE SUMMARY</u>	<u>3</u>
<u>1. INTRODUCTION</u>	<u>8</u>
<u>2. METHODOLOGY</u>	<u>10</u>
2.1. GEOGRAPHICAL DISTRIBUTION: HISTORICAL AND CURRENT	10
2.2. BROOD STOCKS AND FISH DATA	11
2.3. STATISTICAL ANALYSES	13
2.3.1. SURVIVAL AND DETECTION PROBABILITY	13
2.3.2. RELATIONSHIP BETWEEN ANNUAL SURVIVAL RATE AND RIVER FLOW	14
2.3.3. RELATIONSHIP BETWEEN SURVIVAL RATE, RELEASE TIME (MONTH) AND FISH SIZE	15
2.3.4. TRAVEL TIME AND RATE OF MIGRATION OR RATE OF TRAVEL DISTANCE PER DAY	15
2.4.5. SMOLT-TO-ADULT-RETURNS (SAR)	15
2.4.6. AGE COMPOSITION OF ADULT RETURNS	16
<u>3.0. RESULTS AND DISCUSSION</u>	<u>16</u>
3.1. FISH LENGTH	16
3.2. DETECTION PROBABILITIES AT McNARY AND PROSSER	18
3.3. JUVENILE RELEASE-McNARY SURVIVAL PROBABILITY	21
3.3.1. ANNUAL JUVENILE RELEASE-McNARY SURVIVAL RATE AND ITS TEMPORAL TREND	21
3.3.2. RELEASE SITE -TO-McNARY DAM SURVIVAL RATE AMONG RELEASE LOCATIONS AND RELEASE PERIODS	23
3.3.3. COMPARISONS OF SURVIVAL RATES FROM RELEASE SITE TO PROSSER DAM AND FROM RELEASE SITE TO McNARY DAM	26
3.3.4. EFFECT OF RELEASE PERIOD AND FISH SIZE ON SURVIVAL	28
3.4. TRAVEL TIME OR RATE OF MIGRATION	29
3.5. SMOLT-TO-ADULT RETURNS	30
3.6. AGE-AT-RETURN DISTRIBUTION	31
<u>4. ACKNOWLEDGMENT</u>	<u>32</u>
<u>5. REFERENCES</u>	<u>32</u>

Executive Summary

Summer-run Chinook salmon were once widely distributed in the Yakima River basin but were extirpated by the 1970s. Since 2009, building on habitat, passage and instream flow restoration efforts, the Yakama Nation has been implementing a reintroduction program, in which summer chinook eggs are brought from Upper Columbia Basin hatcheries to the Yakama Nation's Marion Drain Hatchery for fertilization, incubation and rearing. Subyearling/presmolts are moved from the hatchery to permanent and mobile acclimation sites upriver for release as smolts into different areas of the Yakima basin. Diverse release strategies, such as releasing from different locations and experimenting with different release dates, have been utilized to maximize the likelihood of achieving stable and abundant returns of natural-origin summer Chinook to the Yakima River basin and to enhance the stability and resiliency of the population against potential environmental changes.

In 2020 a total of 768,378 subyearling summer Chinook were released, with 12,814 (about 1.67 % of the total release and relatively low in comparison to the previous years because of COVID-19 pandemic restriction) tagged for monitoring purposes, especially to evaluate juvenile survival rates and release strategies. This evaluation is an update of ongoing annual monitoring that was initiated with the first reintroductions in 2009. The main objectives of the study are to estimate survival rate of the fish released from each location in the Yakima Basin in 2020 and compare the results with previous years' results to evaluate success and discern trends. We further evaluate whether juvenile survival rate varies significantly among release locations whether the survival rate is a function of release location, release year and month, river flow and size of released fish. For data collected in prior years (2009 through 2018), a logistic regression model (Neeley 2012) was used to estimate survival probability. Since 2019, survival probability from the release locations to downstream dams and detection rate at Prosser and McNary Dams have been estimated using the Cormack-Jolly-Seber (CJS) mark-recapture model, along with other statistical analysis of travel time and the effect of river flow to answer the following research questions:

- **What was the detection rate of juvenile summer chinook at Prosser Dam and McNary Dam, and do the detection rates vary by year? If there is an annual variation in the rate of detection, what factors cause this variability?**

The average rate of detection at McNary Dam for the 2020 PIT-tag release was found to be 7.91% ($\pm 1.51\%$, mean \pm SE); whereas the detection for the fish at Prosser Dam was 27.02% ($\pm 3.65\%$). Over the years 2009-2020, detection rate varied at both dams, and was lowest in 2015, due to poor river conditions in that year. Annual variation can be due to many factors besides river conditions, such as what proportion of fish pass dams via juvenile bypass systems where detectors are installed. As spill has been increased to improve survival of juvenile fish passing dams within the Federal Columbia River Power System, a lower proportion of outmigrants enter juvenile bypass systems where PIT tags can be detected, and variations in spill percentage can make the detection rate vary among years or even from day to day.

- **What was the juvenile survival rate to McNary Dam of each of the groups released to different streams in the Yakima Basin during 2020?**

In 2020, the overall juvenile outmigration survival rates from the release site to Prosser Dam and from Prosser to McNary Dam were 43.7% \pm 4.0 and 33.6% \pm 6.3, respectively. When the survival rate of both segments was combined, survival rate of the summer Chinook in 2020 from the release site to McNary was 14.7% \pm 2.5%, which is double of the survival in the 2019 migration year (7.2% \pm 1.1). The higher survival rate in 2020 migration year might be associated with the early fish release timing and favorable river flow. In 2020, the majority of the fish were released in early (April 10 and May 8), whereas in 2019 they were released in mid-period (May 13, May 17, May 19 and May 20). Earlier releases in 2020 were exposed to more pulse flow events (five events), which might have facilitated their outmigration and increased their survival rate. Previous results have shown that juvenile downstream survival increases as river flow or pulse flow events increase.

Fish were released at three locations (Buckskin, Roza and Prosser) in 2020; but the 2019 release from Wapatox was not repeated in 2020 because of poor 2019 survival related to canal flow. Similar to the previous years, 2020 survival rates varied somewhat among release locations. Survival rate from the release location to Prosser Dam of the group released in Buckskin and Roza during 2020 were found to be 40.9% \pm 3.6% and 40.3% \pm 3.3%, respectively. From release location to McNary Dam, survival rates were 13.6 % \pm 3.4% for the group released in Buckskin, 15% \pm 5.6% for Roza; and 16.9% \pm 5.6% for the below-Prosser release group.

- **What was the overall temporal pattern of survival rates for summer Chinook during the**

study period (from 2009 through 2020), and among the groups released in different locations?

Survival rates varied by year over the period from 2009 through 2020, reflecting interannual variation in river flow, dam operations and environmental conditions such as water temperature that affect salmon during the critical juvenile life stage. The highest average annual survival rate from the release points to McNary Dam was in 2011 ($40.15\% \pm 1.94\%$) and the lowest was in 2015 ($0.73\% \pm 0.47\%$). In general, the survival rate was found to be high when were fish released early (April) as well as when river flow was high or there were more pulse flow events during the outmigration period. The relationship between the average of monthly river flow for May and June (measured below Prosser Dam) and the annual survival rate (release location to McNary Dam) from 2009 through 2020 was strong and statistically significant ($r^2=0.54$, $p=0.01$), indicating that survival rate was a function of river flow in May and June.

Overall survival rates for the period 2009 through 2020 varied among release locations as well. The highest survival rate from release to McNary Dam was for the group released from Stiles Pond ($20.3\% \pm 11.03\%$) and the second highest survival rate was for the Buckskin Slough group ($19.2\% \pm 6.81\%$). The lowest survival rate was for the group released from Wapatox Dam ($0.15\% \pm 0.14\%$) in 2019.

- **With smolts released in different months (April, May and June) to increase temporal distribution, was fish size different for different release dates? What was the effect of fish size and release month on survival rate from the release sites to Prosser Dam, and from Prosser Dam to McNary Dam?**

There was an effect of release period (April, May or June) and fish size (fork length) on the juvenile survival rate for both segments (release site to Prosser Dam; and Prosser Dam to McNary Dam). For example, in an average if the fish size with 50mm released in April, its survival rate from the release location (it can either Buckskin or Roza) to Prosser was estimated to be above 50%; whereas for the same size of fish released in June, the survival rate to Prosser Dam was estimated to be about 10%. From Prosser to McNary Dam, the relationship of size to survival rate was similar for April and May releases, but release in June depressed the Prosser-to-McNary survival rate over the entire range of fish sizes.

- **Did fish released earlier (April) enter the Columbia River estuary earlier (based on detections at Bonneville Dam) than fish released later (June), or did earlier outmigrants travel slower in order to prepare physiologically for saltwater, so that all groups entered the estuary near the same time regardless of when they were released?**

The summer Chinook releases generally exhibited immediate outmigration behavior after release, regardless of release date, but later outmigrants showed greater urgency. Travel days from Prosser Dam to Bonneville Dam for the groups released in April were 73.08 ± 37.77 days, whereas the fish released in June took only 32.70 ± 9.89 days to reach Bonneville Dam.

- **What was the rate of rate of travel from Prosser Dam to Bonneville Dam of the groups released in April, May and June?**

The rate of travel to Bonneville Dam was 7.19 km/day for the group released in April, but the rate of travel more than doubled (16.64 km/day) for the group released in June. This indicates that fish released earlier spent more time in the mainstem in order to go through the series of physiological and morphological changes that allow for a transition to life in salt water. The study suggests that regardless of when they were released, the summer Chinook seemed to enter the ocean at nearly the same time, although outmigration survival rate was higher for the early release.

- **What was the Smolt-Adult-return (SAR) of the group released each year and from each location during the study period (2009-2020)?**

SAR is the percentage of smolts that survive and return to spawn and is the metric that captures most of the cumulative impacts of the hydrosystem and ocean condition on fish, indicating the sustainability of adult returns over time. In this study, SARs were based on the percentage of smolts detected at Bonneville Dam that returned as adults to Bonneville Dam. Since summer Chinook can spend as many as 5 years in the ocean, we estimated SAR of the populations that out-migrated from 2009 through 2017. The SAR estimates varied by year during the study period. The highest SAR was for fish released in 2011 ($10.24 \pm 1.14\%$) and 2012 ($4.24 \pm 0.09\%$), whereas it was zero for the group released in the drought year of 2015. For the group of fish released in other years had about 1% SAR (Bonneville to Bonneville). The variation in SARs among years can be associated with many factors such as smolt size, release and ocean entry timing and ocean conditions.

- **What was the age composition of the adult returns?**

From the total of 1104 returning adult fish with PIT tags were detected at Bonneville Dam from 2009 through 2017, 64% were age 4 (3-year ocean age), 23% of the returns were age 3 (2- ocean), 9% were age 5 (4- ocean) and less than 1% were age of 6 (5-year ocean age). Four percent of the juveniles detected at Bonneville returned as jacks (age-2, 1-ocean).

1. Introduction

The summer Chinook (*Oncorhynchus tshawytscha*) is one of the three historical chinook runs in the Yakima River basin. Adults of the summer run first enter the Yakima River from the ocean in June, and the remainder of the summer run is shaped by flow and temperature in the lower Yakima River, which is strongly influenced by irrigation withdrawals and return flow. Unfavorable conditions can delay entry of the latter part of the summer run from the Columbia River until near the fall spawning season. Juvenile summer Chinook typically leave the Yakima River from late spring to early summer of the year after spawning. Summer Chinook were once widely distributed in the Yakima and Naches rivers (Figure 1) but were extirpated from the Yakima basin by 1970. For decades, several programs such as habitat restoration and species reintroduction were implemented in the Yakima River. With improving spawning and rearing habitat conditions made possible by habitat and instream flow restoration, with improved juvenile and adult passage in the mainstem Columbia River, and with improved ocean conditions, reintroduced adult summer chinook, along with supplemented fall chinook, are returning to the Yakima basin. Annual abundance of summer/fall Chinook at Prosser Dam on the lower Yakima River has increased from an average of just over 1000 fish from 1983 through 1999 to over 4,300 fish on average during the period 2000-2018). We have successfully achieved some level of natural production and local adaptation, however it is still unstable.

Based on 2009-2020 release data, an annual average of 282,775 summer Chinook juveniles were released in the Yakima basin (Table 1). Usually each year, eggs of the species are brought either from the Entiat or Wells hatchery (Entiat and Wells stocks) to the Yakama Nation's Prosser Hatchery for fertilization, incubation and rearing through the fall and winter. The following spring, subyearlings are moved to the acclimation sites upriver and are released directly from permanent acclimation sites on the Yakima and Naches rivers or from temporary mobile acclimation facilities operated on smaller tributary streams. Several release strategies have been utilized to maximize the likelihood of achieving stable and abundant returns of the species to the Yakima River and to enhance the stability and resiliency of the population against potential environmental changes. The strategies include releasing the juveniles into different tributaries (spatial variation) and also different dates (temporal variation). Whether one release strategy performs better than other strategies in terms of juvenile survival and smolt-to-adult return (SAR) are fundamental questions in determining whether

species management and production goals are being reached. On average each year about 12% of the total release is PIT-tagged as part of an ongoing, long-term monitoring program to refine project objectives and strategies, applying what is learned from experimentation, monitoring, evaluation and literature reviews as an adaptive management framework. This evaluation is an update of ongoing annual monitoring that began with the first reintroductions in 2009.

Juvenile survival rates often vary by seasons and years. This variation can be associated with rearing history and environmental conditions. For example, Zabel and Achord (2004) found that juvenile survival rate of wild salmonids was related to fish size (fork length), with larger juveniles having higher downstream survival. Survival rate also increases as river flow increases. Although the Yakima River is highly controlled by storage reservoirs and irrigation and hydropower withdrawals, there is still a large variation in the flow pattern within and across years, which can affect the survival rate of juvenile salmon. Ocean-type summer and fall chinook, which naturally outmigrate from Columbia River tributaries in late spring and early summer, can be harmed by rising water temperature as they attempt to leave the Yakima Basin. Based on the effect of temperature, one can postulate that survival rate should be lower if the fish are released in later months, e.g. June, than fish released as early as April. However, individuals released earlier are likely to be smaller than fish released later and closer to natural outmigration timing. There may be an interaction between fish size and release timing on survival.

The primary objectives of the study are to explore the effect of release date and fish size on survival. More specifically, our objectives are to determine the survival rate from release sites to Prosser Dam or McNary Dam of groups released at different locations in the Yakima Basin during 2020; and understand how other factors (fish size and release date) affect juvenile survival rates using the last 11 years' data (2009-2020). This information is critical for recovery of depressed Chinook stocks.

To achieve these objectives, we focused on the following research questions:

- What was the detection rate of juvenile summer chinook at Prosser Dam and McNary Dam, and do the detection rates vary by year? If there is an annual variation in the rate of detection, what factors cause this variability?
- What was the juvenile survival rate from the release sites to McNary Dam of each of the groups released to different streams during 2020?
- What was the overall temporal pattern of summer Chinook survival rate during the study

period (from 2009 through 2020), and among the groups released in different locations?

- With smolts released in different months (April, May and June) to increase temporal distribution, was fish size different for different release dates? What was the effect of fish size and release month on survival rate from the release sites to Prosser Dam, and from Prosser Dam to McNary Dam?
- What was the rate of rate of travel from Prosser Dam to Bonneville Dam of the groups released in April, May and June?
- Did fish released earlier (April) enter the Columbia River estuary earlier (based on detections at Bonneville Dam) than fish released later (June), or did earlier outmigrants travel slower in order to prepare physiologically for saltwater, so that all groups entered the estuary near the same time regardless of when they were released?
- What was the Smolt-Adult-return (SAR) of the group released in each year (each migration year) and the different locations during the study period (2009-2020)?
- What was the age composition of the adult returns?

2. Methodology

2.1. Geographical distribution: historical and current

Chinook (spring, summer, and fall runs) were native to the Yakima River basin and their historical spawning area was quite widespread in the basin (Figure 1A) but their spawning area has been reduced (Figure 1B). A major objective of the summer-run Chinook reintroduction program, begun in 2009, is to re-establish spawning in the primary historical spawning areas for this run, which are the Yakima River upstream of Wapato Dam through the canyon reach above Roza Dam, and the Naches River from the Yakima River to its confluence with the Tieton River (Figure 1C). The uppermost acclimation and release sites designated in the reintroduction program were located to facilitate adult homing throughout this historical geographical distribution, while the lower sites (Marion Drain downstream to the river mouth) were chosen to maximize survival rates and improve opportunities to collect returning adults as we work to establish a localized brood source (Figure 1D).




Figure 1. Historical (A) and current (B) summer Chinook spawning area; and the locations/tributaries/river segments where summer Chinook juveniles were introduced from 2009 through 2020.

2.2. Brood stocks and fish data

Every year, eggs of summer Chinook have been brought to Yakima basin either from the Wells Hatchery which is located in Pateros, WA (especially for the years from 2009-2020) or Entiat Hatchery (2018-2019) or Wenatchee Stock from Eastbank Hatchery (2010) (See Figure 2). The adult fish were spawned at either Wells or Entiat; green eggs and milt were transferred to the YN Prosser Hatchery for fertilization, incubation and rearing. Prior to migration year 2020, presmolt subyearling juveniles were acclimated at as many as five sites upriver (Stiles Pond, Buckskin Slough (Nelson Springs), Marion Drain Hatchery, Roza Dam and Wapatox Diversion), but for migration year 2020

all subyearling juveniles were transferred to just three locations (Buckskin Slough, Prosser Hatchery and Roza Dam). Temporary mobile acclimation facilities were used in some years in upstream tributary streams of the Naches and Yakima rivers.

On average 33,425 juvenile summer Chinook were PIT-tagged per year (the range was 49,894 in 2011 to 12,814 in 2020) prior to release from April through June (Figure 1D). In 2020, a total of 768,378 subyearling summer Chinook were released from the Buckskin, Roza acclimation sites, along with a group released directly from Prosser Hatchery, including 12,814 fish with PITtags (Table 1 and Table 2) between April 10th and May 12th, 2020 (Table 2). The tagging effort in 2020 compared to previous years was low due to COVID-19 pandemic restrictions.

Table 1. Annual releases of summer Chinook run with and without PIT-tags and the number and percentage of PIT tags in each release.

Year	Total Release		
	Total release (with & without PIT tags)	PIT-tags	PIT tag Percentage (%)
2009	180,911	30,045	16.61
2010	200,747	29,997	14.94
2011	215,770	49,893	23.12
2012	197,103	29,996	15.22
2013	136,563	40,507	29.66
2014	254,881	30,278	11.88
2015	277,448	34,457	12.42
2016	37,000	37,000	100.00
2017	244,499	34,826	14.24
2018	74,000	30,131	40.72
2019	806,000	41,143	5.10
2020	768,378	12,814	1.67
Average	282,775	33,425	12%

All regional PIT tag detection data including release and detection history are available in the PTAGIS database maintained by the Pacific States Marine Fisheries Commission. We queried PTAGIS (<https://www.ptagis.org/>) in April 2021 to retrieve available PIT-tag detection information for all summer Chinook juveniles released in the Yakima Basin from migration year 2009 through 2020 (Table 2). For each fish with a PIT-tag code, we constructed a detection history: a record indicating all detection locations and whether the tagged fish was detected or not detected at each juvenile detection site, focusing on Prosser, McNary, John Day and Bonneville dams (PRO, MCJ, JDJ, B2J, BCC), and by the Estuary Towed Experimental Array (Twx).

Columbia River Power System (FCRPS) to estimate survival rates for juvenile salmon and steelhead (Tuomikoski et al. 2013). The model uses multiple detections of individually marked fish at several dams with PIT-tag detection capabilities (i.e. antenna arrays). One of the assumptions of the CJS model is that there is no immigration or emigration during capture and recapture intervals, which is valid for discrete tag groups migrating through the hydrosystem (which involves passage at several hydroelectric dams) because all fish in the tag group are moving in one direction and over a relatively short period (Conner et al. 2015). All of the assumptions of the CJS models are considered to be met.

To determine how release period (April, May or June) and fish size affect the survival rate from the release location to Prosser, and from Prosser to McNary, we introduced fish size and release period as covariates in the CJS model. This CJS model was built within RMark (Laake 2019) in R, an extension of Program MARK (White and Burnham 1999). The detailed methodology is found in Appendix F of the main report. In 2020, 12,814 tagged fish were tagged, however the first detection dates of 81 tagged fish were found to be earlier than the release date. This might have been due to some of the fish escaped early during tagging or acclimation. We excluded these 81 fish from further analysis.

2.3.2. Relationship between annual survival rate and river flow

Several environmental factors are known to influence downstream smolt survival, and river flow is among the most impactful (Raymond 1968; Connor et al. 2003; Tiffan et al. 2009). We therefore further evaluated whether there was a relationship between the annual survival rate and the average river flow for two summer months (May and June) measured below Prosser Dam. We chose only May and June because most of the juvenile summer Chinook were released from the end of April (29th) to the first week of June (5th) from 2009 through 2020 (See table 2), and they usually leave the Yakima River within 3 or 4 weeks after release. Given this timing, May and June flow can be the most influential factor for the outmigration of this run of Chinook. We downloaded river flow data for the Bureau of Reclamation gaging station (YRPW) located below Prosser Dam in the Yakima River, using the Hydromet site: <https://www.usbr.gov/pn/hydromet/yakima/yakwebarcread.html>, which was accessed in April 2021. A univariate linear relationship between the average survival rate of each migration year and the average river flow (May and June) of each year was built to determine whether the average annual survival rate is a function of river flow.

2.3.3. Relationship between survival rate, release time (month) and fish size

Among the available PIT-tagged fish, only a few had fish length information, so we selected only those tagged fish which had fish length information available for the analysis. Fish release dates were categorized by month. As mentioned under subheading 2.3.1, we used fish length and release month as covariates in the CJS model. Using this model, the average survival rates from release location to Prosser Dam, and from Prosser to McNary Dam were estimated for the release groups with different release months (April, May, June) and the different average fish lengths.

2.3.4. Travel time and rate of migration or rate of travel distance per day

Travel time was estimated as the difference between either the date of release or the date of detection at Prosser Dam (site PRO), and the date of detection at Bonneville Dam (juvenile site B2J or BCC). For fish released below Prosser Dam, we estimated travel time as the difference between the release date and the date of detection at Bonneville Dam. For fish released above Prosser Dam, travel time from Prosser to Bonneville was estimated as the difference between the date of detection at Prosser Dam and the detection date at Bonneville Dam. We estimated travel time for each of three release months (April, May and June). Migration rate or rate of travel was calculated as length of the reach of interest (km) divided by travel time in days for the group.

2.4.5. Smolt-to-Adult>Returns (SAR)

SAR, which is the percentage of smolts that survive and return as an adult to spawn, is a metric that captures most of the cumulative impacts of the hydro-system and ocean conditions on anadromous fish, indicating how sustainable the returns of adults are over time. The SAR was estimated as the percentage of smolts detected at Bonneville Dam returning as adults to Bonneville Dam using the following equation for each year and release group:

$$U_{at\ BON} / J_{at\ BON}$$

Where, $U_{at\ BON}$ is a total number of PIT tagged fish detected at Bonneville Dam both during outmigration as a juvenile and immigration as adults. $J_{at\ BON}$ is the total number of fish detected at Bonneville Dam as juveniles. Because summer Chinook can spend as many as 5 years in the ocean, we estimated SAR of the populations that out-migrated from 2009 through 2017 (migration year).

The variance of SAR estimates for each category was computed by a non-parametric bootstrap re-sampling method (Efron and Tibshirani 1993; Manly 1997). For each sample data set (release group for each migration year), individual capture histories were resampled with replacement. One thousand bootstrap sample data sets were constructed and 1000 estimates of SAR were generated. Statistical bias was assessed as the difference between the mean of the bootstrap replicates and the point estimate derived from the original data (Efron and Tibshirani, 1993). Due to the non-normal distribution of bootstrap SAR estimates, bias correction was used to construct 95% confidence intervals as suggested by Manly (1997).

2.4.6. Age composition of adult returns

Age composition of adult returns was estimated by the proportion of each age class of adult return detected at Bonneville Dam by each migration year.

3.0. Results and discussion

3.1. Fish length

An average 33425 PIT-tagged juvenile summer Chinook were released per year from 2009 through 2020, but only 42,868 had the size information, which was about 13% of the total of PIT-tagged fish released during the study period. Based on the available data, the average size of the fish (fork length) at the time of tagging was 71 mm (Figure 2, Table 3). However, the size of the fish of the groups released in different months (March, April and May) was found to be different. We hypothesized that fish released later would be bigger than the fish released earlier, but we found that fish released in May were bigger when tagged than the fish released in June. The average fork lengths of the groups released in April, May and June were 66.98 ± 0.115 mm, 74.17 ± 0.06 mm, and 63.08 ± 0.026 mm at the time of tagging, respectively. Not getting the same result as we hypothesized might be due to a number of reasons. One possible reason is that the sample sizes (N) were different among the groups released in different months. There was a very large number of lengths in the May release group (38,874); whereas the June release group had only 1844 measured fish (Figure 2, table 3). It is likely the smaller sample size did not represent the actual range of sizes of

the fish released in June. Another reason could be differences in incubation and rearing temperature among groups from different hatcheries with different water sources.

Figure 2. Frequency (count) by fish length (fork length, mm) at the time of tagging for all releases made in April, May and June from 2009 to 2020.




Table 3. Average fish size (mm) at the time of tagging by releasing year and month (April, May, June). The number “n” represents the subset of fish with length data in the PIT Tag Information System (PTAGIS; <http://www.ptagis.org>) database.

Year	April			May			June			Pooled		
	n	Mean	SE	n	Mean	SE	n	Mean	SE	n	Mean	SE
2009							30036	63.17	0.03	30036	63.17	0.03
2010				22711	74.62	0.055				22711	74.62	0.05
2011	1467	67.58	0.14	3619	91.33	0.388				5086	84.48	0.32

2012				3095	68.27	0.131				3095	68.27	0.13
2013				3000	68.51	0.121				3000	68.51	0.12
2014				1268	63.83	0.105	1845	61.89	0.11	3113	62.68	0.10
2015	702	66.75	1.00	3071	69.41	0.182				3773	68.92	0.27
2016				1106	75.65	0.649				1106	75.65	0.65
2017				918	66.20	0.728				918	66.20	0.73
2019				264	75.21	0.423				264	75.21	0.42
2020				4974	75.71	0.094				4974	75.71	0.09
Mean		67.16			72.87			62.53			71.22	

3.2. Detection Probabilities at McNary and Prosser

The rate of detection at Prosser Dam varied with diversion rate; detections in the juvenile bypass system depend on irrigation and hydropower diversions during the outmigration season. Diversion rate in turn depends on Yakima River flow, which fluctuates more than the highly-regulated Columbia River discharge. The rate of detection of juvenile summer Chinook at McNary Dam also varied among years (Table 4), but within a smaller range. Variation in the detection rate at McNary Dam might be due to how surface-passage structures are operated. As with Prosser Dam, the detection rate at Columbia River dams depends upon the proportion of fish that enter juvenile bypass systems where detectors are installed. In recent years, increasing spill and the use of surface-passage structures (spillway weirs) at dams are a primary management strategy to increase survival of juvenile fish passing dams within the Federal Columbia River Power System. Greater use of spillways results in a lower proportion of fish entering juvenile bypass systems where PIT tags can be detected (Widener et al. 2018), and fluctuations in spill and flow can produce variable detection rates among years or within a migration season.

Table 4. Annual detection rate (in percent) at McNary Dam (and its Standard Error, SE) during the period from 2010 through 2020. Enumeration of fish fate (release/detection histories) is coded by detection (1) and no detection (0). For example, at McNary Dam: the code “1.0.0.” means no juvenile detection after release, “1.0.1” means not detected at McNary Dam but detected downstream of McNary Dam, “1.1.0” means detected at McNary Dam but not downstream, and “1.1.1” means detected at both McNary Dam and downstream.

Year	Re-leased	Prosser Dam						McNary Dam					
		Release/detection history (Number of PITtagged fish)				Detection Prob. % (p)		Release/detection history (Number of PITtagged fish)				Detection Prob. % (p)	
		1.0.0.	1.0.1.	1.1.0.	1.1.1.	p	SE	1.0.0.	1.0.1.	1.1.0.	1.1.1.	p	SE
2010	29747	24009	667	4712	359	34.99	1.4	28021	700	865	161	18.7	1.32
2011	49365	45699	2392	1187	87	3.5	0.3	44591	2295	2151	328	12.5	0.6
2012	29821	26562	891	242	126	12.38	1.03	27335	1469	830	187	11.3	0.7
2013	30186	20328	1232	8210	416	28.57	0.2	27618	920	1360	288	23.9	1.2
2014	30524	24590	278	5506	150	35.04	2.3	29796	300	361	67	18.3	2
2015	33829	33150	17	662	0	1.95	0	33785	27	15	2	6.88	4.7
2016	35546	Released all fish below Prosser Dam						32451	932	1933	230	19.8	1.16
2017	17534	15051	289	2098	96	24.93	0.24	16545	604	308	77	11.3	1.21
2018	30130	28241	126	1749	14	10	2.53	29867	123	11	27	18	3.14
2019	41151	37765	185	3161	40	17.77	2.5	40592	334	199	26	7.22	1.36
2020	12729	10823	108	1758	40	27.02	3.65	12290	291	123	25	7.91	1.51

In 2020, a total 12,729 juvenile summer Chinook with PIT tags were released from the 3 locations (Buckskin Slough, Roza juvenile bypass and below Prosser Dam). The average rate of detection at McNary Dam for the 2020 release was found to be 7.91% ($\pm 1.51\%$ SE, see table 4), whereas the detection at Prosser was $27.02 \pm 3.75\%$. The highest detection rate for a release group at McNary Dam was for Prosser releases ($9.5 \pm 3.4\%$, see table 4). The group released below Prosser would be expected to have low mortality from release to McNary Dam compared to the groups released upstream and considerably farther from McNary Dam. In general, travel distance is considered to be an important factor influencing survival rate. As travel distance increases, mortality also increases due to higher risk of predation and changing environmental conditions.

For all upstream release groups combined, the average (pooled) detection rate at Prosser for 2020 groups was about $27.75 \pm 3.6\%$; whereas the average detection rate at McNary was only $7.91 \pm 1.31\%$.

Among the release groups, the highest detection rate at Prosser was $46.0 \pm 7.0\%$ for the group released below Roza Dam (Table 4A), whereas the detection rate of this group at McNary Dam was about $6.7 \pm 2.6\%$ (Table 5B).

Table 5. Detection rate at Prosser (PRO) and McNary Dam (McN) for the groups of sub-yearling summer Chinook released from 2010 to 2020.

5A. Detection Probability at Prosser Dam (PRO)

Migration year	Stiles	Buckskin	Marion Drain	Roza	Prosser	Yakima mouth	Wapatox	Pooled
2010	0.35 ± 0.015							0.35 ± 0.015
2011	0.011 ± 0.004	0.048 ± 0.005						0.035 ± 0.003
2012		0.251 ± 0.022	0.072 ± 0.013		NA			0.124 ± 0.01
2013		0.271 ± 0.004		0.304 ± 0.004				0.286 ± 0.002
2014		0.321 ± 0.025		0.488 ± 0.056		NA		0.35 ± 0.023
2015		$0.948 \pm 15655^*$		$0.022 \pm 0.001^*$	NA			0.02 ± 0
2016					NA			
2017								0.249 ± 0.002
2018				0.106 ± 0.027			0.026 ± 0.001	0.1 ± 0.025
2019		0.462 ± 0.138		0.507 ± 0.061	NA		$0.939 \pm 1857^*$	0.178 ± 0.025
2020		0.321 ± 0.064		0.46 ± 0.07	NA			0.270 ± 0.036

5B. Detection Probability at McNary Dam (McN)

Migration year	Stiles	Buckskin	Marion Drain	Roza	Prosser	Yakima mouth	Wapatox	Pooled
2010	18.7 ± 1.3							18.7 ± 1.32
2011	10.9 ± 1	13.5 ± 0.9						12.5 ± 0.6
2012		9.5 ± 1.1	12.6 ± 1.3		12.4 ± 1.7			11.29 ± 0.7
2013		25.7 ± 1.6		21.5 ± 1.9	11.8 ± 7.8			23.89 ± 1.2
2014		18.7 ± 2.3		14.1 ± 4.3	33.3 ± 15.7	NA		18.25 ± 2
2015		$0.9485 \pm 15655^*$		$0.0222 \pm 0.001^*$	19.8 ± 1.2			6.88 ± 4.7
2016					12.5 ± 3.7			19.79 ± 1.16
2017								11.3 ± 1.21
2018				18 ± 3.3			18.2 ± 11.6	18 ± 3.14
2019		5.6 ± 5.4		5.9 ± 2.4	7.9 ± 1.7		$0.94 \pm 1857^*$	7.22 ± 1.36
2020		7.9 ± 2.2		6.7 ± 2.6	9.5 ± 3.4			7.91 ± 1.51

* Model convergence issue due to no downstream detections.

Note: Some of the juveniles were detected at John Day Dam but not detected at BON. The number of detections attributed to Bonneville Dam (BON) includes fish that were detected either at John Day Dam (JDJ), Bonneville Dam (B2J or BCC), or by the Estuary Towed Experimental Array (TWX).

3.3. Juvenile Release-McNary Survival Probability

3.3.1. Annual juvenile Release-McNary Survival rate and its temporal trend

The survival rate of juvenile summer Chinook from release to McNary Dam varied among years (Figure 3; Table 6). The highest average annual survival rate was in 2011 ($40.15 \pm 1.94\%$) and the lowest was in 2015 ($0.73 \pm 0.47\%$). In 2020 the average annual survival rate from all release locations to McNary Dam was $14.7 \pm 2.5\%$, which was higher than in 2019 (7.22 ± 1.35) and 2018's survival rate ($2.58 \pm 0.41\%$, Figure 3; Table 6).




Figure 3. Average annual survival rate (release to McNary Dam) of juvenile summer Chinook released from 2010 through 2020.

It is important to understand why the survival rate varied among years. The juvenile survival might have been affected by many factors such as different brood stocks, release timing or river flow and including other variables. On an average the survival rate in 2011 was high (Table 6 and Figure 3). Looking at individual groups in 2011, the highest survival rate was for the group released into Buckskin Slough on the lower Naches River, which was released before May 10th, and its brood

stock was Wenatchee (Eastbank hatchery, see Table 2). For Stiles Pond, also on the lower Naches, in 2011, the survival rate was also high even though these fish were released in the middle period (May 11 through May 25th). The brood stock for Stiles was from Priest Rapids Hatchery.

Despite different brood stocks, release times and release locations, both groups (Stiles and Buckskin) had relatively high survival rates in 2011 compared to other years. These results suggest that other external factors might have played a role in increasing the survival rate. We explored whether Yakima River flow below Prosser Dam had an effect on survival rate. We built the univariate relationship between the average river flow for May and June and the annual survival rate, and found that survival rate was strongly influenced by May and June average river flow ($R^2=0.54$, $p=0.01$, see Figure 4). It indicates that survival rate was a function of river flow, however the river flow was able to explain only about 54% of the annual variation in survival rate. Temperature or predation or interactions between temperature and flow or other factors might also have affected the survival rate. Further investigations, especially into how release period and fish size affected survival rate, are discussed in a later section (See 3.3.4. Effect of release period and fish size on survival).

Table 6. Total released smolt population, survival rate from release locations to McNary Dam and its Standard Error (SE) and the average river flow for May and June of each year from 2010 through 2020.

Outmigration /Release Year	Smolts Released	Survival Rate (%)		Average River flow (cfs) (May & June)
		Average	SE	
2010	29747	18.44	1.22	2896
2011	49321	40.15	1.94	9305
2012	29821	30.20	1.89	7102
2013	30186	22.89	1.09	3842
2014	30524	7.68	0.79	3131
2015	33829	0.73	0.47	699
2016	35546	30.74	1.73	2559
2017	17534	19.41	1.88	5400
2018	30028	2.58	0.41	4064
2019	41071	7.22	1.35	1307
2020	12729	14.70	2.50	1795




Figure 4. Relationship between average May-June river flow and the annual survival rate of juvenile summer Chinook from all release sites to McNary Dam for the years 2010 through 2020. Each point with error bar is the average survival rate and its 95% confidence interval (CI) for each year. The dotted line with the shaded area is the predicted linear trend (survival rate vs. river flow) and its 95% CI.

3.3.2. Release site -to-McNary Dam survival rate among release locations and release periods

As mentioned above, the average annual survival rate from all release sites to McNary Dam varied by year. The survival rate also varied by release location (Table 7 and Figure 5 and 6). However, when the data were pooled by release period (Early, Mid and Late), the groups released earlier had about $19.39 \pm 10.75\%$ survival rate, whereas the mid and late releases had survival rates of $16.27 \pm 3.23\%$ and $7.6 \pm 4.48\%$, respectively. When releases were pooled by location (see figure 6), the highest survival rate was for the group released from Stiles Pond ($20.3 \pm 11.03\%$) and the second

highest survival rate was for the Buckskin Slough group ($19.2 \pm 6.81\%$). The lowest survival rate was for the group that was released from the Wapatox bypass in 2018 and 2019 ($0.15 \pm 0.14\%$, Figure 5). Low survival for the release was most likely due to the low flow in the bypass because the bypass was designed for a much higher flow. (The Wapatox power plant was closed in 2003, with only irrigation deliveries remaining in the Wapatox canal.) There were no releases from Wapatox in 2020 while a solution to the problem was put in place. For the 2021 release we deployed a pipe from the mobile acclimation units directly to the bypass exit pipe to the river, avoiding the slack water in front of the canal's fish screens. In 2021, we also released two raceways directly into the river near the Wapatox diversion for comparison.

Table 7. Survival rate (%) of summer Chinook from each release site to McNary Dam from 2009 through 2020 for the 7 release sites. The survival rate and its standard Error (SE) are given for the 2019 and 2020 estimates. Early, Mid and Late releases correspond to the period through May 10; May 11 through May 25; and the period after May 25 respectively.

Year	Stiles		Buckskin			Marion drain	Roza			Prosser		Yakima River Mouth	Wapatox
	Mid	Late	Early	Mid	Late	Mid	Early	Mid	Late	Early	Mid	Early	Mid
2009		1.5											
2010	19.7												
2011	39.7		43.7										
2012				37.2		35.8					20.8		
2013				29.8					20.9				
2014				18.3	3.2				4.8				
2015				0.01	0			0.07	0		2.6		
2016												31.2	
2017								19.4			19.6		
2018								4.9					0.3
2019				2.3 ± 2.1				11.0 ± 4.2			17.9 ± 3.7		0.0
2020				13.6 ± 3.4				15.0 ± 5.6			16.9 ± 5.5		

Note: the survival rate estimates from 2009 through 2018 are from a previous report (Neeley 2019, Appendix G).




Figure 5. Juvenile survival rate from release site to McNary Dam for summer Chinook groups released at different locations from 2009 through 2020.




Figure 6. Average survival rate of juvenile summer Chinook to McNary Dam by release location from 2009 through 2020. Marion Drain and the Yakima River mouth each had only one estimate so that there was no variance.

3.3.3. Comparisons of survival rates from release site to Prosser Dam and from release site to McNary Dam

Survival rates from release site to McNary in 2020 were much lower than survival from release site to Prosser. For example, the survival rate for the group released from Buckskin Slough was about 40.9% to Prosser; but from Prosser to McNary it was only 33.4%; and for overall survival rate from release site-to-McNary was 13.6% (Table 8 and Figure 7). Mortality from Prosser to McNary is likely concentrated in the lower Yakima river and the delta at its confluence with the Columbia River and related to high water temperature and documented heavy predation. From the delta, fish must travel 69 river kilometers (rkm) down the Columbia River to detection facilities at McNary Dam in addition to the 76 rkm from Prosser Dam to the delta, but on the basis of Columbia River smolt survival studies it is likely that most of the observed juvenile summer Chinook mortality occurs in the Yakima River from Prosser to the delta.

Table 8. Juvenile summer Chinook survival rate from each release site to Prosser Dam, from Prosser Dam to McNary Dam, and from release site to McNary in 2019 and 2020. “N” is the number of PIT tags. In 2020, 81 Buckskin fish were found detected earlier than their release date, and were excluded from the analysis.

Year	Release Site	N	Release site to PRO		PRO to McN		Release site to McN	
			Survival rate	SE	Survival rate	SE	Survival rate	SE
2019	Buckskin	10365	0.214	0.042	0.105	0.100	0.023	0.021
	Roza	10254	0.387	0.029	0.284	0.110	0.110	0.042
	Prosser	10266	NA	NA	0.179	0.037	0.179	0.037
	Wapatox	10266	0.001	74.100*	0.000	0.000	0.000	0.000
	Pooled	41151	0.356	0.028	0.202	0.037	0.072	0.012
2020	Buckskin	4920	0.409	0.036	0.334	0.087	0.136	0.034
	Roza	4996	0.403	0.033	0.372	0.141	0.150	0.056
	Prosser	2813	NA	NA	0.169	0.055	0.169	0.056
	Pooled	12729	0.437	0.029	0.336	0.063	0.147	0.025

* Indicates the model convergence issue due to no downstream detections.




Figure 7. Juvenile summer Chinook survival rate from each release site-to-Prosser Dam, from Prosser-to-McNary Dam, and from release site-to-McNary in 2019 and 2020.

3.3.4. Effect of release period and fish size on survival

As mentioned in the methodology section, we built the CJS model with release period (month) and fish size as covariates using the fish size information available (N=42,868, see chapter 3.1, Fish Size). Figure 9 (left side) shows that release period affected juvenile survival and that for fish with fork length of 50mm in April, the survival rate to Prosser Dam exceeded 50%, whereas 50mm fish released in June had a survival rate of approximately 10%. For the largest fish, there seemed to be no effect of release timing on the survival rate.

From Prosser-to-McNary Dam (right side of Figure 9), the relationship of size to survival rate was similar for April and May releases, but release in June depressed the Prosser-to-McNary survival rate over the entire range of fish sizes. Standard errors for the groups released in April and May were found to be large, which might be due to small sample size. As mentioned in 3.1., the sample size was relatively low for the group release in April (2,155) and June (1,844) compared to May release (38,874).




Figure 8. Effect of release period and fish size on the rate of survival from the release site to Prosser Dam, and from Prosser Dam to McNary Dam. The shaded area is the standard Error (SE).

In 2020 the overall juvenile outmigration survival rates from the release site to Prosser Dam and

from Prosser Dam to McNary Dam were $43.7\% \pm 4.0$ and $33.6\% \pm 6.3$, respectively, but the overall survival rate of summer Chinook in 2020 from the release site to McNary was $14.7\% \pm 2.5\%$, which is double what it was in the 2019 migration year ($7.2\% \pm 1.1$). The higher survival rate in 2020 migration year can be associated with either early fish release timing or favorable river flow or a combination of both. In 2020, the majority of the fish were released in the early period (April 10 and May 8), whereas in 2019 they were released in the middle period (May 13, May 17, May 19 and May 20, Table 2). Since these fish were released early they were exposed to 5 pulse flow events (reservoir releases specifically for fish outmigration, Figure 9), compared to 4 such events in 2019, which might have facilitated their migration down the Yakima River. Previous results have shown that juvenile downstream survival increases as river flow increases.




Figure 9. Daily river flow and water temperature during the summer chinook outmigration period (March -July) and release window for 2019 and 2020.

3.4. Travel time or rate of migration

Summer Chinook generally exhibited immediate outmigration behavior after release, regardless of release date, but later outmigrants showed greater urgency. Travel times from Prosser Dam to

Bonneville Dam for the groups released in April were about 73.08 ± 37.77 days, whereas the fish released in June took only 32.70 ± 9.89 days to reach Bonneville Dam (Table 9).

Table 9. Travel days \pm SE and rate of travel (km/day \pm SE) from Prosser to Bonneville Dam for the groups released in April, May and June from 2010 through 2020.

Release Month	Number of PIT Tags	Travel days	Rate of migration (km/day)
April	24,555	73.08 ± 37.77	7.19 ± 0.10
May	28,318	65.08 ± 14.03	8.15 ± 0.04
June	20,140	32.70 ± 9.89	16.64 ± 0.03

The distance between Prosser Dam and Bonneville Dam is normally given as 381 rkm and the rate of travel over that distance was 7.19 km/day for the group released in April; but the rate more than doubled (16.64 km/day) for the group released in June. The slower rate of travel for earlier releases indicates that fish released earlier spent more time in the mainstem in order to go through the series of physiological and morphological changes that allow for a transition to life in salt water. Before entering the ocean, anadromous species must change their osmoregulation process, undergoing physical adaptations of their gills and kidneys that build a tolerance to salt water. The study suggests that regardless when they were released, the summer Chinook seemed to enter the ocean at nearly the same time, although outmigration survival rate was higher for the early release.

3.5. Smolt-to-Adult Returns

SAR which is the percentage of smolts that survive and return to spawn and captures most of the cumulative impacts of the hydro system and ocean condition on fish, telling us how sustainable the returns of adults are over time. The SAR estimate was based on the percentage of smolts detected at Bonneville Dam that returned as adults to Bonneville Dam. In general, the SAR varied by year during the study period. The highest SAR was for the fish released in 2011 ($10.24 \pm 1.14\%$) and 2012 ($4.24 \pm 0.09\%$), whereas it was zero for the group released in 2015 (see Table 10). The groups of fish released in other years averaged about 1% SAR from Bonneville juvenile to Bonneville adult. The variation in SAR among years can be associated with many factors such as smolt length, release timing, ocean conditions etc. Since SAR and juvenile survival both were high in 2011 and 2012 compared to other years, the higher SAR seems to be related to higher juvenile downstream survival.

Table 10. Smolt-adult returns (based on Juvenile and adult detection at Bonneville Dam) for each release over migration years 2010-2017. The value with yellow color indicates the value is subject to revision if 4-ocean adults may return in 2021 from the 2017 releases.

Migration year	Stiles	Buckskin	Marion Drain	Roza	Prosser	Yakima mouth	Wapatox	Pooled
2010	1.25 ± 0.46							1.25 ± 0.46
2011	10.2 ± 2.06	10.22 ± 1.35						10.21 ± 1.14
2012		4.10 ± 1.4	3.29 ± 1.18		6.89 ± 2.71			4.24 ± 0.9
2013		2.08 ± 0.86		1.46 ± 0.81				1.80 ± 0.60
2014		0.69 ± 0.6		0				0.69 ± 0.6
2015		0		0	0			0
2016					1.07 ± 0.48			1.07 ± 0.48
2017				0.88 ± 0.49	1.97 ± 1.90			1.02 ± 0.53
2018	With incomplete returns, no SAR was calculated for 2018-2020 releases							
2019								
2020								

3.6. Age-at-return distribution

From the total of 1104 returning adult fish with PIT tags were detected at Bonneville Dam from 2009 through 2017, 64% were age 4 (3-year ocean age), 23% of the returns were age 3 (2- ocean), 9% were age 5 (4- ocean) and less than 1% were age of 6 (5-year ocean age). Four percent of the juveniles detected at Bonneville returned as jacks (age 2, 1-ocean; Table 11).

Table 11. Total number of PIT-tagged fish detected at return to Bonneville Dam by ocean age (years). Values shaded yellow are subject to change based on 4-ocean returns.

Migration Year	Number of returning adults						Percentage (%)				
	Age 1	Age 2	Age 3	Age 4	Age 5	Total	Age 1	Age 2	Age 3	Age 4	Age 5
2010	7	21	79	19	0	126	5.56	16.67	62.70	15.08	0.00
2011	33	170	339	53	2	597	5.53	28.48	56.78	8.88	0.34
2012	0	19	106	32	0	157	0.00	12.10	67.52	20.38	0.00
2013	1	49	40	8	0	98	1.02	50.00	40.82	8.16	0.00
2014	1	2	14	1	0	18	5.56	11.11	77.78	5.56	0.00
2016	4	26	47	2	0	79	5.06	32.91	59.49	2.53	0.00
2017	2	3	24	0	0	29*	6.90	10.34	82.76	0.00	0.00*
2018											

2019	With incomplete returns, no SAR was calculated for 2018-2020 releases					
2020						
Average		4.23	23.09	63.98	8.66	0.05

4. Acknowledgment

We thank all of the crews whose collective fish-tagging efforts over the years made this study possible. We are also grateful to David Lind who reviewed, edited and provided valuable comments in the draft report. We would also like to thank Daylen Isaac who provided the hydrological data.

5. References

- Conner, MM, S. N. Bennett, W. C. Saunders, and N. Bouwes. 2015. Comparison of tributary survival estimates of steelhead using Cormack-Jolly-Seber and Barker models: Implications for sampling effort and designs. *Transactions of the American Fisheries Society*, 144:1, 34—47.
- Laake, J. 2013 ‘RMark: an R interface for analysis of capture-recapture data with MARK’. Seattle WA: US Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Alaska Fisheries Science Center.
- Lebreton, JD, Burnham, KP, Clobert, J & Anderson, DR. 1992. Modeling survival and testing biological hypotheses using marked animals: a unified approach with case studies, *Ecological Monographs*, 62, pp. 67-118.
- Neeley, D. 2012. Prosser-Passage Estimation Issues. Appendix F in Sampson, Fast, and Bosch, Yakima/Klickitat Fisheries Project Monitoring and Evaluation, Final Report for the Performance Period May 1, 2011 through April 30, 2012. Yakama Fisheries, Toppenish, WA.
- Tuomikoski, J, J. McCann, B. Chockley, H. Schaller, S. Haeseker, J. Fryer, R. Lessard, C. Petrosky, E. Tinus, T. Dalton, and R. Ehlke. 2013. Comparative survival study (CSS) of PIT-tagged spring White, GC., and K. P. Burnham. *Program MARK: survival estimation from populations of marked animals. Bird Study* 46(Supplement): 120—138.
- Widener, DL., James R. Faulkner, Steven G. Smith, Tiffani M. Marsh, and Richard W. Zabel. 2018. Survival Estimates for the Passage of Spring-Migrating Juvenile Salmonids through Snake and

- Columbia River Dams and Reservoirs, 2017. Northwest Fisheries Science Center, NOAA.
Submitted to Bonneville Power Administration, 2018 February
- Williams, B. K., Nichols, JD, & Conroy, MJ. 2002. Analysis and Management of Animals Populations. San Diego, CA: Academic Press.
- Zabel, RW, Achord S. 2004. Relating size of juveniles to survival within and among populations of Chinook salmon. *Ecology* 85:795–806
- Zydlowski, GB, Stich DS & McCormick SD. 2014. Photoperiod control of downstream movement of Atlantic salmon *Salmo salar* smolts. *Journal of Fish Biology* 85: 1023– 1041.