




Monitoring and evaluation efforts in the Yakima River Basin are the result of a cooperative effort by many individuals from a variety of agencies including the Yakama Nation Fisheries Program, the Washington Department of Fish and Wildlife, the United States Fish and Wildlife Service, the Columbia River Inter-Tribal Fish Commission, the National Oceanic and Atmospheric Administration Fisheries department as well as some consultants and contractors. We also wish to acknowledge and thank the Yakama Nation Tribal Council, U.S. Bureau of Reclamation, Yakima Subbasin Fish and Wildlife Recovery Board, Pacific States Marine Fisheries Commission, U.S. Forest Service Naches Ranger District, University of Idaho, Lars Moberg and associates, University of Washington, and Central Washington University for their many contributions to this project including both recommendations and data services.

Regional Assessment of Supplementation Project (1992)

“Supplementation is the use of artificial propagation in an attempt to maintain or increase natural production while maintaining the long term fitness of the target population, and keeping the ecological and genetic impacts on nontarget populations within specified limits”.

Given this mouthful and the high priority placed on this project from the beginning to research and evaluate hatchery effects/uncertainties, a LOT of work went into design and implementation of a comprehensive monitoring program. This talk is a “whirlwind tour” of what we’ve learned so far.



Some background on the Cle Elum spring chinook program in the upper Yakima. The project was designed to implement the “gravel-to-gravel” hatchery reform concept where a central facility is used to rear fish, but fish are released from acclimation sites near spawning and rearing areas so that returning fish can return to these areas, spawn naturally, and “integrate” with returning natural-origin fish. The 1st brood cycle began in 1997 with brood collection. The 1st age-4 returns spawning in the wild returned in 2001. The 1st generation age-4 returns from integrated (HoR and NoR) spawners returned in 2005, the 2nd generation returns in 2009, etc. Only NoR fish (unmarked and untagged fish) are used for brood stock.

The Naches River is being used as a control stream. Both the upper Yakima and Naches systems experience very similar environmental conditions, e.g., droughts and floods rarely if ever occur in one stream without impacting the other as well. Also, historical data suggest there are virtually no upper Yakima fish which stray into the Naches system. Thus, differences in these two populations over time can be attributed to supplementation.

Brood Years 1997-2001
Optimum Conv. vs Seminatural Rearing
Fast et al. 2008



“We found insufficient evidence to conclude that seminatural treatment resulted in higher survival indices than did optimum conventional treatment”.


This study compared a traditional hatchery raceway (optimum conventional treatment-OCT) with a semi-natural treatment (SNT; i.e., rearing in camouflage-painted raceways with surface and underwater structures and underwater feeders) designed to induce more natural coloration and behavior in juvenile fish.

Brood Years 1997-2001
Wild vs Integrated Hatchery Traits
Knudsen et al. 2006, 2008




Reported differences in lengths, weights, spawn timing, female egg mass, and fecundity suggesting likely fitness reductions in integrated HOR fish.

Brood Years 2002-2004
 Growth Modulation Studies
 Larsen et al. 2006, 2013




Release Size
 HI: 104.1mm LO: 95.7mm
 <SAR indices to Bonneville Dam

Age-at-Return Distribution
 Overall Biomass:
 HI about 21% > LO



This study used a more natural, or slower growth regime (LO) and compared it to a traditional hatchery feeding regime (HI). As expected, the HI feeding regime produced larger fish at release which outperformed smaller (LO) fish in terms of smolt-to-adult return (SAR) indices. However, science has only recently started to look more closely at age-at-return; in every case the age-at-return distribution was older (which equates to larger returning adults) for the LO treatment relative to the HI growth treatment.


Brood Year 2002-2014 Update Bosch et al. 2022, in review



This is an opportunistic study that investigated demographic, environmental condition, and survival relationships for PIT-tagged fish detected exiting the acclimation sites over 13 brood years using fork length at PIT-tagging as a surrogate for smolt release size. These graphs are pooled data for 3068 PIT-tagged fish released as juveniles from CESRF and detected at Bonneville Dam as returning fish over 13 brood years. The upper two slides depict the main findings in Bosch et al. 2022 (submitted to *Environmental Biology of Fishes* and in peer review) – larger fish at release exited acclimation sites earlier and returned at younger ages than did smaller fish at release. The lower panels depict temporal and origin trends in the data that warrant further investigation as to potential causes, e.g., changes in feeding/growth rearing regime, changes in feed nutrient content, changes in rearing water temperature, or possibly changes in wild/natural fish used for brood stock, i.e., a genetic change that might be causing younger age-at-return (see Waters et al. 2021).


Brood Years 2002-2010
Homing and Spatial Distribution
Dittman et al. 2010

“Final spawning location depended strongly on where fish were released as smolts, but many fish also spawned in the vicinity of the central rearing hatchery”.




A Synthesis of Findings after 3 generations of spawning in the natural environment Fast et al. 2015

Upper Yakima vs Naches Redds, 1981-2021



Upper Yakima vs Naches Natural-Origin Returns, 1982-2021



The observed increase in redd abundance in the supplemented Upper Yakima from the pre- to post-supplementation period ($p=0.06$) is more than 4 times greater than that observed in the unsupplemented Naches population ($p=0.49$). Note that only 10 reds were observed in the Naches system in 2019. Changes in natural-origin returns were not significant in either case, but note that the downward trend in the Naches population post-supplementation is about 3 times greater than that observed in the Upper Yakima population.



Other Ecological Risks

- Proportions of mini-jacks and jacks concerning
- Ecological interactions within adopted guidelines
- Stray rates < 5%
- Pathogen and BKD risk profiles very low

References: Fritts and Pearsons 2004, 2006, 2008; Fritts et al. 2007; Galbreath et al. 2021; Larsen et al. 2004, 2010, 2019; Pearsons and Hopley 1999; Pearsons 2008; Pearsons et al. 2007, 2009; Pearsons and Temple 2007, 2010; Pierce et al. 2021; Temple and Pearsons 2012; Temple et al. 2017.

Relative Reproductive Success in an artificial spawning channel, brood years 2001-2005
Schroder et al. 2008, 2010, 2012




Females:
comparable egg
deposition rates, but
better survival to fry
for W/N

Males: comparable
breeding success
but W/N larger, more
dominant

HONOR. PROTECT. RESTORE.

This spawning channel was constructed for the specific purpose of studying relative reproductive success. From 2001 to 2005, we DNA-sampled and placed returning wild and hatchery-origin adults into the channel, let them spawn on their own, then collected DNA from a sub-sample of juvenile offspring and conducted a parent/progeny pedigree analysis to determine the number of progeny sired by hatchery- and wild-origin parents. Results were published in 2008 and 2010.



Since 2007 we have also been taking DNA samples from every spring Chinook returning through the Roza Dam adult monitoring facility for the purpose of doing a parent/progeny pedigree analysis for all fish destined to spawn in the Upper Yakima River over several years. In the illustration the circles represent the brood year with red lines corresponding to the year that aged 3 through 5 adults return to Roza Dam and are sampled prior to being released to spawn naturally upstream. This example shows fish from BY 2007, which return to spawn alongside their natural-origin counterparts in 2010 (age 3, “jacks”), 2011 (age 4), and 2012 (age 5). Mating among hatchery-reared and natural-origin fish occurred in every subsequent year to create wild-born F₂s, which return 3 to 5 years later. The example follows fish (returning to Roza in 2007) that spawned in the wild and whose progeny returned as adults in year 2010-12, and produced naturally-spawned fish (F₂s) that returned in years 2015 through 2017.

2007-2011 Parent/Progeny Preliminary Results Koch et al. 2022



These are the results of the Roza / whole river pedigree evaluation for the first brood cycle (brood year 2007-2011 fish returning to spawn in 2010-2016). The study is expected to be completed in the next two years with updated data from fish returning through 2021 to include grandparentage analysis.






Figure 4 from paper updated to include fourth generation (BY2014) fish. Density plot of individuals from the wild founders (P_1 Founders, black) and three generations of the integrated (INT, blue colors) and segregated (SEG, red colors) hatchery lines along the first discriminant function from the discriminant analysis of principal components (DAPC). Population genetic differentiation, F_{ST} , for each of the generations compared to the P_1 founders was low in all pairwise comparisons, ranging from 0 to 0.0108 (Table S3). The F_1 hatchery fish and segregated line steadily diverged from the P_1 founders over time; the F_2 and F_3 SEG populations were significantly differentiated from the founders ($F_{ST} = 0.0049$ and 0.0108 respectively, $P < 0.001$). In contrast, the integrated line did not exhibit the same temporal trend of increasing genetic divergence.



Supplementation has resulted in an increasing long-term trend in overall spring Chinook returns to the Yakima River Basin; however, the long-term trend in natural-origin returns is virtually flat.



“Salmonid management based largely on hatchery production, with no overt and large-scale ecosystem level recovery program, is doomed to failure”.

Meffe, GK. 1992. *Techno-Arrogance and Halfway Technologies...*

Conservation Biology 6:351-352

Conservation and fisheries professionals have made statements to regional and national policy makers very similar to this for over 150 years as European immigrants proceeded to occupy and transform the Columbia Basin to better serve their worldview which prioritized material wealth over ecosystem health. As Lisa Wilson of the Lummi Nation stated in a recent Op-ed, “re-establishing properly functioning ecosystems that support natural salmon production to meet these needs is preferable [to hatchery production] from a tribal perspective”. That is why we are working hard throughout the Columbia Basin to restore habitats. In the Yakima Basin...

Yakima Basin Integrated Plan




- Fish Passage Improvements
- Instream Flow Enhancement
- Floodplain Enhancement
- Habitat Restoration Actions
- For Example:
 - Remove Bateman causeway
 - Lower Yakima habitat focus
 - Cold water refugia
 - Predator mgmt./removal
 - Isolate and fix passage issues
 - Lake Cle Elum juvenile passage
 - Basinwide habitat work
 - Infrastructure upgrades




HONOR. PROTECT. RESTORE.

A lot of investment in terms of human and economic resources is occurring through the Yakima Basin Integrated Plan, a collaborative effort of tribal, city, county, irrigation, non-governmental, and other agencies and interests. But is this work sufficient to qualify as a “large-scale ecosystem level recovery program”? Or are there other factors still needing improvement...



And it is not just large barriers that are sources of mortality for juvenile and adult salmon. In the Yakima R. Basin there are still over 30 water use diversions to serve irrigation districts, source: BOR Yakima River Basin Study 2011, pp. 5-7.
<https://www.usbr.gov/pn/programs/yrbwep/reports/tm/6modreliabtyflow.pdf>



Managed rivers and climate change are reducing flows and increasing temperatures during critical juvenile and adult migration periods.

Predation/Non-natives (ISAB 2008)



Smallmouth bass




Channel Catfish



Walleye


HONOR. PROTECT. RESTORE.

Highly altered river systems have created conditions that favor species such as these and ...



Hinrichsen et al. (2013) reported that dam construction and alterations to the temperature and discharge regimes of the Columbia River have likely contributed to the increase in abundance and spatial distribution of American Shad *Alosa sapidissima* and Haskell et al. (2013) reported that this is altering food webs potentially affecting Pacific salmon.

Human Population Growth (Lackey 2000, 2001)



US and Canada censuses. State and regional district projections for 2010 and 2020

Since 2000:

Yakima County
+11%

WA State
+18%

I think we're all generally familiar with these human population trends (note OR and WA are reversed). More humans = more pressure to serve the needs and economic interests of those humans which historically has resulted in further alteration or elimination of habitats available to fish and wildlife.

Conclusions and Recommendations



- Expectations need to be consistent with reality
- Hatcheries aren't the cause of poor productivity
- Adaptive Management is important
 - Explore adjustments to rearing and release strategies to mitigate trend towards younger age-at-return
 - Growth modulation
 - Photoperiod and temperature manipulation
 - PRAS equipment



HONOR. PROTECT. RESTORE.

Western science: what can we learn about organisms; traditional ecological knowledge: what can we learn from organisms. Both ways of learning are communicating the same message: if we want thriving, wild salmon populations we need healthy, wild ecosystems. If we are unable or unwilling to “re-wild” the ecosystems, we cannot and should not expect hatcheries to replace the wild productivity we willingly sacrifice to serve the needs of an ever-growing human economy.

Until we transition from a society that asks, “How can we continue to manage resources to maximize benefits to humans and the economy while minimizing risks to fish and wildlife” to a society that asks, “How can we restore healthy and whole ecosystems while minimizing sacrifices to humans and the economy” we will need hatchery programs to provide some fishery benefits and meet minimal obligations. Therefore we need to be looking at modifying hatchery operations in all ways possible to optimize their performance in terms of returning adults to fisheries and the spawning grounds.

Yakima Basin Levi George / Cle Elum Supplementation and Research
AFS 2022 Presentation and Project-related References

- Beckman, B.R. and D.A. Larsen. 2005. Upstream Migration of Minijack (Age-2) Chinook Salmon in the Columbia River: Behavior, Abundance, Distribution, and Origin. *Transactions of the American Fisheries Society* 134:1520-1541.
- Beckman, B.R., B. Gadberry, P. Parkins, and D.A. Larsen. 2008. The Effect of Yakima River Spring Chinook Salmon Sire Life History Type on Emergence Timing and Size of Progeny. *Transactions of the American Fisheries Society* 137:1285-1291.
- Blodgett, B., M. Johnston, and B. Bosch (editors). 2021. Yakima/Klickitat Fisheries Project Monitoring and Evaluation – Yakima Subbasin; Final Report for the performance period May 1, 2020 through April 30, 2021, Project No. 1995-063-25, 388 electronic pages. <https://www.cbfish.org/Document.mvc/Viewer/P186823>.
- Bosch, W.J., S.N. Pandit, B.P. Sandford, G.M. Temple, M.V. Johnston, and D.A. Larsen. 2022. Effects of volitional emigration timing and smolt size on survival and age-at-return in a Pacific Salmon hatchery population. *Environmental Biology of Fishes* (in peer review).
- BPA (Bonneville Power Administration). 1996. Yakima Fisheries Project. Final Environmental Impact Statement. Bonneville Power Administration. Washington Department of Fish and Wildlife. Yakama Indian Nation. January, 1996. DOE/EIS-0169. DOE/BP-2784. Portland, OR.
- Busack, C., T. Pearsons, C. Knudsen, S. Phelps, Washington Department of Fish and Wildlife, B. Watson, M. Johnston, Yakama Nation, U.S. Department of Energy, Bonneville Power Administration, Division of Fish and Wildlife. 1997. Yakima Fisheries Project spring Chinook supplementation monitoring plan. Project Number 195-065, Contract Number DE-BI79-1996 BPA64878. <https://pisces.bpa.gov/release/documents/documentviewer.aspx?doc=64878-1>
- Busack, C., S. L. Schroder, C. M. Knudsen, T. N. Pearsons, and A. L. Fritts. 2006. Natural Production and Domestication Monitoring of the Yakima Spring Chinook Supplementation Program: December 2005 Revision. Chapter 7 in Busack et al., "Yakima/Klickitat Fisheries Project Genetic Studies; Yakima/Klickitat Fisheries Project Monitoring and Evaluation", 2005-2006 Annual Report, Project No. 199506325, 205 electronic pages, ([BPA Report DOE/BP-00022370-5](#)).
- Busack, C. and C.M. Knudsen. 2007. Using factorial mating designs to increase the effective number of breeders in fish hatcheries. *Aquaculture* 273:24-32.
- Busack, C., C.M. Knudsen, G. Hart, and P. Huffman. 2007. Morphological Differences Between Adult Wild and First-Generation Hatchery Upper Yakima River Spring Chinook Salmon. *Transactions of the American Fisheries Society* 136:1076-1087.
- Cuenca, M.L., T.W.H. Backman, and P.R. Mundy. 1993. The use of supplementation to aid in natural stock restoration. Pages 269-293 in J.G. Cloud and G.H. Thorgaard, editors. *Genetic conservation of salmonid fishes*. Plenum Press, New York.
- Dittman, A.H., D. May, D.A. Larsen, M.L. Moser, M. Johnston, and D. Fast. 2010. Homing and spawning site selection by supplemented hatchery- and natural-origin Yakima River spring Chinook salmon. *Transactions of the American Fisheries Society* 139:1014-1028.
- Fast, D.E., D. Neeley, D.T. Lind, M.V. Johnston, C.R. Strom, W.J. Bosch, C.M. Knudsen, S.L. Schroder, and B.D. Watson. 2008. Survival Comparison of Spring Chinook Salmon Reared in a

- Production Hatchery under Optimum Conventional and Seminatural Conditions. Transactions of the American Fisheries Society 137:1507–1518.
- Fast, D.E., W.J. Bosch, M.V. Johnston, C.R. Strom, C.M. Knudsen, A.L. Fritts, G.M. Temple, T.N. Pearsons, D.A. Larsen, A.H. Dittman, and D. May. 2015. A Synthesis of Findings from an Integrated Hatchery Program after Three Generations of Spawning in the Natural Environment. North American Journal of Aquaculture 77:377-395.
- Fritts, A.L., and T.N. Pearsons. 2004. Smallmouth bass predation on hatchery and wild salmonids in the Yakima River, Washington. Transactions of the American Fisheries Society 133:880-895.
- Fritts, A.L. and T.N. Pearsons. 2006. Effects of Predation by Nonnative Smallmouth Bass on Native Salmonid Prey: the Role of Predator and Prey Size. Transactions of the American Fisheries Society 135:853-860.
- Fritts, A.L., J.L. Scott, and T.N. Pearsons. 2007. The effects of domestication on the relative vulnerability of hatchery and wild spring Chinook salmon to predation. Canadian Journal of Fisheries and Aquatic Sciences 64:813-818.
- Fritts, A.L., and T.N. Pearsons. 2008. Can nonnative smallmouth bass, *Micropterus dolomieu*, be swamped by hatchery fish releases to increase juvenile Chinook salmon, *Oncorhynchus tshawytscha*, survival? Environmental Biology of Fishes 83:485–494.
- Galbreath, P.F., B.A. Staton, H.M. Nuetzel, C.A. Stockton, C.M. Knudsen, L.R. Medeiros, I.J. Koch, W.J. Bosch, and A.L. Pierce. 2021. Precocious Maturation of Hatchery-Raised Spring Chinook Salmon as Age-2 Minijacks Is Not Detectably Affected by Sire Age. Transactions of the American Fisheries Society <https://doi.org/10.1002/tafs.10343>.
- Ham, K.D., and T.N. Pearsons. 2001. A practical approach for containing ecological risks associated with fish stocking programs. Fisheries 25(4):15-23.
- Haskell, C.A., K.F. Tiffan, and D.W. Rondorf. 2013. The Effects of Juvenile American Shad Planktivory on Zooplankton Production in Columbia River Food Webs. Transactions of the American Fisheries Society, 142: 606-620.
- Hess, M. A., C. D. Rabe, J. L. Vogel, J. J. Stephenson, D. D. Nelson, and S. R. Narum. 2012. Supportive breeding boosts natural population abundance with minimal negative impacts on fitness of a wild population of Chinook Salmon. Molecular Ecology 21:5236–5250.
- Hinrichsen, R.A., D.J. Hasselman, C.C. Ebbesmeyer, and B.A. Shields. 2013. The Role of Impoundments, Temperature, and Discharge on Colonization of the Columbia River Basin, USA, by Nonindigenous American Shad. Transactions of the American Fisheries Society, 142: 887-900.
- Independent Scientific Advisory Board (ISAB). 2008. Non-native species impacts on native salmonids in the Columbia River Basin. ISAB Non-native Species Report ([ISAB 2008-4](#)). Northwest Power and Conservation Council. Portland, Oregon.
- Janowitz-Koch, I., C. Rabe, R. Kinzer, D. Nelson, M.A. Hess, and S.R. Narum. 2019. Long-term evaluation of fitness and demographic effects of a Chinook Salmon supplementation program. Evolutionary Applications 12:456–469.
- Knudsen, C.M., S.L. Schroder, C.A. Busack, M.V. Johnston, T.N. Pearsons, W.J. Bosch, and D.E. Fast. 2006. Comparison of Life History Traits between First-Generation Hatchery and Wild Upper Yakima River Spring Chinook Salmon. Transactions of the American Fisheries Society 135:1130–1144.

- Knudsen, C.M., S.L. Schroder, C. Busack, M.V. Johnston, T.N. Pearsons, and C.R. Strom. 2008. Comparison of Female Reproductive Traits and Progeny of First-Generation Hatchery and Wild Upper Yakima River Spring Chinook Salmon. *Transactions of the American Fisheries Society* 137:1433-1445.
- Knudsen, C.M., M.V. Johnston, S.L. Schroder, W.J. Bosch, D.E. Fast, and C.R. Strom. 2009. Effects of passive integrated transponder tags on smolt-to-adult recruit survival, growth, and behavior of hatchery spring Chinook salmon. *North American Journal of Fisheries Management* 29:658-669.
- Koch, I.J., T.R. Seamons, P.F. Galbreath, H.M. Nuetzel, A.P. Matala, K.I. Warheit, D.E. Fast, M.V. Johnston, C.R. Strom, S.R. Narum, and W.J. Bosch. 2022. Effects of Supplementation in Upper Yakima River Chinook Salmon. *Transactions of the American Fisheries Society* <https://doi.org/10.1002/tafs.10354>
- Lackey, R. T. 2000. Restoring wild salmon to the Pacific Northwest: chasing an illusion? Pages 91–143 in P. Koss and M. Katz, editors. *What we don't know about Pacific Northwest fish runs: an inquiry into decision making*. Portland State University, Portland, Oregon.
- Lackey, R. T. 2001. Defending reality. *Fisheries* 26(6):26–27.
- Larsen, D.A., B.R. Beckman, K.A. Cooper, D. Barrett, M. Johnston, P. Swanson, and W.W. Dickhoff. 2004. Assessment of High Rates of Precocious Male Maturation in a Spring Chinook Salmon Supplementation Hatchery Program. *Transactions of the American Fisheries Society* 133:98-120.
- Larsen, D.A., B.R. Beckman, C.R. Strom, P.J. Parkins, K.A. Cooper, D.E. Fast, and W.W. Dickhoff. 2006. Growth Modulation Alters the Incidence of Early Male Maturation and Physiological Development of Hatchery-Reared Spring Chinook Salmon: A Comparison with Wild Fish. *Transactions of the American Fisheries Society* 135:1017-1032.
- Larsen, D.A., B.R. Beckman, and K.A. Cooper. 2010. Examining the Conflict between Smolting and Precocious Male Maturation in Spring (Stream-Type) Chinook Salmon. *Transactions of the American Fisheries Society* 139:564-578.
- Larsen, D.A., D.L. Harstad, C.R. Strom, M.V. Johnston, C.M. Knudsen, D.E. Fast, T.N. Pearsons, and B.R. Beckman. 2013. Early Life History Variation in Hatchery- and Natural-Origin Spring Chinook Salmon in the Yakima River, Washington. *Transactions of the American Fisheries Society* 142:2, 540-555.
- Larsen, D.A., D.L. Harstad, A.E. Fuhrman, C.M. Knudsen, S.L. Schroder, W.J. Bosch, P.F. Galbreath, D.E. Fast, and B.R. Beckman. 2019. Maintaining a wild phenotype in a conservation hatchery program for Chinook salmon: The effect of managed breeding on early male maturation. *PLOS ONE* <https://doi.org/10.1371/journal.pone.0216168>.
- Major, W.W. III, J.M. Grassley, K.E. Ryding, C.E. Grue, T.N. Pearsons, D.A. Tipton, and A.E. Stephenson. 2005. Abundance and consumption of fish by California gulls and ring-billed gulls at water and fish management structures within the Yakima River, Washington. *Waterbirds* 28:366-377.
- Meffe, G.K. 1992. Techno-arrogance and halfway technologies: Salmon hatcheries on the Pacific coast of North America. *Conservation Biology* 6:351-52.
- Mobrand, L.E., J. Barr, L. Blankenship, D.E. Campton, T.T.P. Evelyn, T.A. Flagg, C.V.W. Mahnken, L.W. Seeb, P.R. Seidel, and W.W. Smoker. 2005. Hatchery Reform in Washington State: Principles and Emerging Issues. *Fisheries* 30:11-23.

- Nehlsen, W., J.E. Williams, and J.A. Lichatowich. 1991. Pacific Salmon at the Crossroads: Stocks at Risk from California, Oregon, Idaho, and Washington. *Fisheries* 16(2):4-21.
- Paquet, P. J., T. Flagg, A. Appleby, J. Barr, L. Blankenship, D. Campton, M. Delarm, T. Evelyn, D. Fast, J. Gislason, P. Kline, D. Maynard, L. Mobrand, G. Nandor, P. Seidel, and S. Smith. 2011. Hatcheries, Conservation, and Sustainable Fisheries—Achieving Multiple Goals: Results of the Hatchery Scientific Review Group's Columbia River Basin Review. *Fisheries* 36:11, 547-561.
- Pearsons, T.N., and C.W. Hopley. 1999. A practical approach for assessing ecological risks associated with fish stocking programs. *Fisheries* 24(9):16-23.
- Pearsons, T.N. 2002. Chronology of ecological interactions associated with the life-span of salmon supplementation programs. *Fisheries* 27(12):10-15.
- Pearsons, T.N. and G.M. Temple. 2007. Impacts of Early Stages of Salmon Supplementation and Reintroduction Programs on Three Trout Species. *North American Journal of Fisheries Management* 27:1-20.
- Pearsons, T.N., A.L. Fritts, and J.L. Scott. 2007. The effects of hatchery domestication on competitive dominance of juvenile spring Chinook salmon. *Canadian Journal of Fisheries and Aquatic Sciences* 64:803-812.
- Pearsons, T.N. 2008. Misconception, reality, and uncertainty about ecological interactions and risks between hatchery and wild salmonids. *Fisheries* 33:278-290.
- Pearsons, T.N., C.L. Johnson, B.B. James, and G.M. Temple. 2009. Abundance and Distribution of Precociously Mature Male Spring Chinook Salmon of Hatchery and Natural Origin in the Yakima River. *North American Journal of Fisheries Management* 29:778-790.
- Pearsons, T.N. and G.M. Temple. 2010. Changes to Rainbow Trout Abundance and Salmonid Biomass in a Washington Watershed as Related to Hatchery Salmon Supplementation. *Transactions of the American Fisheries Society* 139:502-520.
- Pearsons, T.N. 2010. Operating Hatcheries within an Ecosystem Context Using the Adaptive Stocking Concept. *Fisheries* 35:23-31.
- Pierce, A.L., L.R. Medeiros, B. Hoffman, I.J. Koch, S.R. Narum, P.F. Galbreath, and J.J. Nagler. 2021. Dietary tetradecylthioacetic acid supplementation during the fall prevents an increase in body lipid levels but does not influence precocious male maturation rate in juvenile spring Chinook salmon. *Aquaculture Research* 52:5483–5492. <https://doi.org/10.1111/are.15422>
- RASP (Regional Assessment of Supplementation Planning). 1992. Supplementation in the Columbia River Basin, Parts 1-5. Report DOE/[BP 01830-11](#), Bonneville Power Administration.
- Schroder, S.L., C.M. Knudsen, T.N. Pearsons, T.W. Kassler, S.F. Young, C.A. Busack, and D.E. Fast. 2008. Breeding Success of Wild and First-Generation Hatchery Female Spring Chinook Salmon Spawning in an Artificial Stream. *Transactions of the American Fisheries Society*, 137:1475-1489.
- Schroder, S.L., C.M. Knudsen, T.N. Pearsons, T.W. Kassler, S.F. Young, E.P. Beall, and D.E. Fast. 2010. Behavior and Breeding Success of Wild and First-Generation Hatchery Male Spring Chinook Salmon Spawning in an Artificial Stream. *Transactions of the American Fisheries Society*, 139:989-1003.
- Schroder, S.L., C.M. Knudsen, T.N. Pearsons, T.W. Kassler, E.P. Beall, S.F. Young, and D.E. Fast. 2012. Breeding Success of four male life history types of spring Chinook Salmon spawning in an artificial stream. *Environmental Biology of Fishes*, 94:231-248.

- Temple, G. M., and T. N. Pearsons. 2012. Risk management of non-target fish taxa in the Yakima River Watershed associated with hatchery salmon supplementation. *Environmental Biology of Fishes* 94:67-86.
- Temple, G.M., T. Newsome, T.D. Webster, and S.W. Coil. 2017. Evaluation of Rainbow Trout Abundance, Biomass, and Condition Following Coho Salmon Reintroduction in Taneum Creek, Washington. *Northwest Science* 91:54-68.
- U.S. Bureau of Reclamation. 2012. [Yakima River Basin Integrated Water Resource Management Plan](#), Final Programmatic Environmental Impact Statement. Pacific Northwest Region, Yakima, WA.
- Venditti, D.A., R.N. Kinzer, K.A. Apperson, B. Barnett, M. Belnap, T. Copeland, M.P. Corsi, and K. Tardy. 2018. Effects of hatchery supplementation on abundance and productivity of natural-origin Chinook salmon: two decades of evaluation and implications for conservation programs. [Canadian Journal of Fisheries and Aquatic Sciences](#) 75:1495-1510. See also: <https://nwcouncil.app.box.com/s/gsolcxk9nv1w3897am4th5nl7xe6108g>.
- Waters, C.D., J.J. Hard, M.S.O. Briec, D.E. Fast, K.I. Warheit, R. Waples, C.M. Knudsen, W.J. Bosch, and K.A. Naish. 2015. Effectiveness of managed gene flow in reducing genetic divergence associated with captive breeding. *Evolutionary Applications* 8:956-971. DOI: [10.1111/eva.12331](https://doi.org/10.1111/eva.12331).
- Waters, C.D., A. Clemento, T. Aykanat, J.C. Garza, K.A. Naish, S. Narum, and C.R. Primmer. 2021. Heterogeneous genetic basis of age at maturity in salmonid fishes. *Molecular Ecology* 30:1435-1456.
- Williamson, K., A. R. Murdoch, T. N. Pearsons, and E. J. Ward. 2010. Factors influencing the relative fitness of hatchery and wild spring Chinook salmon (*Oncorhynchus tshawytscha*) in the Wenatchee River, Washington, USA. *Canadian Journal of Fisheries and Aquatic Sciences* 67:1840-1851.