



Factors Affecting Recolonization of the Teanaway River by Spring Chinook Salmon

**Andrew Dittman
Darran May
Mary Moser
Anthony Fritts
Mark Johnston
Dave Fast**



Outline

- NOAA/UW/USGS Research
- Teanaway River Spring Chinook
- Factors affecting homing fidelity
- PIT tag analysis of Spring Chinook migratory behavior in
Teanaway River



Imprinting, Homing, and Straying

Homing research and the NWFSC mission




- Recovery science: Spatial structure and patterns of dispersal for population/ESU viability
- Ecological and genetic interactions between naturally-spawning (stray) hatchery and wild salmon
- Identification of Essential Fish Habitat for conservation and recolonization
- Anthropogenic impacts on homing/spawning success (e.g. contaminants, rural development, dams and transport, irrigation and water transport)

Monitoring –Redd mapping and analysis

(1981-2010)


- Weekly redd Surveys
- Float or walk reaches
 - 1-5 times during season
- Marked with flagging
- Georeferenced to reaches (2002-2010)




May et al. in prep.

Comprehensive carcass surveys of upper Yakima Basin (2002-2010)



- GPS location (3 m accuracy); date
- hatchery/wild
- male/female; jack, precocious
- length, scale
- tag location, recovery
- egg retention; disease
- dna (2003-2010; 3000 samples),
- otolith (2002-2010; ~1500 samples)
- ~15,000 carcasses sampled




Cle Elum R.




Teanaway R.




Yakima R.



Tradeoffs between homing and habitat quality (Cram et al., in review)



Spatial patterns of fine sediment infiltration (Klett et al., in prep.)




Outline

-NOAA/UW/USGS Research


-Teanaway River Spring Chinook

-Factors affecting homing fidelity


-PIT tag analysis of Spring Chinook migratory behavior in
Teanaway River




Utilization of the Teanaway River by Spring Chinook spawners




Utilization of the Teanaway River by Natural Origin spawners




Effects of supplementation on Teanaway River spawning




Effects of supplementation on Teanaway River spawning




Effects of supplementation on Teanaway River spawning




Effects of supplementation on Teanaway River spawning



Effects of supplementation on Teanaway River spawning



Effects of supplementation on Teanaway River spawning



Outline

-NOAA/UW/USGS Research


-Teanaway River Spring Chinook

-Factors affecting homing fidelity


-PIT tag analysis of Spring Chinook migratory behavior in
Teanaway River





Distribution of Jack Creek Fish 2010




Homing fidelity to the Teanaway R. is highly variable between years






Relationship between homing fidelity and Teanaway R. flow




Relationship between homing fidelity and Teanaway R. flow



Spawning site location (selection?) is size dependent

Jack Creek releases (female)



Outline

-NOAA/UW/USGS Research


-Teanaway River Spring Chinook

-Factors affecting homing fidelity


**-PIT tag analysis of Spring Chinook migratory behavior in
Teanaway River**




PIT tag analysis of Spring Chinook migratory behavior in Teanaway River




PIT Tag Recoveries – Carcass survey




Migratory patterns of Spring Chinook into Teanaway River




Gender specific migratory patterns of Spring Chinook into Teanaway River



PIT Tag Detections –Teanaway array



Size selective emigration of JC fish



Acknowledgements

Yakama Nation

Bill Bosch

Yakama biologists

WDFW

PIT Tag crew

Lynne Anderson

Field help

Most of NWFSC staff &
especially Physiology team

Funding