FLOODPLAIN FOREST REGENERATION AND FLUVIAL PROCESSES:

Status of Cottonwood Forest on the Kittitas Reach and Restoration Implications

Katrina Strathmann
Restoration Ecologist
Mid-Columbia Fisheries Enhancement Group
katrina@midcolumbiafisheries.org

Yakima Basin Science Conference June 14-15, 2023



OVERVIEW

- Overview of cottonwood regeneration
- Status of forests on the Kittitas reach
- Implications for restoration

Cottonwood is foundation species in arid west floodplains.

- Bank stability and slowed erosion
- Roughness, slowing flood velocities
- Shade, cooling water temperatures
- Input nutrients for aquatic insects
- Source of large wood
 - Create instream complexity
 - Facilitate channel movement
 - Aggrade sediments



Vegetative reproduction = <u>stand</u> <u>maintenance</u>

Sexual reproduction = <u>stand</u> <u>creation</u>

- Annually, seed released early summer
- Seed is viable for short period





Photo by Spencer Creek Nursery

Vegetative reproduction = <u>stand</u> maintenance

Sexual reproduction = <u>stand</u> <u>creation</u>

- Annually, seed released early summer
- Seed is viable for short period





Conditions required for cottonwood regeneration


- Floodplains connected to river
- Significant habitat-forming flows
 - Create bare substrate
- Spring freshet
 - Moist soil at the time of seed release
- Flow recession rate that does not exceed the rate of root growth

Stochastic events result in episodic regeneration

Less frequent floods, ice scour, drought

Cottonwood age distribution


- On unregulated rivers, seedlings dominate age classes
- > 80% of trees are <12 yrs old





Regeneration Status

Cottonwood age distribution


 33% of trees are <12 yrs old









Reach 6 – Downstream dynamic reach






ASSESSMENT: FOREST REGENERATION


Reach 2 – Confined






Forest Condition

- 27% of all trees are in moderate to extreme stress
- 10% of all trees in severe to extreme stress
- Condition varies by reach
 - Reaches 5, 6 contain fewest stressed trees



IMPLICATIONS FOR RESTORATION

Environmental Flows for Cottonwood Regeneration



IMPLICATIONS FOR RESTORATION

Incorporating Regeneration into Floodplain Restoration

Replanting:

 Depth to groundwater < 6 ft during summer high ET

Stand creation using regeneration conditions:

- Created seedbed surfaces → contouring
- Moist soil at seed release

 elevations/irrigation
- Propagule source

 assisted seeding
- Groundwater available during early root growth → elevations/irrigation




Managing Cottonwood Forests



Thank you to funders:

- SRFB
- Dept of Ecology





THANK YOU TO OUR PARTNERS

- Many, many private landowners
- Washington Recreation and Conservation Office - Salmon Recovery Funding
- Yakama Nation Dept of Natural Resources
- Natural Resources Design
- Kittitas County Public Works
- Washington Department of Fish and Wildlife
- City of Ellensburg
- Washington Dept. of Transportation
- Bureau of Reclamation
- Bureau of Land Management

MCF Field Technicians:
 Marissa Mei-Lien Louie
 Kyrsten Wolterstorff
 Tonya Morrey

